

ASSESSMENT OF SOIL FERTILITY STATUS IN MARORI BLOCK OF PILIBHIT DISTRICT, UTTAR PRADESH

ABSTRACT

This study aims to evaluate the agricultural productivity potential of soils in several villages in the Marori block of Pilibhit district during 2022. 40 soil samples were collected using a random sampling technique, air-dried, and analyzed for physical and chemical properties. The results showed that the soil content varied significantly, with sand content ranging from 8.2-78.4% silt content from 9.2-67.3%, and clay content from 4.2-39.2%. Bulk density, particle density, porosity, pH levels, electrical conductivity, and organic carbon were also measured. The available nutrients varied, with nitrogen, phosphorus, potassium, calcium, magnesium, and sulphur ranging from 149.13-265.63 kg/ha, 13.59-49.72 kg/ha, 99.34-300.53 kg/ha, 3.3-6.9 cmol(P+)/kg, 1.53-4.20 and 11.29-19.23 kg/ha, respectively. The results showed that 99.5% of the soil samples had normal pH levels, and 0.5% was acidic. The majority of the soil samples had low organic carbon, with 73% having the highest available P. The majority of the soil samples were adequate in terms of exchangeable Ca and Mg content, and 60% had the greatest available sulphur. In a nutshell, the study highlights the importance of soil fertility in sustainable agricultural production.

Key words: GIS, Marori block, Nutrient index, Soil fertility

INTRODUCTION

Soil is a vital natural dynamic body that sustains all living things on Earth (Jones, 2012). Soil fertility evaluation is the most fundamental decision-making tool for efficiently planning a specific land use system (Havlin *et al.*, 2010). There are various procedures for determining soil fertility status, but soil testing is the most widely used and most suited. Soil testing offers information about nutrient availability in soils, which is used to make fertilizer recommendations for crop productivity. Soil analysis comprises physical features (texture, structure, color, bulk density, and so on) as well as chemical properties (soil pH, organic matter, macro and micronutrients, and so on), which are required for long-term soil management (Panda, 2010). According to Wilding and Lin (2006), soil contains various

amounts of air, water, minerals, organic materials, untold numbers of creatures, and other elements that are necessary for life. The availability of nutrients to growing crop plants as well as the total amount of nutrients present affect the overall crop growth and development (Shreesanthe *et al.*, 2018). Therefore, it is important to regularly and systematically examine the sufficiency of major, secondary, and micronutrients as well as nutrient deficits. Therefore, the data from soil tests is the most accurate source of information on the availability of plant nutrients and how to modify fertilizer recommendations for different crops. Prior to the development of new technologies like Global Positioning Systems (GPS) and Geographic Information Systems (GIS), it was challenging to describe the spatial diversity of soil fertility throughout a field. For the purpose of creating thematic soil fertility maps, it is crucial to collect soil samples using GPS (Mishra *et al.*, 2016). Keeping the aboveaid matter in view, for improving productivity in rice-based systems, soil fertility was assessed using the following parameters: organic matter content (OC), soil pH, accessible sulphur, electrical conductivity (EC), total nitrogen (TN), C/N ratio, available phosphorus, exchangeable calcium (Ca), magnesium (Mg), and texture (sand, silt, and clay). For the aforementioned fertility indicators, they used average weighted data from topsoil samples taken at depths ranging from 0 to 15 cm (Delsouz, 2017). In this paper, the fertility rates of various villages in the Pilibhit area of Uttar Pradesh are discussed. It might be helpful in suggesting the best crops, cropping patterns, and methods for managing the soil for the regions sustained yields.

MATERIAL AND METHODS

Site Description:

The present investigation carried out in the Marori block which is situated between latitude 28°64' North and longitude 79°81' East (Table 1; Fig. 1). The existing crops in this block are rice, wheat, sugarcane, mustard, and seasonal vegetables etc. Pilibhit experiences winter from November to February. It experiences pleasant windy days, clear skies, and cool nights from November to the end of February. The day temperature hovers around 14 °C (57°F) while night temperature is below 7°C (45°F) for most of December and January, often dropping to 3°C (37°F) or 4°C (39°F). Rain is expected to be occurs in February with rainfall

distribution of 1300-1400mm. The climate of the area is sub humid with max temp. 38.4°C and min temp 4.3°C.




Fig 1:Representing study map of Pilibhit area




Fig 2:Representing sampling map of site area

Soil sampling and analysis

Altogether 40 surface soil samples (0-15cm depth) were collected prior to the sowing of crops during the month of January 2023 from Marori block covering different villages Araji Chant, Baldevpur, Bilgaon, ChidiaDih, Deshnagar, Devipura, Himmatnagar, Junapuri, Kaim, Lalpuriya SahibSingh, Mohanpur, Mundiaratanpuri (Fig. 2). The soil samples were collected in zigzag manner from the different locations of crop field & near about 0.5 kg collected samples from each field were stored in the polythene bag. The samples were collected along with the geo-coordinates, recorded from each location by using global positioning system (GPS) tool. A varied number of soil samples were gathered from each hamlet and placed in a polythene bag with appropriate labeling. While collecting the soil samples, the specifics of the farmer's field, crops cultivated in that area, and fertilizers applied were noted. Soil samples were dried at room temperature and pulverized with a wooden roller on a wooden plank before being sieved at 2 mm. Then, soil samples that had been homogenized were placed in polythene bags for further analysis to determine the physicochemical properties of the soil. The department of Ag chemistry and Soil Science's macronutrient lab examined soil samples in accordance with standard operating procedure.

Nutrient index evaluation:

The soils of the individual blocks were classified into three fertility classes based on nutrient index values obtained from soil test summaries and their percentage distribution into low, medium, and high categories. Parker et al. (1951) devised the nutritional index.

Nutrient Index

$$= \frac{[\% \text{ in high category} \times 3 + \% \text{ in medium category} \times 2 + \% \text{ in low category} \times 1]}{100}$$

In this percent assessment a nutrient index less than 1.5 denotes low category and that falls between 1.5 and 2.5 represents the medium fertility class. Value of 2.5 and above (maxi 3.00) signifies a high fertility class in respect of the particular nutrient (Ghosh and Hasan, 1976).

Statistical analysis:

Snedecor and Cochran's (1967) technique for doing simple statistical analysis was used. These analyses included maximum, minimum, mean, coefficient of variation and correlation. The Coefficient Variation was determined by using formula:

$$C.V. = \frac{\text{Standard Deviation}}{\text{Mean}} \times 100$$

The relationship between relevant soil properties and available cationic macronutrient of soils were calculated by using standard statistical methods. The correlation coefficient was determined by using the formula:

Where:

r = Correlation coefficient

SP (xy) = Sum product of x, y variables

SS (x) = Sum of square of x variable

SS (y) = Sum of square of y variable.

RESULTS AND DISCUSSION:

Physical properties

In the soils of Marori block, the sand, silt and clay content of samples ranged from 8.2-78.4, 9.2-67.3 and 4.2-39.2(Table 4) respectively. There is no trend in the study area (Khadka *et al.* (2019).The bulk density of the studied soils varied from 1.12-1.47 g/cm³. The lowest bulk density was found in samples S₃₀ and S₄₀ from the villages of Deshnagar and Junapuri; this may be due to the presence of high organic carbon content.The bulk density exhibited a significant and negative correlation (r = -0.810*) with organic carbon (Table 6).Similar result was observed by Khadka *et al.* (2019) in the Agricultural Research Station, Vijayanagar, Jumla, Nepal. A similar result was observed by Patel *et al.* (2017) in the Jaunpur District of eastern Uttar Pradesh.The average particle density of the studied soils was 2.25 g/cm³. Devipura, ChidiaDih, Junapuri and Divipura village samples S₇, S₂₃, S₃₃ and S₃₇ had the highest value of particle density, while Himmatnagar village samples S₂ had the lowest. The Particle density exhibited a significant and negative correlation (r = -0.811**) with organic carbon (Table 6). Deolin *et al.* (2020) found comparable outcomes in the Saheshpur block of Dehradun.The distributions of physical properties are presented in Fig. No.3.




Fig. 3: Spatial variability physical properties in Marori block of Pilibhit district, Uttar Pradesh.

Chemical Properties

The pH of Marori block ranged from 5.9-7.7 having mean value 7.08 (Table 5). The results indicate that 98% pH was normal (6.5-7.5) and 1% of samples were acidic (pH <6.5). Kumar *et al.* (2018) recorded similar results in different blocks of Rampur district Uttar Pradesh. A positive and significant correlation with porosity ($r = 0.346^*$) in the soil (Table 6) (Ramana *et al.* 2015; Kashiwaret *et al.* 2018). The electrical conductivity of the analyzed soil sample ranged from 0.03-0.37. The coefficient of variation and standard deviation were 0.55 and 56.65, respectively (Table 5). Bilgaon was found to have the highest electrical conductivity in S_1 , while Deshnagar had the lowest in S_{13} . Kumar *et al.* (2013) reported similar trends in the soil fertility status of some Muzaffarnagar, Uttar Pradesh. On the basis of the limits suggested by Muhar *et al.* (1963) for judging the salt problem of soils, most of the samples (99%) were found normal ($EC < 1.0$ dS/m) and the remaining 1% samples were found in the category of

soluble salt content critical for germination (EC 1 to 2 dS/m). The organic carbon content of the soils under investigation exhibited a range of 0.23-0.83% (Table 5). In the study conducted, it was seen that S₂ and S₃₁ Himmatnagar and Deshnagar villages exhibited the highest organic carbon. Conversely, S₃₉ situated in Junapuri village displayed relatively low organic carbon content. Furthermore, 50% of the soil samples analyzed exhibited low organic carbon levels, while 40% exhibited medium levels, and the remaining 10% exhibited high organic carbon content. The organic carbon exhibited a significant and negative correlation ($r = -0.810^{**}$), ($r = -0.811^{**}$) with bulk density and particle density (Table 6). Sonkar *et al.* (2023) documented similar occurrences in the Sakaldiha block situated inside the industrial region of Chandauli District in Uttar Pradesh. The distributions of chemical properties are presented in Fig. No.4.




Fig. 4: Spatial variability chemical properties in Marori block of Pilibhit district, Uttar Pradesh.

Primary and secondary nutrient status

The soil in the research region exhibited a range of available nitrogen content, spanning from 149.13-265.63 kg/ha with a mean value of 181.25 kg/ha (Table 5). Among the samples, sample number S₄₀ from the village Junapuri exhibited the greatest nitrogen content, while sample number S₉ from the village ArajiChant exhibited the lowest nitrogen content (Amara *et al.* 2017; Kumar *et al.* 2018). Out of 40 soil samples, 82.48% of samples had less than 280 kg/ha, 17.52% of samples had between 280 and 560 kg/ha, and no sample had more than 560 kg/ha based on the classification proposed by Subbiah and Asija (1956) (Devastu *et al.* 2017). Nitrogen has a significant and positive correlation ($r = 0.472^{**}$) with organic carbon (Table 6) Bharteey *et al.* (2023). The phosphorus concentration in the soil of the research region exhibited a range of 13.59-49.72 (kg/ha), with a mean value of 30.81 P₂O₅ (kg/ha). In the study, it was seen that sample number S₂₂ from ChidiaDih village exhibited the highest phosphorus content of 49.72. Additionally, it was found that 73% of the soil samples had the highest P₂O₅ (kg/ha) level, while 27% of the soil samples exhibited a medium P₂O₅ (kg/ha) content. Based on the limit proposed by Muhr *et al.* (1963). The available potassium content in these soils ranged from 99.34-300.53 kg/ha, with a mean of 193.13 kg/ha. No. S₇ village Devipura has the maximum potassium content, while No. S₅ village Mohanpur has the lowest potassium content. 75% of 40 soil samples had medium potassium content, 25% had low potassium content, and no samples had high potassium content. Singh *et al.* (2016) reported a similar trend in the soil fertility status of Araji line block Varanasi district Uttar Pradesh. The available calcium ranged from 3.3-6.9 (Cmol(p+)/kg) in the soils of the Marori block of the Pilibhit district, with a mean value of 5.62 (Cmol(p+)/kg). Lalpuriya village had the highest calcium availability, while Kaim village had the lowest calcium availability. A similar result was also obtained by Sharma *et al.* (2013) in the vegetable-growing soil of the Varanasi district with a mean value of 1764 ppm of available calcium. The available magnesium ranged from 1.53-4.20 (Cmol(p+)/kg), Lalpuriya village sample No. S₄ had the highest Mg²⁺, while Kaim village sample no. S₁₇ had the lowest Mg²⁺. Total samples were found to be within the adequate range for Mg²⁺ availability. Similar result is also obtained by Sharma *et al.* (2013). The range of sulphur content in Marori block soils was between 11.29-28.22 kg/ha. Sample No. S₄ from the village of Lalpuriya had the highest available sulphur content, while sample no. S₅ from the village of Mohanpur had the lowest. Similar results were also found by Rai *et al.* (2018) in black soil of Varanasi district of eastern Uttar Pradesh. The geographical distribution of available primary and secondary nutrients in the study area is presented in Fig 5.




Fig 5: Spatial variability map of Available primary and secondary nutrient Marori in block of Pilibhit District, Uttar Pradesh

Nutrient index value of study area

The Nutrient index value (NIV) for available primary nutrients *i.e.* N, P, K, and S. Marori block of Pilibhit district were given below in Fig. 6. The nutrient index value for the soils of Marori block was low for the available nitrogen, Phosphorus, potassium and Sulphur. It was analyzed that NIV for N, P, K and S were 1.20, 0.8, 0.81 and 0.98 respectively, against the nutrient index value less than 1.5 for low, 1.5 to 2.5 for medium and greater than 2.5 for high (Singh *et al.* 2016).




Fig 6: Nutrient Index value of Marori block of Pilibhit district

CONCLUSION

It can be concluded the Marori block of Pilibhit district's soil can be classified as sandy loam, clay loam, and silt clay loam, with a normal pH. Out of 40 soil samples, 50% had low organic carbon, 40% medium, 10% high, and 100% low available nitrogen. 73% had high available P, 75% medium and 25% low available K. The soil contained exchangeable Ca and Mg was 100% sufficient. This could be attributed to Soil deterioration in the district as a result of bad agricultural practices, intensive farming, monoculture and excessive irrigation. Integrated nutrient management (INM), which rationally combines chemical fertilizers with biofertilizers and organic manures, should be put into place to address these problems of Pilibhit District which could be environmentally safe and economically viable. The advantages of biofertilizers and bio inoculants in crop production for increasing soil fertility and nutrient status should be made known to farmers. Soil mapping using GIS may generate an idea about the soil fertility state, and it could be very useful for fertilizer management for a specific area simply by applying previously analyzed soil data, leading to site-specific smart nutrient management. Camps, rallies, training sessions, and frontline demonstrations should be arranged to have a greater and more persistent impact on farmers regarding the benefits of soil testing.

REFERENCES

- Black, C.A. (1965). Methods of soil analyses. Black CA ed. Madison Wisconsin, USA, 1-2, 1572.
- Bouyoucos, G.J. (1962). Hydrometer Method Improved for Making Particle Size Analysis of Soils. *Agronomy Journal*, 54: 464-465.
- Cheng, K.L., Bray, R.H. (1951). Determination of calcium and magnesium in soil and plant material. *Soil science* 72(6): 449-458.
- Chesnin, L, Yien, C.H.(1950). Turbidimetric determination of available sulfur. *Proceeding of Soil Sci Am*, 149.
- Delsouz Khaki, B., Honarjoo, N., Davatgar, N., Jalalian, A., Torabi Golsefidi, H. (2017). Assessment of two soil fertility indexes to evaluate paddy fields for rice cultivation. *Sustainability*, 9(8): 1299.
- Deoli, B.K., Shefali, A., Madhuben, S., Anwar, S.N. (2020). Assessment of Soil Quality Using Physiochemical Parameter of Soil In Dehradun District of Uttarakhand, 1580-1590.
- Desavathu, R.N., Nadipena, A.R., Peddada, J.R. (2018). Assessment of soil fertility status in Paderu Mandal, Visakhapatnam district of Andhra Pradesh through Geospatial techniques. *The Egyptian Journal of Remote Sensing and Space Science* 21(1): 73-81.
- Hanway, J.J., Heidel, H. (1952). Soil analysis methods as used in Iowa state college soil testing laboratory. *Iowa agriculture* 57.
- Havlin, H.L., Beaton, J.D., Tisdale, S.L., Nelson, W.L.(2010). *Soil Fertility and Fertilizers: An Introduction to Nutrient Management*. 7th Edition, PHI Learning Private Limited, New Delhi. India, 516.
- Jackson, M.L., Miller, R.H., Forkiln, R.E. (1973). Soil chemical analysis Prentic-Hall of India Pvt.And Ltd. New Delhi: 2nd Indian Rep.
- Jones Jr, J.B. (2012). Plant nutrition and soil fertility manual. CRC press.
- Khadka, D., Lamichhane, S., Amgain, R., Joshi, S., Shree, P., Kamal, S. A. H., Ghimire N H. (2019). Soil fertility assessment and mapping spatial distribution of Agricultural Research Station, Bijayanagar, Jumla, Nepal. *Eurasian Journal of Soil Science* 8(3): 237-248.
- Kumar, R., Singh, M., Kumar, S., Singh, M., and Kumar, P. (2018). Estimation of Soil Fertility Status under Sugar Cane-Wheat Farming System in Different Blocks of Rampur District of Uttar Pradesh. *Journal of Krishi Vigyan* 6(2): 101-104.
- Kumar, R., Singh, M., Mishra, A.K., Singh, R., Tripathi, N.C (2013). Assessment of soil quality under maize-wheat cropping system of Milak block, district Rampur, Uttar Pradesh. *Journal Homepage URL*, 3(1), 76-80.
- KVK Pilibhit District.(2015). [http:// Krishi Vigyan Kendra, Pilibhit \(kv4.in\)](http://KrishiVigyanKendra.Pilibhit(kv4.in)).
- Mishra, A., Das, D., Saren, S., Dey, P. (2016). GPS and GIS based soil fertility maps of Nayagarh district, Odisha. *Annals of Plant and Soil research* 18(1), 23-28.

- Muhr, G.R., Datta, N.P., Shankara, S.N., Dever, E., Lacey, V.K. Dovahue, R.R. (1963). Soil Testing in Indian USAID Mission to India.
- Olsen, S.R., Cole, C.V., Watanable, F.S. Dean, L.A. (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate. USDA Circular 939.
- Panda, S.C. (2010). Soil Management and Organic Farming. Agrobios. Bharat Printing Press, Jodhpur, India. Soil fertility mapping in Dindur sub-watershed of Karna-taka for site specific recommendations. *Journal of the Indian Society of Soil Science*, **64**(4): 381-390.
- Parker, F.W., Nelson, W.L., Winters, E., Miles, I.E. (1951). The broad interpretation and application of soil test information. *Agronomy Journal*, **43**(3): 105-112.
- Patel, A., Verma, S., Singh, S.K., Singh, R.K. (2017). Soil fertility status of Jaunpur District in Eastern Uttar Pradesh. *Journal of Pharmacognosy and Phytochemistry*, **6**(6): 949-952.
- Prakash, S., Singh, A., Naresh, R.K., Pal, D., Kumar, A. (2019). Assessment of productivity and soil fertility of Saharanpur in irrigated agro-ecosystem of western Uttar Pradesh. *International Journal of Chemical Studies*, **7**(6): 2225-2227.
- Rai, A., Singh, S. (2018). Available nutrients status in black soils of Varanasi district of eastern part of Uttar Pradesh. *Journal of Applied and Natural Science*, **10**(4): 1238-1242.
- Sharma R.P., Yadava R.B., Lama T.D., Bahadur A., Singh K.P. (2013). Status of Secondary Nutrients vis-à-vis Soil Site-characteristics of Vegetable growing Soils of Varanasi. *Vegetable Science*, **40** (1), 65-68.
- Shreekanth, S., Anita, E.K., Rekha, M.V., Champa, B.V., Nagaraja, M.S. (2018). Secondary and Micronutrient Status in Soils of Grape Orchards of Vijayapura Taluka in Northern Karnataka, India. *International Journal of Current Microbiology and Applied Sciences*, **7**(5): 1393-1401.
- Singh, A., Srikanth, B.H., Kumari, K. (2021). Determining the black soldier fly larvae performance for plant-based food waste reduction and the effect on biomass yield. *Waste Management* **130**: 147-154.
- Singh, S.K., Dey, P., Sharma, P.K., Singh, Y.V., Latore, A.M., Singh, C.M., Varma, S.S. (2016). Primary and cationic micronutrient status of soils in few districts of eastern Uttar Pradesh. *Journal of the Indian Society of Soil Science*, **64**(4): 319-332.
- Snedecor, G.W., Cochran, W.G. (1967). *Statistical methods* 6th Edition Iowa State University press Ames. Iowa USA.
- Sonkar, V., Jha, S.K., Tiwari, S., Akarsh, A., Priya, S., Sharan, P., and Sharan, A. (2023). Characterization and explanation of the soil fertility state of Sakaldiha block in the Chandauli District of Uttar Pradesh's industrial area. *World Journal of Advanced Research and Reviews*, 2023, **18**(01): 063–072.
- Subbiah, B.W., Asija, G.L. (1956). A rapid procedure for the estimation of available micronutrient in soils. *Current Science*, **25**: 259-260.

Walkley, A., Black, I.A. (1934). An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. Soil science, **37**(1): 29-38.

Wilding, L.P., Lin, H. (2006). Advancing the frontiers of soil science towards a geoscience. Geoderma, **131**(3-4): 257-274.

Table 1: Location of soil samples collected in different villages of Marori block of Pilibhit district

S. No.	Village name	Longitude (DD)	Latitude (DD)	Cropping History	
				Previous year	Current year
S ₁	Bilgaon	28.6480	79.8330	Sugarcane	Sugarcane
S ₂	Himmatnagar	28.6626	79.8500	Sugarcane	Sugarcane
S ₃	Bilgaon	28.6527	79.8355	Sugarcane	Sugarcane
S ₄	Lalpuria			Paddy Mustard,	Sugarcane
	SahibSingh	28.6390	79.8455		
S ₅	Mohanpur	28.6381	79.8677	Wheat, Sugarcane	Sugarcane
S ₆	Devipura	28.6313	79.8589	Sugarcane	Wheat
S ₇	Devipura	28.6185	79.8586	Sugarcane	Sugarcane
S ₈	ChidiaDih	28.6230	79.8440	Sugarcane	Wheat
S ₉	Araji Chant	28.6580	79.8510	Sugarcane	Wheat
S ₁₀	Kaim	28.6370	79.8610	Sugarcane	Wheat
S ₁₁	Devipura	28.6260	79.8540	Sugarcane	Paddy
S ₁₂				Mustard,	Paddy
	Deshnagar	28.6320	79.8260	Sugarcane	
S ₁₃	Deshnagar	28.6360	79.8470	Sugarcane	Paddy
S ₁₄	Baldevpur	28.6620	79.8590	Paddy Wheat	Wheat
S ₁₅	Mohanpur	28.6430	79.8650	Paddy	Sugarcane
S ₁₆	Lalpuria			Paddy Mustard	Sugarcane
	SahibSingh	28.6460	79.8510		
S ₁₇	Kaim	28.6310	79.8660	Sugarcane	Sugarcane
S ₁₈				Mustard,	Paddy
	Himmatnagar	28.6660	79.8530	Sugarcane	
S ₁₉	Bilgaon	28.6470	79.8420	Paddy, Sugarcane	Paddy
S ₂₀	Bilgaon	28.6430	79.8330	Sugarcane	Sugarcane
S ₂₁	ChidiaDih	28.6298	79.8442	Paddy, Sugarcane	Sugarcane
S ₂₂	ChidiaDih	28.6199	79.8342	Paddy	Sugarcane
S ₂₃	ChidiaDih	28.6290	79.8340	Sugarcane	Sugarcane
S ₂₄	Araji Chant	28.6570	79.8440	Paddy Mustard	Sugarcane
S ₂₅	Kaim	28.6290	79.8770	Wheat Paddy	Sugarcane
S ₂₆	Kaim	28.6240	79.8730	Paddy Mustard	Sugarcane
S ₂₇	kaim	28.6360	79.8690	Sugarcane	Paddy
S ₂₈	Mundia			Sugarcane	Sugarcane
	Ratanpuri	28.6580	79.8730		
S ₂₉	Deshnagar	28.6370	79.8370	Sugarcane	Paddy
S ₃₀	Deshnagar	28.6410	79.8270	Sugarcane	Sugarcane
S ₃₁	Baldevpur	28.6560	79.8580	Sugarcane	Sugarcane
S ₃₂	Junapuri	28.6450	79.8580	Paddy Mustard	Wheat
S ₃₃	Junapuri	28.6490	79.8570	Sugarcane	Paddy
S ₃₄	Baldevpur	28.6520	79.8480	Sugarcane	Sugarcane

S ₃₅	Mohanpur	28.6440	79.8760	Paddy Mustard	Sugarcane
S ₃₆	Mundia ratanpuri	28.6620	79.8690	Sugarcane	Sugar cane
S ₃₇	Devipura	28.6230	79.8650	Sugarcane	Sugarcane
S ₃₈	Junapuri	28.6465	79.8713	Sugarcane	Paddy
S ₃₉	Junapuri	28.6519	79.8686	Sugarcane	Sugarcane
S ₄₀	Junapuri	28.6534	79.8745	Sugarcane	Wheat

Table 2: Procedure used for physico-chemical analysis of soil of different villages of Marori block of Pilibhit district

a. Physical properties		Method applied	Reference
1	Particle size analysis (%)	Hydrometer method	Bouyoucos (1962)
2	Bulk density (g/cm ³)	Pycnometer	Black <i>et al.</i> (1965)
3	Particle density (g/cm ³)	Pycnometer	Black <i>et al.</i> (1965)
b. Chemical properties			
1	Soil reaction (pH)	pH meter	Jackson (1973)
2	Electrical Conductivity (dS/m)	EC bridge	Jackson (1973)
3	Organic Carbon (%)	Wet oxidation method	Walkley and Black (1934)
c. Available Nutrient			
4	Available N (kg/ha)	Alkaline KMnO ₄ method	Subbiah and Asija (1956)
5	Available P (kg/ha)	Olsen's method	Olsen <i>et al.</i> (1954)
6	Available K (kg/ha)	Ammonium acetate method	Hanway and Heidal (1952)
7	Available S (kg/ha)	Calcium chloride method	Chesnin and Yien (1950)
8	Exchangeable Ca and Mg (Cmol(p ⁺)/kg)	EDTA Method	Cheng and Bray (1951)

Table 3: Characterization of soil test values for different nutrients

Nutrient	Rating of the soil test values		
	Low	Medium	High
E.C. (dS/m)	<0.8	0.8-2.5	>2.5
O.C. (%)	< 0.5	0.5-0.75	> 0.75
Available N (kg/ha)	<280	280-560	>560
Available P (kg/ha)	<10	10-25	>25
Available K (kg/ha)	<108	108-280	>280
Available S (mg/kg)	<10	10-20	>20
	Deficient	Sufficient	Insufficient
Ex. Ca (Cmol(p ⁺) kg ⁻¹)	<1.5	1.5-4.5	>4.5
Ex. Mg (Cmol(p ⁺) kg ⁻¹)	<1.5	1.5-4.5	>4.5

Table 4:Physical properties of soils of different villages in Marori block of Pilibhit district, Uttar Pradesh

S. No.	Sand (%)	Silt (%)	Clay (%)	PD (g/cm ³)	BD (g/cm ³)	Porosity (%)
S ₁	50.3	45.5	4.2	2.2	1.35	38.64
S ₂	14.2	60.5	25.3	1.98	1.17	40.91
S ₃	35.5	45.8	18.7	2.1	1.36	35.24
S ₄	33.9	41.7	24.4	2.35	1.34	42.98
S ₅	40.5	38.2	21.3	2.26	1.32	41.59
S ₆	23.2	67.3	9.5	2.28	1.36	40.35
S ₇	60.3	16.5	23.2	2.36	1.42	39.83
S ₈	54.1	20.6	25.3	2.22	1.43	35.59
S ₉	72.4	16.1	11.5	2.24	1.39	37.95
S ₁₀	22.3	47.6	30.1	2.02	1.18	41.58
S ₁₁	8.6	66.1	25.3	2.04	1.19	41.67
S ₁₂	36.4	25.5	38.1	2.24	1.33	40.63
S ₁₃	16.7	64.1	19.2	2.29	1.36	40.61
S ₁₄	64.4	25.2	10.4	2.25	1.34	40.44
S ₁₅	12.1	62.3	25.6	2.28	1.47	35.53
S ₁₆	24.3	58.6	17.1	2.31	1.43	38.10
S ₁₇	13.1	65.2	21.7	2.28	1.34	41.23
S ₁₈	74.4	16.2	9.4	2.04	1.13	44.61
S ₁₉	20.3	45.5	34.2	2.32	1.45	37.50
S ₂₀	35.1	38.3	26.6	2.25	1.38	38.67
S ₂₁	59.6	13.3	27.1	2.21	1.44	34.84
S ₂₂	78.4	9.2	12.4	2.33	1.39	40.34
S ₂₃	73.7	18.1	8.2	2.36	1.42	39.83
S ₂₄	27.2	56.5	16.3	2.31	1.38	40.26
S ₂₅	53.1	23.3	23.6	2.35	1.33	43.40
S ₂₆	72.3	12.4	15.3	2.19	1.39	36.53
S ₂₇	21.5	65.3	13.2	2.28	1.37	39.91
S ₂₈	27.7	38.2	34.1	2.24	1.4	37.50
S ₂₉	8.2	55.3	36.5	2.26	1.35	40.27
S ₃₀	35.3	31.6	33.1	2.04	1.12	45.10
S ₃₁	68.2	12.7	19.1	2.26	1.34	40.71
S ₃₂	24.4	41.1	34.5	2.28	1.25	45.18
S ₃₃	36.7	24.1	39.2	2.36	1.38	41.53
S ₃₄	64.3	23.6	12.1	2.34	1.42	39.32
S ₃₅	18.1	48.4	33.5	2.45	1.45	40.82
S ₃₆	37.5	51.2	11.3	2.33	1.42	39.06
S ₃₇	26.2	48.1	25.7	2.36	1.45	38.56
S ₃₈	28.4	42.2	29.4	2.27	1.39	38.77
S ₃₉	29.6	32.1	38.3	2.31	1.37	40.69
S ₄₀	58.1	27.5	14.4	2.02	1.14	43.56
Max.	78.40	67.30	39.20	2.45	1.47	45.18
Min.	8.20	9.20	4.20	1.98	1.12	34.84
Mean	39.02	38.53	22.46	2.25	1.35	39.99
SD	21.07	18.00	9.61	0.11	0.09	2.54
SE	3.33	2.85	1.52	0.02	0.01	0.40
CV	54.01	46.71	42.79	5.01	6.94	6.36

Table 5: Chemical and Nutrient properties of soils of different villages in Marori block of Pilibhit district, Uttar Pradesh

S. No.	pH	EC (dS/m)	OC (%)	N (kg/ha)	P (kg/ha)	K (kg/ha)	S (kg/ha)	Ca (cmol(p+)/kg)	Mg (cmol(p+)/kg)
S ₁	5.9	0.37	0.68	172.66	22.50	225.89	25.40	4.8	2.67
S ₂	7.0	0.17	0.83	201.45	31.95	139.65	16.93	6.7	3.7
S ₃	6.9	0.11	0.64	169.87	27.47	176.31	22.58	5.8	2.55
S ₄	7.5	0.11	0.53	161.76	27.63	103.50	28.22	6.9	4.2
S ₅	7.4	0.09	0.68	173.54	22.74	99.34	11.29	6.5	3.96
S ₆	7.1	0.06	0.53	161.32	13.59	250.87	22.58	5.4	2.7
S ₇	7.1	0.10	0.39	199.65	22.67	300.53	14.34	6.3	3.2
S ₈	6.7	0.23	0.48	155.44	24.43	210.32	18.54	5.8	2.75
S ₉	6.9	0.12	0.38	149.13	40.15	222.41	22.37	5.7	2.97
S ₁₀	7.5	0.13	0.80	210.67	36.63	180.45	14.11	6.2	2.42
S ₁₁	7.0	0.12	0.79	255.94	31.58	103.50	17.78	5.9	2.36
S ₁₂	6.9	0.09	0.68	179.61	27.41	199.67	12.56	5.3	2.21
S ₁₃	7.2	0.03	0.67	171.43	40.53	279.21	26.81	6.3	3.89
S ₁₄	6.8	0.08	0.68	175.32	31.67	240.56	20.89	5.7	2.32
S ₁₅	6.7	0.07	0.29	166.56	27.58	222.60	18.91	5.2	2.41
S ₁₆	6.9	0.08	0.38	156.31	31.94	198.32	15.81	5.0	1.97
S ₁₇	7.3	0.10	0.63	153.27	18.73	240.43	16.88	3.3	1.53
S ₁₈	7.5	0.08	0.76	250.87	36.14	140.31	26.81	5.3	2.25
S ₁₉	7.5	0.09	0.24	152.76	45.34	280.17	13.56	6.4	3.22
S ₂₀	7.2	0.07	0.53	153.14	31.32	252.65	25.12	5.8	2.87
S ₂₁	7.5	0.10	0.45	163.35	22.38	221.69	16.89	4.7	2.22
S ₂₂	7.3	0.08	0.42	211.47	49.72	231.38	18.63	6.5	3.23
S ₂₃	7.1	0.06	0.27	171.51	27.65	241.13	14.87	5.8	2.79
S ₂₄	7.0	0.10	0.42	169.48	40.50	226.54	27.10	4.3	2.22
S ₂₅	7.1	0.06	0.47	168.76	31.87	293.67	12.53	6.2	3.21
S ₂₆	7.3	0.12	0.56	164.56	27.78	132.83	23.14	6.5	3.1
S ₂₇	7.1	0.07	0.39	169.37	18.00	139.63	13.83	6.1	2.96
S ₂₈	7.2	0.10	0.54	182.54	27.43	156.45	21.25	5.7	2.59
S ₂₉	7.3	0.08	0.50	230.45	36.76	133.61	14.96	4.8	2.24
S ₃₀	7.1	0.10	0.83	200.89	31.50	210.66	19.34	5.4	2.21
S ₃₁	7.2	0.14	0.53	166.32	49.50	154.98	21.56	6.2	3.25
S ₃₂	7.7	0.08	0.45	162.38	27.65	123.54	15.52	5.2	2.2
S ₃₃	6.8	0.07	0.35	240.46	40.85	130.16	23.71	4	1.87
S ₃₄	6.6	0.08	0.36	177.12	22.35	220.63	21.17	6.3	3.4
S ₃₅	7.0	0.10	0.24	152.53	31.56	123.76	26.25	6.4	3
S ₃₆	6.9	0.10	0.35	156.56	40.22	130.50	14.39	5.6	2.43
S ₃₇	6.7	0.07	0.38	222.34	27.39	220.75	23.99	6.2	2.86
S ₃₈	7.2	0.05	0.23	149.76	22.76	160.68	20.32	5	2.2
S ₃₉	6.8	0.06	0.27	153.97	40.85	295.67	16.46	4.9	1.97
S ₄₀	7.4	0.06	0.78	265.63	23.67	110.21	11.85	4.7	2.22
Max.	7.70	0.37	0.83	265.63	49.72	300.53	28.22	6.90	4.20
Min.	5.90	0.03	0.23	149.13	13.59	99.34	11.29	3.30	1.53
Mean	7.08	0.10	0.51	181.25	30.81	193.13	19.23	5.62	2.71
SD	0.33	0.06	0.18	31.65	8.46	59.59	4.85	0.79	0.61
SE	0.05	0.01	0.03	5.00	1.34	9.42	0.77	0.12	0.10

CV	4.62	56.66	35.19	17.46	27.45	30.85	25.24	14.04	22.36
----	------	-------	-------	-------	-------	-------	-------	-------	-------

Table 6: Correlation between physico-chemical and nutrient properties of different villages of Marori block of Pilibhit district, Uttar Pradesh

	Sand	Silt	Clay	BD	PD	Porosity	pH	EC	OC	N	P	K	S	Ca	Mg
Sand	1	-.891**	-.525**	.038	.014	-.045	-.033	.126	-.016	.042	.109	.137	.061	.174	.168
Silt	-.891**	1	.080	-.035	-.045	.003	-.090	-.038	.080	-.077	-.208	-.114	.029	-.145	-.085
Clay	-.525**	.080	1	-.019	.054	.091	.241	-.205	-.115	.052	.149	-.087	-.188	-.110	-.209
BD	.038	-.035	-.019	1	.808**	-.719**	-.311	-.066	-.810**	-.589**	-.024	.369*	.127	.040	.121
PD	.014	-.045	.054	.808**	1	-.172	-.144	-.269	-.811**	-.485**	.062	.315*	.109	.034	.170
Porosity	-.045	.003	.091	-.719**	-.172	1	.346*	-.213	.392*	.423**	.117	-.240	-.086	-.031	-.010
pH	-.033	-.090	.241	-.311	-.144	.346*	1	-.506**	.129	.102	.112	-.258	-.233	.134	.100
EC	.126	-.038	-.205	-.066	-.269	-.213	-.506**	1	.280	-.072	-.097	-.079	.157	.004	.074
OC	-.016	.080	-.115	-.810**	-.811**	.392*	.129	.280	1	.472**	-.114	-.261	-.007	.033	.019
N	.042	-.077	.052	-.589**	-.485**	.423**	.102	-.072	.472**	1	.122	-.349*	-.078	-.080	-.168
P	.109	-.208	.149	-.024	.062	.117	.112	-.097	-.114	.122	1	.067	.125	.132	.064
K	.137	-.114	-.087	.369*	.315*	-.240	-.258	-.079	-.261	-.349*	.067	1	-.012	-.062	-.012
S	.061	.029	-.188	.127	.109	-.086	-.233	.157	-.007	-.078	.125	-.012	1	.027	.152
Ca	.174	-.145	-.110	.040	.034	-.031	.134	.004	.033	-.080	.132	-.062	.027	1	.848**
Mg	.168	-.085	-.209	.121	.170	-.010	.100	.074	.019	-.168	.064	-.012	.152	.848**	1

** . Correlation is significant at the 0.01 level (2-tailed).

* . Correlation is significant at the 0.05 level (2-tailed).