

# Zooplankton of Sevsak Stream (Elazığ-TURKIYE)

## ABSTRACT:

The zooplankton and its seasonal distribution of Sevsak Stream has been investigated in 2021. A total of 31 zooplankton species have been recorded in the stream. Among the zooplankton 19 species identified from Rotifer 9 species from Cladocera and 3 species from Copepoda. In every season zooplankton species were recorded in various number. Highest number of taxa were recorded in spring. In this season also an increase occurred in the individual numbers of species. *Keratella cochlearis*, *Polyarthra dolichoptera* and *Cyclops vicinus* were the most recorded species in the stream. As in the other aquatic habitat, rotifera group was determined as the most recorded species in terms of taxa and number of individuals.

**Key words:** *Zooplankton, Rotifera, Copepoda, Cladocera, Sevsak Stream*

## 1. INTRODUCTION

Rotifera, Cladocera and Copepoda are the largest groups of zooplankton, which are the main links of the food chain in the aquatic environment. Species diversity and abundance in unit volume provide information about the biological characteristics of reservoirs and ponds. Zooplanktonic organisms constitute the main food source of fish in freshwater sources and they constitute the main food of many pelagic-fed fish species and young periods of demersal-fed fish [1,2].

In addition, zooplankton species are also used to determine water quality, trophic status of the lake and water pollution. With the increasing importance of zooplanktonic organisms, researches on Rotifera, Cladocera and Copepoda, which constitute an important part of zooplankton in our country as well as in the world, have been started. Zooplankton of stagnant waters has great attention by the scientists in Turkey but researches on the streams are relatively few. By some of these researches, the rotifer fauna of Gümüldür Stream [3], diversity of Rotifera in Tigris River [4], the rotifer fauna and its seasonal variations of Fırat River [5], the rotifer fauna and its seasonal variations of Zıkkım Stream [6], the rotifer fauna of Asi River [7], the zooplankton fauna of some rivers in Mediterranean Region [8], the Copepoda and Cladocera fauna of Asi River [9], the Rotifera fauna of Euphrates River basin [10] zooplankton structure of Karaman Stream [11] zooplankton succession of the Asi River [12] were determined. In this research, we aim to identify the zooplankton fauna and its seasonal composition of Sevsak Stream, where no previous relevant data are available.

## 2. MATERIALS AND METHODS

Hazar Lake, which is an important water source, within the borders of Elazığ province, 22 km from the city center. It is a tectonic lake. Its average depth is 93 m, its maximum length is 20 km, It has a maximum width of 5.4 km and an altitude of 1248 m. The streams feeding the Hazar Lake are called "fur streams" together with Sevsak Stream, Zıkkım Stream, Kürk Stream, Behramaz Stream and Mogal Stream. Sevsak Stream is located in the north east of the lake. The stream born from the high hills is poured into the lake through a channel after passing the state

highway. Sevsak Stream is a small stream and can dry out completely in the hot months of some years [13,14,15]. Sampling stations in Sevsak Stream is given in Fig 1.



**Fig1. Sampling stations in Sevsak Stream**

Zooplankton samples were collected Hydro-Bios plankton net (25 cm in diameter, 100 cm in length, with a 55-µm mesh size) and immediately fixed with a 4% solution of formalin. Samples were taken from stagnant or slow-running, vegetation-rich areas of the stream basin. To identify zooplankton species, various resources were used [16-22].

Some physicochemical parameters, such as water temperature, electrical conductivity, pH, dissolved oxygen were measured on-site simultaneously with the sampling time.

### 3. RESULTS AND DISCUSSION

During the study 31 zooplankton species have been recorded in the stream. Among the zooplankton 19 species identified from Rotifera 9 species from Cladocera and 3 species from Copepoda. The list of the species and their seasonal distribution was given in Table 1.

As shown in the Table 1 in spring in every station the highest number of species were recorded. The highest number of taxa were recorded in the second station of spring period (18 species), the least taxa were recorded in the first station of the stream in winter. *K. cochlearis* and *P. dolichoptera* from Rotifera were determined in 10 samplings. *C. retikulata* from Cladocera and *C. vicinus* from Copepoda were recorded the dominant species. *C. lacustris* recorded only the last stations. *D. forcipatus*, *T. tetractis* recorded only in one sampling. In every season zooplankton species have been identified from the stream.

Rotifera was found as the dominant group in terms of number of species and individuals. The relative density of Rotifera was 61.3% followed by Cladocera 29% and Copepoda 9.7%.

Comment [U1]: was

Comment [U2]: in

Comment [U3]: founded

**Table 1. The seasonal distribution of zooplankton in Sevsak Stream**

	Seasons											
	Winter			Spring			Summer			Autumn		
	Stations											
	1	2	3	1	2	3	1	2	3	1	2	3
<b>ROTIFERA</b>												
<i>Asplanchna priodonta</i> Gosse, 1850		+		+	+	+		+	+		+	+
<i>Brachionus angularis</i> Gosse, 1851				+		+	+		+			+
<i>Brachionus calyciflorus</i> Pallas, 1766				+				+			+	+
<i>Colurella obtusa</i> (Gosse, 1886)												
<i>Dicranophorus forcipatus</i> (O.F.Müller, 1786)				+								+
<i>Euchlanis dilatata</i> Ehrenberg, 1832					+	+		+	+	+	+	+
<i>Kellicottia longispina</i> (Kellicott, 1879)								+	+			
<i>Keratella cochlearis</i> (Gosse, 1851)	+		+	+	+	+	+	+		+	+	+
<i>Keratella quadrata</i> (Müller, 1786)					+	+		+	+		+	
<i>Lecane bulla</i> (Gosse, 1886)				+	+							
<i>Lecane closterocerca</i> (Schmarda, 1859)	+							+				
<i>Lecane flexilis</i> (Gosse, 1886)		+			+			+				
<i>Lecane luna</i> (O.F.Müller, 1776)					+		+					
<i>Lecane lunaris</i> (Ehrenberg, 1832)				+		+						
<i>Lepadella ovalis</i> (O.F.Müller, 1786)				+	+							
<i>Notholca squamula</i> (O.F.Müller, 1786)				+	+			+			+	
<i>Polyarhtra dolichoptera</i> Idelson, 1925	+		+		+	+	+	+	+	+	+	+
<i>Synchaeta oblonga</i> Ehrenberg, 1831						+	+					
<i>Trichotria tetractis</i> (Ehrenberg, 1830)					+							
<b>CLADOCERA</b>												
<i>Bosmina longirostris</i> (O.F.Müller, 1785)		+	+		+		+					
<i>Chydorus sphaericus</i> (O.F. Müller, 1776)					+							+
<i>Ceriodaphnia reticulata</i> (Jurine, 1820)				+	+	+	+	+	+			+
<i>Coronetella rectangula</i> Sars, 1862					+		+					+
<i>Cornigerius lacustris</i> (Spandl 1923-1924)			+			+			+			+
<i>Daphnia longispina</i> O.F. Müller 1875				+	+	+		+				+
<i>Diaphanosoma lacustris</i> Korinek, 1981							+	+				
<i>Macrothrix laticornis</i> (Fischer, 1851)						+						+
<i>Pleuroxus aduncus</i> (Jurine, 1820)		+						+		+		
<b>COPEPODA</b>												
<i>Acanthopodiaptomus denticornis</i> (Wierzejski, 1887)		+	+	+							+	
<i>Cyclops vicinus</i> Ulanin, 1875	+		+	+	+				+		+	+
<i>Nitokra hibernica</i> (Brady, 1880)						+			+		+	
<b>TOTAL</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>11</b>	<b>18</b>	<b>13</b>	<b>9</b>	<b>14</b>	<b>10</b>	<b>6</b>	<b>10</b>	<b>11</b>

In the stream pH values were changed between 7.1-7.6, dissolved oxygen 6.9-2 mg/L, electrical conductivity, 298-345 µS/cm and temperature 5.5.-16 °C (Table 2).

**Table 2. Some physical and chemical parameter values of Sevsak Stream**

Parameters	Winter	Spring	Summer	Autumn
El. Conductivity (µS/cm)	315	298	345	302
pH	7.1	7.3	7.6	7.2
Dissolved oxygen (mg/L)	8.9	9.2	6.9	7.6
Temperature °C	5.5	9	16	12

In recent years, many studies [23-37] showed that rotifers are the dominant species in running waters. The reason is that rotifers are less affected than other groups by the deterioration of water quality and display better adaptation to these conditions [1]. The importance of rotifers

Comment [U4]: adapted

increases in comparison to cladocerans when the abundance of the latter is low [38]. Furthermore, rotifers are known as opportunistic species in extreme conditions [39]. Within the last decade, because of the degradation of water quality of many wetlands for a variety of reasons (pollution, eutrophication, etc., and the effect of global warming), rotifers have become dominant species in many aquatic habitats [40]. In line with that finding, this study identified that most of the zooplankton species belonged to phylum Rotifera.

Comment [U5]: the dominant, species

It was observed that seasonal changes are important factors effecting seasonal distribution of zooplanktonic groups in Sevsak Stream. It is thought that water temperature effects the distribution and amount of zooplankton in the stream. The fact that zooplankton is more abundant in spring and summer supports this data.

Comment [U6]: effect

There are also zooplankton researches carried out in previous years in other streams flowing into the Hazar Lake, such as the Sevsak Stream. In Kürk Stream *Keratella cochlearis* from Rotifera *Cyclops vicinus* from Copepoda and *Ceriodaphnia reticulata* from Cladocera were taken the first places according to frequency of occurrence [27]. In Behramaz Stream *Kellicottia longispina*, *K. cochlearis* and *C. vicinus* were recorded as dominant species [29]. In Hoşrük Stream [41] *Polyarthra dolichoptera* from rotifers has been detected in 10 months and is the most common species among all zooplankton species. *Cyclops vicinus* from Copepoda was the second dominant species observed for 9 months.

Comment [U7]: taken

Comment [U8]: at

A one-year comprehensive zooplankton survey was conducted in Lake Hazar. As a result of the survey, 52 zooplankton species were found in the lake. In Hazar Lake *Brachionus quadridentatus* *Keratella quadrata* and *Synchaeta verrucosa* has been most recorded Rotifera species. *Cyclops vicinus* from Copepoda was observed every month. *Alona rectangula*, *Cornigerus lacustris* and *Diaphanosoma lacustris* from Cladocera were the most abundant cladoceran species [42].

The common species in Hazar Lake and the streams flowing into the lake (Kürk, Behramaz Hoşrük Streams) *K. cochlearis*, *P. dolichoptera* and *C. vicinus* species [27,29,41]. These species have also been recorded in Sevsak Creek. Another common finding is that rotifers are the most common species of all zooplankton in these wetlands. In addition, in these studies, it was reported that the most abundant zooplankton species in terms of the number of species and the number of individuals were in spring and summer. This finding is consistent with the zooplankton distribution of the Sevsak Stream.

*Cornigerus* species are endemic to the Ponto-Caspian, with the exception of *Cornigerus lacustris*, which is endemic to freshwater Lake Hazar in the Euphrates basin [43]. This species was recorded only in the last station next to Hazar Lake. This species living habitat known as Hazar Lake. The fact that the species was found at the last station connected to the lake indicates that this species is not a species belonging to this stream. This species was not recorded in the researches carried out before in the Hoşrük Stream [41] Zikkım Stream [37] and Behramaz Stream [29] which spills into the lake. In Hazar Lake *C. lacustris* was found every month [42].

Comment [U9]: lives, habitat

The QB/T index shows the rate of the number of *Brachionus* to the number of *Trichocerca*. The Q index is evaluated in three groups for the lake's trophic state, that Q=1 means oligotrophy, Q = 1.0-2.0 means mesotrophy, and Q>2 means eutrophy. In this study, Sevsak Stream was determined (2 species of *Brachionus*, *B. angularis* and *B. calyciflorus* and 1 species of *Trichocerca*, *T. tetractis*)  $QB/T \ 2/1=2$ . According to this, the Sevsak Stream showed mesotrophic property.

## CONCLUSION

The zooplankton species found during the research are important as they are the first record for Sevsak Creek. Hazar Lake has got a blue flag. Sevsak Stream, which is one of the streams pouring into the lake, should be protected in order not to be affected by pollutants and not to pollute the lake.

## REFERENCES

1. Kolisko R, Plankton Rotifers Biology and Taxonomy. E. Schweizerbart'sche Verlagsbuchhandlung (Nageleu. Obermiller), 1974, Stuttgart,Germany.
2. Wetzel RG, Limnology. W. B. Saunders Company, Philadelphia,1975, 743 p.
3. Ustaoglu MR, Balık S, Aygen C, Özdemir D, Rotifera fauna of Gümlüdü Stream (Izmir). E.Ü. Su Ürünleri Fakültesi, Su Ürünleri Dergisi 1996,13 (1-2): 163-169, Turkish.
4. Bekleyen A, Gokot B, Varol M, Thirty-four new records and the diversity of the Rotifera in the Turkish part of the Tigris River watershed, with remarks on biogeographically interesting taxa. Sci Res Essays. 2011, 6(30):6270-6284. doi: 10.5897/SRE11.355
5. Saler (Emiroglu) S, Şen B, Şen D, The seasonal variations of rotifers of Kömürhan region of River Fırat. Su Ürünleri Sempozyumu, 2000, Sinop, 385-396.
6. Saler (Emiroglu) S, Şen, B. Rotifers of Zikkım Stream which flows into Hazar Lake and their seasonal variations. XI. Ulusal Su Ürünleri Sempozyumu, 2001,1: 261-271
7. Bozkurt A, Gökusu MZL, Sarihan E, Taşdemir M, Rotifera fauna of Asi River (Hatay, Türkiye). Ege University Journal of Fisheries & Aquatic Sciences 2002, 19, (1-2): 63-67
8. Bozkurt A, Preliminary observation on the zooplankton fauna of some rivers in Mediterranean Region. Turk J Aqua Life. 2004, 2(3):65-70, Turkish.
9. Gökusu MZL, Bozkurt A, Taşdemir M, Sarihan E, Copepoda and Cladocera (Crustacea) fauna of Asi River (Hatay, Türkiye). Ege J Fish Aqua Sci. 2005, 22(1-2):17-19, Turkish.
10. Akbulut (Emir) N, Yıldız K, The Rotifera fauna of Euphrates of River Basin (Turkey). Hacettepe Journal Biology and Chemistry, 2005, 34: 93-105.
11. Altındağ A, Buyurgan Ö, Kaya M, Özdemir E, Dirican S, A survey on some physico-chemical parameters and zooplankton structure in Karaman Stream, Antalya, Turkey. J Anim Vet Adv. 2009, 8(9):1710-1716.
12. Bozkurt A, Güven SE, Zooplankton succession of the Asi River (Hatay-Turkey). J FisheriesSciences.com. 2010, 4(4):337-353, Turkish.
13. Özmen H, Kulaççı F, Çukurovalı A, Doğru M. Concentrations of heavy metal and radioactivity in surface water and sediment of Hazar Lake (Elazığ, Turkey). Chemosphere, 2004, 55(3) : 401 - 408
14. KoçerMAT, Water quality and phytoplankton distribution in the open water of Lake Hazar, Fırat University, Institute of Science and Technology, PhD Thesis, 2008, 357p, Turkish.
15. Rashid RF, Saler S, Çoban MZ. Water Quality of a Tectonic Lake: Hazar Lake. Asian Journal of Fisheries and Aquatic Research, 2022, 16(5):35-44
16. Koste W, Radertiere Mitteleuropas. 2. Berlin, Germany: Tafelband, 1978.
17. Edmonson WT, Freshwater Biology. 2nd edition. New York, NY, USA: John Wiley and Sons Inc.; London: Chapman and Hall Limited, 1959.
18. Harding JP, Smith WA, A Key to the British Freshwater Cyclopoid and Calanoid Copepods. 2nd edition. Ambleside, UK: Freshwater Biological Association, 1974.
19. Nogrady T, Pourriot R, Notommatidae and Scariidae, Guides to the Identification of the Microinvertebrates of the Continental Waters of the World 8, Dumont, H.J.F. (ed.), SPB Academic Publishing, 1995, 3: 248.
20. Segers H, The Lecanidae (Monogononta). Ghent, Belgium: Ghent University, 1995.
21. Smet WH, Proalidae (Monogonanta), Guides to the Identification of the Microinvertebrates of the Continental Waters of the World 9 (Co. Ed: Dumont, H.J.F.), SPB Academic Publishing, 1996, 4, 102.
22. Smirnov NN, Cladocera: the Chydorinae and Syciinae (Chydoridae) of the World. Guides to the identification of the of the Continental Waters of the World, Vol. 11. Amsterdam, the Netherlands: SPB Academic Publishing, 1996.

23. Özbay H, Altındağ A. Zooplankton abundance in the River Kars, Northeast Turkey: Impact of environmental variables. *Afr. J. Biotechnol.*,2009,8(21):5814-5818.
24. Saler S, Zooplankton of Munzur River (Tunceli-Turkey). *J Anim Vet Adv.*, 2011,10(2):192-194.
25. Saler S, İpek A, N, Zooplankton composition of Tohma Stream (Malatya - Turkey). *Journal of Aquaculture Engineering and Fisheries Research*,2016, 2: 1, 30-35
26. Saler S, Eroğlu M, Haykır H, Zooplankton of Peri Stream (Tunceli-Türkiye). *e-Journal of New World Sciences Academy, Ecological Life Sciences*,2010,6(2):14-20,Turkish.
27. Saler S, İpek N, Arslan S, Zooplankton of Kürk Stream (Elazığ-Turkey). *Journal of Fisheries Sciences.com*,2011,5(3):219-255.
28. Saler S, Bulut H, Birici N, Tepe R, Alpaslan K, Zooplankton of Karasu (Erzincan). *Eğirdir Su Ürün Fak Derg.*, 2015, 11(1):10-16, Turkish.
29. Saler S, Çelik S, Yüce S, Zooplankton of Behramaz Stream (Elazığ). *Ecology 2017 11-13 May, Kayseri* . 2017,397 p,Turkish.
30. Güher H, The investigation of zooplanktonic organisms (Rotifera, Copepoda, Cladocera) of Meriç River (Turkey). *J Anim Vet Adv.*,2012,11(24):4673-4677.
31. Gaygusuz Ö, Dorak Z, Species composition and diversity of the zooplankton fauna of Darlık Stream (Istanbul-Turkey) and its tributaries. *J FisheriesSciences. Com*, 2013, 7(4):329-343. doi: 10.3153/jfsc.com.2013037
32. Saler S, Haykır H, Zooplankton composition of Pülümür Stream (Tunceli-Turkey). *J Anim Vet Adv.*,2011, 10(11):1401-1403.
33. İpek N, Saler S, Zooplankton of Görgüşan Stream and Geban Stream (Elazığ-Turkey). *J FisheriesSciences.com*, 2012, 6(2):155-163. Turkish.
34. İpek N, Saler S, Zooplankton community structure of Ohi Stream (Elazığ-Turkey). *J FisheriesSciences.com*, 2013. 7(1):83-88
35. Bulut H, Saler S, Zooplankton variation of Murat River (Elazığ-within the borders Palu district). *Turk J Agrie-Food Sci Tech.*, 2014, 2(1):13-17,Turkish.
36. Bulut H, Saler S, Effect of physicochemical parameters on zooplankton at a freshwater body of Euphrates Basin (Elazığ-Turkey). *Cellular and Molecular Biology*, 2019, 65 (1), 8-13.
37. Bulut H, SalerS, Zooplankton of Zikkım Stream (Elazığ). 6th Asia Pasific International Modern Sciences Congress,2021, 1:352-35
38. Wallace RL, Snell TW, Ricci C, Nogrady T, Rotifera biology,ecology and systematics. In: Segers H, Dumont HJ, editors.*Guides to the Identification of the Microinvertebrates of the Continental Waters of the World*. 2nd edition. Ghent, Belgium:Kenobi Productions; Leiden, the Netherlands: BackhuysAcademic Publishing BV.2006.
39. Gannon JE, Stremberger RS, Zooplankton (Especially Crustaceans and Rotifers) as indicators of water quality transactions of the American Microscopical Society , 1978, 97, 1: 16-35
40. Gürbüz P, Buyurgan Ö, Tekatlı Ç, Altındağ A, Species diversity and community structure of zooplankton in three different types of water body within the Sakarya River Basin, Turkey. *Turk J Zool.*,2017, 41: 848-859.
41. Saler S, Yüce S, Çelik, B., Bulut, H. Zooplankton of Hoşrük Stream (Elazığ-Türkiye). *Türk Tarım – Gıda Bilim ve Teknoloji Dergisi*, 2018,6(5): 607-612, Turkish.
- 42.Çelik B, Saler S, Karakaya G, Yüce S, Özbey N, Şeker T,Evaluation of zooplankton of HazarLake (Elazığ) by means of geographic information system.*Ecological Life Sciences*, 2022,17 (1), 24-35, Turkish.
43. Dumont H J, Vierstraete A, Akbulut N, Cornigerius lacustris of Lake Hazar, Turkey, a synonym of Cornigerius maeticus (Pengo) of the Ponto-Caspian (Cladocera: Onychopoda). *Plazi.org taxonomic treatments database*. Checklist dataset, 2019. <https://doi.org/10.11646/zootaxa.4619.1.9> accessed via GBIF.org on 2022-07-04