

## Original Research Article

# Influence of chitosan coating on shelf-life, biochemical properties and nutrient elements of carrot (*Daucus carota*) during postharvest storage

### ABSTRACT

Post-harvest loss of different vegetables in Bangladesh is widespread due to the lack of suitable technologies. A laboratory trial was conducted to assess the effect of chitosan coating on weight loss, shelf-life, biochemical qualities and mineral elements of carrots at postharvest storage. At room temperature ( $\approx 23\text{-}25^{\circ}\text{C}$ ), four treatments were selected with different chitosan solutions (0.00%, 0.10%, 0.20% and 0.30%). Matured carrots were collected at 2, 4, 6 and 8 days after postharvest storage (DAPS) for weight loss, while the shelf-life data were measured at 4 and 8 DAPS. On the other hand, carrots were collected at 0 (fresh) and 8 DAPS to determine total sugar, phenolic and mineral contents (Ca, Mg, P, S, Na, and K). Compared to control, the application of 0.30% chitosan coating preserved weight loss by 1.48% at 8 DAPS. The results also showed that applying 0.30% chitosan coating significantly ( $P \leq 0.05$ ) extended the shelf-life of carrots up to 35% at 8 DAPS compared to the control. Chitosan coatings also enhanced the loss of total sugars and contents of Mg in carrots during postharvest storage. However, total phenolic contents in carrots decreased significantly ( $P \leq 0.05$ ) during postharvest storage, and the application of chitosan was unable to protect them. The study concluded that chitosan coating with 0.30% solution might be used to prevent weight loss, extend shelf-life, and improve some nutritional qualities of carrots viz., Ca, Na, K, P and S.

*Keywords: Carrot, chitosan, mineral nutrients, storage life, sugar, total phenol*

### 1. INTRODUCTION

Carrot (*Daucus carota* L.) is a nutrient-rich root vegetable that belongs to the Apiaceae family. It is one of the widely produced vegetables in winter during rabi cropping season (September to mid-December) in Bangladesh and is consumed both as salad and cooked vegetables [1]. It contains a greater content of vitamins including carotene (precursor of vitamin A), thiamine and riboflavin, proteins, carbohydrates, fiber, and mineral nutrients such as potassium, sodium and high in sugar [2]. It is also rich in phenolic content, which represents the antioxidant compounds in carrots [3]. Due to its high nutritional value, the production and consumption of carrots are increasing gradually both locally and globally. According to the Bangladesh Bureau of Statistics, nationally the carrot was grown in 5085 acres of land with 19246 Metric tonnes of production in the year of 2018 to 2019, whereas the production was 16306 Metric tonnes in the year of 2016 to 2017 [1]. However, to keep up with the pace of the demand throughout the year and to ensure the nutritional status of carrot quality post-harvest storage is inevitable. A daily newspaper reported that “Thousands of tonnes of vegetables and fruits go to waste annually in Bangladesh due to a lack of

sufficient technologies and knowledge on post-harvest processes" [4]. The report showed that among the vegetables, postharvest losses in country bean 24.29%, cabbage 24.44% and tomato 27.64% and losses of vegetables at traders' level are much higher than at farmers' level.

Besides the degradation during handling, storage and transportation to distribution post; losses due to active metabolism, water transpiration, respiration and spoilage through microorganisms are highly considerable [5]. The losses of water shrink the vegetable and lose luster and consumer preferences. The microbial infection by several bacteria and fungus causes rotting, discoloration and reduces the market value to a greater extent [6]. Postharvest diseases can be accounted for more than 25% of total yield loss in developed countries and 50% or more in underdeveloped countries [7]. Methods including controlled atmosphere storage, low temperature and edible coating are available to extend the shelf life and maintain the quality of post-harvest vegetables. The consumer acceptability of the addible food coating has increased in recent decades and is used widely around the globe [8-11].

Chitosan is a widely used food coating that has gained popularity for prolonging the post-harvest shelf life of fruits and vegetables in recent decades [12- 16]. After cellulose, chitosan is the second most abundant polysaccharide found in nature which is a deacetylated derivative of chitin [17]. It is known to be biodegradable, easily dissolvable in organic acids, non-toxic, biocompatible, biofunctional, and have strong antifungal and antimicrobial properties [18-21]. Chitosan has also been known to reduce post-harvest water loss from fruits and vegetables by maintaining firmness [22-23]. Study results showed that chitosan coating with a 0.2% solution in tomato fruits extended the shelf-life at postharvest storage by decreasing the weight loss and decay incidence [24]. In vivo investigations have shown that chitosan treatment can delay the postharvest deterioration of fruits and vegetables [24-25]. Despite many beneficial effects of chitosan on the post-harvest quality of vegetables only a few studies have been conducted so far on the effect of chitosan on the post-harvest quality of vegetables and fruits in Bangladesh [24, 26-28]. Moreover, studies are very limited regarding the effect of chitosan on the post-harvest shelf life, biochemical properties and nutrient content of carrots in Bangladesh for its successful application as a vegetable coating. Therefore, this study was designed to assess the influence of chitosan application at postharvest storage of carrots in Bangladesh and determine its biochemical and nutritional properties.

## **2. MATERIAL AND METHODS**

### **2.1 Experimental Site**

The present investigation was conducted at the Laboratory of Plant Nutrition and Environmental Chemistry, Dept. of Agricultural Chemistry, Bangladesh Agricultural University (BAU), Mymensingh, Bangladesh, from February to June 2020. The main objective of this study was to find out the most suitable application dose of chitosan for postharvest storage of carrots in Bangladesh condition as well as to study its effect on major biochemical and nutritional properties of carrots.

### **2.2 Collection and Sorting of Carrots**

Twelve (12) kg of fully matured carrots were collected from the farmer's field of the Sadar Upazila of Mymensingh district. After that all the carrot samples were transported to the Laboratory of Plant Nutrition and Environmental Chemistry, Dept. of Agricultural Chemistry, BAU, Mymensingh, Bangladesh. Then sorting/ screening of carrots was done manually based on their shape, size, and colour. Decomposed and pest-infested carrots were discarded at this stage. Finally, almost similar shapes, sizes, and coloured carrots were chosen for the laboratory trial.

### **2.3 Treatments and Their Preparation**

Chitosan was obtained from Research-Lab Fine Chem Industries, Maharashtra, India (CAS No. 9012-76-4; Deacetylation >80%). Four treatments were selected with different chitosan solutions Control (no chitosan), 0.10%, 0.20% and 0.30% following the method [24] to prepare the treatment solutions.

### **2.4 Postharvest Application of Chitosan**

A total of 5-6 carrots were dipped for 30 seconds in each chitosan treatment, and the same number of carrots were likewise dipped in distilled water with a pH of 5.0 (control). At a temperature of 25 °C, all treated carrots were left to air dry for 1 hour. One group was considered a replicate with three replications. There were 12 (4x3) groups of carrots in this experiment. The treated and control carrots were placed in zip-lock bags to keep the relative humidity (RH) between 90 to 95% and then stored at room temperature (25 °C).

### **2.5 Data Recorded at Postharvest Storage**

Data on the shelf-life of carrots were recorded at 4 and 8 days after storage (DAS), while the weight loss data of carrots were measured at 2, 4, 6 and 8 days after post-harvest storage. For chemical analyses, two carrots from each replication were randomly selected at 8 DAS.

### **2.6 Measurement of Biochemical Quality of Carrots**

Two (2) carrots samples from each replication were collected at 8 DAS to determine total sugar and phenol contents. Total phenol estimation in carrots was carried out with the Folin-Ciocalteu reagent [29]. The concentration of phenols in carrots was calculated against the catechol standard curve and expressed as mg phenols/100 g material.

Total sugar content in carrot is estimated by determining the volume of the unknown sugar solution required to reduce a measured volume of Fehling's solution [29]. Representative carrot slice was extracted using methanol solution at 1:10 ratio and distilled water was used to make the final volume of 50 mL. Then 1.0 mL of extract was evaporated using a water bath near to dryness. Exactly 50 mL of distilled water was then added to dilute and dissolve the remaining content. 4 mL anthrone reagent was added in a test tube containing 1.0 mL of sample solution and heated for 5 minutes in a boiling water bath. Then an uv-spectrophotometer was used to record absorbance at 620 nm wavelength [29]. The external standard was sourced from glucose, and gram glucose equivalents per 100 g of fresh carrot (g/100 g FW) was used to express the total sugar content.

### **2.7 Measurement of Mineral Elements of Carrots**

To determine different mineral elements (Ca, Mg, Na, K, P, and S), collected carrots samples were chopped into small pieces with a sharp stainless-steel knife and dried for roughly 72 hrs in an electric oven at 50 °C temperature. The samples were then pulverized in a grinding mill and utilized to make an extract by wet oxidation technique using a di-acid mixture [30]. Among the mineral elements, Ca and Mg were assessed titrimetrically, P and S were measured by spectrophotometry, and Na and K were estimated by flame photometry [30].

## **2.8 Statistical Analysis**

The data were statistically analyzed using the software package Minitab 17 following analysis of variance (ANOVA) and a general linear model. The least significant differences (LSD) were used to separate treatment means at the 5% level of probability at a specific time of data collection [31].

## **3. RESULTS AND DISCUSSION**

### **3.1 Weight Loss of Carrots at Storage**

The addition of chitosan had no significant effect on reducing the weight loss of carrots most of the time compared to the control except by 0.30% chitosan solution at 2 DAPS (Fig. 1). The addition of 0.30% chitosan solution significantly ( $P \leq 0.05$ ) reduced the weight of carrots compared to control at 2 DAPS. In general, the addition of chitosan reduced the weight loss of the carrot and prolonged the post-harvest storage time ranging from 4.61% weight loss to 25.09% compared to control which always had the highest percentage of weight loss ranging from 5.81% to 25.09% most of the time except by 0.10% chitosan solution at 2 DAPS. Among all the chitosan treatments the rate of weight loss of carrots decreased as the rate of chitosan increased from 0.10% to 0.30%. At all the times, the lowest weight loss was recorded by the 0.30% chitosan solution ranging from 4.61% to 23.61% weight loss at 2 to 8 DAPS, respectively. An almost similar reduction in weight loss has been reported after the addition of chitosan in tomato and grapefruit stored at room temperature [24, 32]. Chien et al. [33] found that citrus fruits covering with low molecular weight chitosan reduced weight loss considerably. They also reported that postharvest water retention prevents quick deterioration of fruits due to shriveling and that postharvest water loss may modify metabolism and, in certain cases, increase fruit ripening before shriveling becomes visible. As a result, limiting water loss from the fruit during storage or ripening aids in preserving fruit quality. However, present study results inferred that postharvest application of chitosan coating could be used to prevent weight loss of carrots.

**Figure 1: Effect of chitosan application on weight loss (%) of carrots at different days after post-harvest storage (DAPS) at room temperature. Each value is the mean for three replicates, and the vertical bar indicates the standard error. Red-colored bars are indicating the least significant difference (LSD) value at P-value  $\leq 0.05$ .**

### **3.2 Shelf-life of Carrots at Storage**

Compared to the fresh carrots (0 DAPS), the shelf life of all the carrots with all the treatments decreased as the time of storage progressed (Fig. 2). However, the addition of chitosan had significantly ( $P \leq 0.05$ ) reduced the loss of shelf-life of carrots compared to the control most of the cases except by 0.20% chitosan solution at 4 DAPS (Fig. 2). At 4 DAPS the loss of shelf-life of carrots showed no significant difference compared with control. The control had the highest loss of shelf-life of carrots at both the 4 and 8 DAPS recording 94.30% and 41.70% shelf-life, respectively whereas the lowest loss of shelf-life was observed by 0.10% chitosan solution at 4 DAPS (99.30% shelf-life) and by 0.20% chitosan solution at 8 DAPS (88.70% shelf-life). However, no significant difference in shelf-life of carrots was observed among 0.10%, 0.20% and 0.30% chitosan solution treatments at both 4 DAPS and 8 DAPS. Sultana *et al.* [24] reported a similar finding that the addition of chitosan during the postharvest storage increased the shelf-life of the tomatoes. According to Liu *et al.* [34], gray mold (caused by *Botrytis cinerea*) and blue mold (caused by *Penicillium expansum*) in tomato fruits stored at room and refrigeration temperature have been greatly reduced by the application of 0.5 and 1% chitosan solution. Furthermore, Romanazzi *et al.* [35] stated that both preharvest and postharvest application of chitosan had shown promising effects in disease control. They also stated that chitosan has a dual mechanism of action on pathogens and plants. According to their findings, chitosan inhibits the growth of decay-causing fungus and foodborne pathogens and induces resistance reactions in the tissues of the host plant. Thus, the study results suggest that chitosan

coating could be utilized to improve the shelf-life of carrots during postharvest storage, which could be owing to chitosan's ability to suppress postharvest damages of carrots due to various pathogens/microbes.

**Figure 2: Effect of chitosan application on shelf-life (%) of carrots at different days after post-harvest storage (DAPS) at room temperature. Each value is the mean for three replicates, and the vertical bar indicates the standard error. Red-colored bars are indicating the least significant difference (LSD) value at P-value  $\leq 0.05$ .**

### **3.3 Biochemical Properties of Carrots at Storage**

#### **3.3.1 Total sugar content**

The total sugar content of the carrots was decreased over time ranging from 2.02% to 2.19 % during storage compared to the fresh carrot (2.62 %) (Table 1). Chitosan addition at the rate of 0.10% and 0.20% were recorded a higher sugar content ranging from 2.14 to 2.19% compared to the control (2.05%) whereas 0.30% chitosan solution was recorded a slightly lower sugar content (2.02%). However, no significant difference was observed in the sugar content of carrots among any of the treatments. The degradation of sugar during post-harvest storage of carrots has also been reported in previous studies [36-37]. Similarly, the addition of chitosan was also reported to reduce sugar loss in pear compared to control [38]. Respiration has been accounted as the main reason for the loss in sugar content during storage in carrots. Sugar being utilized as a source of energy during respiration and other metabolic activities [36]. However, the chitosan coating is responsible for reducing respiration and other metabolic activities in carrots during storage [39] which was also may be the case in this study with the addition of chitosan.

### **3.3.2 Total phenolic content**

Similar to sugar content, the phenolic content of the carrots was decreased over time ranging from 0.48 to 0.78 mg/100 g compared to fresh carrots (0.90 mg/100 g) (Table 1). All the chitosan treatments significantly ( $P \leq 0.05$ ) reduced the phenolic content of carrots compared with control at 8 DAPS. There was no significant difference in phenolic content of carrots was observed between the 0.10% and 0.20%; 0.20% and 0.30% chitosan solution. The lowest phenolic content was recorded by 0.30% chitosan solution (0.48 mg/100 g) and was significantly ( $P \leq 0.05$ ) different compared to control and 0.10% chitosan solution. The decrease of phenol content in carrots during storage has also been reported [3]. Similar results were also reported for long-time storage of beans by Coelho *et al.* [40]. In addition to storage time, the phenolic content of vegetables during storage depends on the storage environment, methods, processing and genotypes. The storage of carrots with low oxygen has been reported to reduce phenolic content [41]. Thermal processing reduced the number of phenolic compounds, tannins, and antioxidant activity in vegetables [42]. Beninger *et al.* [43] reported that the total phenolic content of several pinto bean genotypes fluctuated significantly during storage. Thus, it can be inferred that several factors, viz. storage, genotypes, processing methods, and environmental conditions, play a vital role in the total phenolic contents of carrots.

### **3.3 Mineral Element Contents of Carrots at Storage**

The addition of chitosan coating increased the nutrient contents (Ca, Mg, Na, K, P and S) in carrots at 8 DAPS compared to the fresh carrots (0 DAPS) (Table 2). The chitosan coating significantly ( $P \leq 0.05$ ) increased the Ca content in carrots compared to the control whereas showed no significant difference in case of P and S. The addition of 0.30% chitosan solution recorded significantly lowest Mg contents in carrots at 8 DAPS whereas was significantly highest in case of Na and K content. According to Leclerc *et al.* [44], the carrot contains 0.34-0.37% Ca, 0.12-0.17% Mg, 2.80-3.21% K and 0.30-0.34% P depending on the types of fertilization. The nutrient content in carrots was slightly different in this study compared to the report by Leclerc *et al.* [44] which may be due to the difference in the genotype and variety. A similar finding was reported by Khazaei and Vandenberg [45], where stated that mineral nutrient contents in the bean were different depending on the genotype, environmental variation, and their interactions. The present study also showed minimum changes in the nutrient content of the carrot with chitosan treatments which may also be evident that chitosan can be safely added with the carrots during post-harvest storage as a food coating.

### **3.4 Correlations Among the Nutritional Qualities of Carrots**

Pearson correlation coefficients among the nutritional qualities of carrots harvested at 8 DAPS are summarized in Tables 2. Total phenol content of carrots showed a significant negative correlation with the Ca, Na and K. Thus, it can be inferred that total phenol contents in carrots negatively affected by the contents of Ca, Na and K at 8 DAPS. On the other hand, the total sugar content of the sugar of carrots showed no correlation with any of the nutrient elements determined in this study. Among the nutrient elements Ca showed a significant positive correlation with the Na, K and P. In addition, Na showed a significant positive correlation with the K, P and S; K showed a significant positive correlation with P and S; P showed a significant positive correlation with S. Magnesium showed negative relationship with all other nutrient elements of carrots, and among those, the relationship in between Mg and S was significantly negative ( $r = -0.562^*$ ). However, such inverse relationships indicated that the content of these mineral elements is moved in the reverse direction, i.e., when the content of any one of these elements is increased, the other is decreased with the same magnitude and vice-versa.

**Table 1: Effect of chitosan application on biochemical properties (total sugar and total phenol) and concentration ( $\pm$  SE) of different mineral elements of carrot samples (Ca, Mg, Na, K, P, S) at 8 days after post-harvest storage (DAPS) at room temperature. Different letters indicating statistical significance at P-value  $\leq$  0.05.**

Treatments (Chitosan solution)	Biochemical properties		Nutrient content (%) in carrots					
	Sugar (%)	Phenol (mg %)	Ca	Mg	Na	K	P	S
Control	2.05 $\pm$ 0.30(a)	0.78 $\pm$ 0.02(a)	0.26 $\pm$ 0.01(c)	0.26 $\pm$ 0.01(a)	0.61 $\pm$ 0.02(c)	1.09 $\pm$ 0.07(b)	0.23 $\pm$ 0.01(a)	0.093 $\pm$ 0.003(a)
0.10%	2.14 $\pm$ 0.24(a)	0.56 $\pm$ 0.02(b)	0.35 $\pm$ 0.02(b)	0.26 $\pm$ 0.01(a)	0.70 $\pm$ 0.03(bc)	1.14 $\pm$ 0.05(b)	0.23 $\pm$ 0.01(a)	0.096 $\pm$ 0.003(a)
0.20%	2.18 $\pm$ 0.15(a)	0.51 $\pm$ 0.03(bc)	0.40 $\pm$ 0.01(a)	0.27 $\pm$ 0.01(a)	0.77 $\pm$ 0.03(ab)	1.36 $\pm$ 0.06(a)	0.24 $\pm$ 0.01(a)	0.097 $\pm$ 0.006(a)
0.30%	2.02 $\pm$ 0.18(a)	0.47 $\pm$ 0.02(c)	0.41 $\pm$ 0.01(a)	0.22 $\pm$ 0.01(b)	0.86 $\pm$ 0.03(a)	1.50 $\pm$ 0.07(a)	0.27 $\pm$ 0.02(a)	0.095 $\pm$ 0.005(a)
LSD <sup>#</sup>	0.64	0.068	0.037	0.026	0.082	0.182	0.043	0.012
Fresh (0 DAPS)	2.62 $\pm$ 0.12	0.90 $\pm$ 0.02	0.28 $\pm$ 0.01	0.22 $\pm$ 0.01	0.54 $\pm$ 0.04	1.02 $\pm$ 0.05	0.22 $\pm$ 0.01	0.020 $\pm$ 0.003

<sup>#</sup>LSD: Least significant difference at P-value  $\leq$  0.05

**Table 2: Pearson correlation coefficients for nutritional qualities of carrots collected at 8 days after postharvest storage (DAPS) (n=12)**

	T-Phenol	T-Sugar	Ca	Mg	Na	K	P
T-Sugar	-0.078 <sup>ns</sup>						
Ca	-0.927 <sup>***</sup>	0.113 <sup>ns</sup>					
Mg	0.322 <sup>ns</sup>	0.201 <sup>ns</sup>	-0.153 <sup>ns</sup>				
Na	-0.821 <sup>***</sup>	0.003 <sup>ns</sup>	0.938 <sup>***</sup>	-0.285 <sup>ns</sup>			
K	-0.705 <sup>**</sup>	0.008 <sup>ns</sup>	0.871 <sup>***</sup>	-0.212 <sup>ns</sup>	0.968 <sup>***</sup>		
P	-0.429 <sup>ns</sup>	-0.114 <sup>ns</sup>	0.610 <sup>**</sup>	-0.076 <sup>ns</sup>	0.796 <sup>***</sup>	0.853 <sup>***</sup>	
S	-0.195 <sup>ns</sup>	-0.019 <sup>ns</sup>	0.443 <sup>ns</sup>	-0.562 <sup>*</sup>	0.504 <sup>*</sup>	0.542 <sup>*</sup>	0.733 <sup>***</sup>

Notes: T-Sugar: Total Sugar; T-Phenol: Total Phenol

<sup>ns</sup> non-significant; <sup>\*\*\*</sup> indicating significant at P<0.01, <sup>\*\*</sup> significant at P<0.05, <sup>\*</sup> significant at P<0.1

#### 4. CONCLUSION

The application of chitosan coating at different doses showed a remarkable positive effect to preserve weight loss and increase the shelf-life of carrots. Similarly, chitosan coatings also reduce the loss of total sugar contents in carrots during postharvest storage. Furthermore, the study found that chitosan coatings have the potential for enhancing some mineral nutritional aspects. However, total phenolic contents in carrots decreased considerably at storage, and the application of chitosan was unable to protect them. So, it can be inferred from this study that chitosan coating with 0.30% solution may be used to prevent weight loss, extend shelf-life, and improve some nutritional qualities of carrots. But peoples of our country are not familiar yet with such coatings in fruits and vegetables. Usually, they treated such coating as toxic and would like to avoid such coated agricultural commodities. Thus, consumer acceptance of such coated fruits and vegetables will need to be investigated in the future, and 'chitosan coating is non-toxic and safe'- a message must be circulated through print and electronic media.

#### REFERENCES

1. BBS. *Report on Agriculture and Rural Statistics*. Agriculture and Rural Statistics Survey (ARSS) Project-2017, Bangladesh Bureau of Statistics, Statistics and Informatics Division, Ministry of Planning, Government of the People's Republic of Bangladesh, Dhaka-1207. 2019. pp. 78.
2. Cheena J, Prashanth N, Seenivasan HN, Saidaiah. Effect of postharvest treatments on storage and quality of carrot cv. *New Kuroda* at ambient temperature. *Int J Curr Microbiol App Sci*. 2020; 9:2034-2040. <https://doi.org/10.20546/ijcmas.2020.909.253>
3. Haq RU, Prasad K. Antioxidant activity, phenolic, carotenoid and color changes in packaged fresh carrots stored under refrigeration temperature. *J Food Meas Charac*. 2017; 11(4):1542-1549. <https://doi.org/10.1007/s11694-017-9533-2>
4. The Daily Star. Tonnes of vegetables, fruits go to waste annually. Published on October 01 (2013). Available from <https://www.thedailystar.net/news/tonnes-of-vegetables-fruits-go-to-waste-annually>
5. Linares-Morales JR, Gutiérrez-Méndez N, Rivera-Chavira BE, Pérez-Vega SB, Nevárez-Moorillón GV. Biocontrol processes in fruits and fresh produce, the use of lactic acid bacteria as a sustainable option. *Front Sustain Food Syst*. 2018; 2:50. <https://doi.org/10.3389/fsufs.2018.00050>
6. Punja ZK, Collyer D, Scott C, Lung S, Holmes J, Sutton D. Pathogens and molds affecting production and quality of *Cannabis sativa* L. *Front plant sci*. 2019; 10:1120.

<https://doi.org/10.3389/fpls.2019.01120>

7. Katiyar D, Hemantaranjan A, Singh B. Chitosan as a promising natural compound to enhance potential physiological responses in plant: a review. *Indian J Plant Physiol.* 2015; 20(1):1-9. <https://doi.org/10.1007/s40502-015-0139-6>
8. Yadav A, Kumar N, Upadhyay A, Sethi S, Singh A. Edible coating as postharvest management strategy for shelf-life extension of fresh tomato (*Solanum lycopersicum* L.): An overview. *J Food Sci.* 2022; 87(6):1-35. <https://doi.org/10.1111/1750-3841.16145>
9. da Silva Rios DA, Nakamoto MM, Braga AR, da Silva EM. Food coating using vegetable sources: importance and industrial potential, gaps of knowledge, current application, and future trends. *Appl Food Res.* 2022; 2(1):100073. <https://doi.org/10.1016/j.afres.2022.100073>
10. Horská E, Šedík P, Mušínská K, Savitskaya T, Grinshpan D, Kačániová M. Acceptability of Edible Food Packaging in Slovakia: A Case Study on Young Generation. *Front Sustain Food Syst.* 2021; 5:720700. <https://doi.org/10.3389/fsufs.2021.720700>
11. Maringgal B, Hashim N, Tawakkal IS, Mohamed MT. Recent advance in edible coating and its effect on fresh/fresh-cut fruits quality. *Trends Food Sci Technol.* 2020; 96:253-267. <https://doi.org/10.1016/j.tifs.2019.12.024>
12. Meitha K, Pramesti Y, Signorelli S, Kriswantoro JA. Postharvest chitosan application maintains the quality of spinach through suppression of bacterial growth and elicitation. *Hort Environ Biotechnol.* 2022; 63(2):217-227. <https://doi.org/10.1007/s13580-021-00397-0>
13. Hesami A, Kavooosi S, Khademi R, Sarikhani S. Effect of chitosan coating and storage temperature on shelf-life and fruit quality of *Ziziphus Mauritiana*. *Int J Fruit Sci.* 2021; 21(1):509-518. <https://doi.org/10.1080/15538362.2021.1906825>
14. Lin Y, Li N, Lin H, Lin M, Chen Y, Wang H, Ritenour MA, Lin Y. Effects of chitosan treatment on the storability and quality properties of longan fruit during storage. *Food Chem.* 2020; 306:125627. <https://doi.org/10.1016/j.foodchem.2019.125627>
15. Romanazzi G, Feliziani E, Sivakumar D. Chitosan, a biopolymer with triple action on postharvest decay of fruit and vegetables: Eliciting, antimicrobial and film-forming properties. *Front Microbiol.* 2018; 9:2745. <https://doi.org/10.3389/fmicb.2018.02745>
16. Camatari FO, Santana LC, Carnellosi MA, Alexandre AP, Nunes MI, Goulart MO, Narain N, Silva MA. Impact of edible coatings based on cassava starch and chitosan on the post-harvest shelf life of mango (*Mangifera indica*) 'Tommy Atkins' fruits. *Food Sci Technol.* 2018; 38:86-95. <https://doi.org/10.1590/1678-457X.16417>
17. Kaczmarek MB, Struszczyk-Swita K, Li X, Szczęśna-Antczak M, Daroch M. Enzymatic modifications of chitin, chitosan, and chitooligosaccharides. *Front Bioeng Biotechnol.* 2019; 7:243. <https://doi.org/10.3389/fbioe.2019.00243>
18. Jiménez-Gómez CP, Cecilia JA. Chitosan: a natural biopolymer with a wide and varied range of applications. *Molecules.* 2020; 25(17):3981. <https://doi.org/10.3390/molecules25173981>
19. Mahira S, Jain A, Khan W, Domb AJ. *Chapter 1: Antimicrobial Materials- An Overview.* In: *Antimicrobial Materials for Biomedical Applications* (Eds. Domb, A. J., Kunduru, K. R. and Farah, S.). The Royal Society of Chemistry, UK. 2019. pp. 1-37.
20. Liang J, Yan H, Puligundla P, Gao X, Zhou Y, Wan X. Applications of chitosan nanoparticles to enhance absorption and bioavailability of tea polyphenols: A review. *Food Hydrocoll.* 2017; 69:286-92. <https://doi.org/10.1016/j.foodhyd.2017.01.041>
21. Hosseinnejad M, Jafari SM. Evaluation of different factors affecting antimicrobial properties of chitosan. *Int J Biol Macromol.* 2016; 85:467-475. <https://doi.org/10.1016/j.ijbiomac.2016.01.022>
22. Qi H, Hu W, Jiang A, Tian M, Li Y. Extending shelf-life of fresh-cut 'Fuji' apples with chitosan-coatings. *Innovative Food Sci Emerging Technol.* 2011; 12(1):62-66. <https://doi.org/10.1016/j.ifset.2010.11.001>
23. Xiao Z, Luo Y, Luo Y, Wang Q. Combined effects of sodium chlorite dip treatment and

- chitosan coatings on the quality of fresh-cut d'Anjou pears. *Postharvest Biol Technol.* 2011; 62(3):319-326. <https://doi.org/10.1016/j.postharvbio.2011.07.007>
24. Sultana N, Zakir HM, Parvin MA, Sharmin S, Seal HP. Effect of chitosan coating on physiological responses and nutritional qualities of tomato fruits during postharvest storage. *Asian J Adv Agril Res.* 2019; 10(2):1-11. DOI: 10.9734/AJAAR/2019/v10i230027
  25. Bautista-Baños S, Hernandez-Lauzardo AN, Velazquez-Del Valle MG, Hernández-López M, Barka EA, Bosquez-Molina E, Wilson CL. Chitosan as a potential natural compound to control pre and postharvest diseases of horticultural commodities. *Crop Prot.* 2006; 25(2):108-118. <https://doi.org/10.1016/j.cropro.2005.03.010>
  26. Rahman MA, Islam MA, Sultan MT, Kamal AH, Khan RA, Razzak M, Chowdhury MT, Khan MS, Mollah MZ. Assessment of chitosan as preservative on shelf life and major nutrient contents on fruits and vegetables. *Am J Agril Sci.* 2019; 6(1):1-10.
  27. Parvin MA, Zakir HM, Sultana N, Kafi A, Seal HP. Effects of different application methods of chitosan on growth, yield and quality of tomato (*Lycopersicon esculentum* Mill.). *Arch Agric Environ Sci.* 2019; 4(3):261-267. DOI: 10.26832/24566632.2019.040301
  28. Hossain MS, Iqbal A. Effect of shrimp chitosan coating on postharvest quality of banana (*Musa sapientum* L.) fruits. *Int Food Res J.* 2016; 23(1):277-283.
  29. Sadasivam S, Manickam A. *Biochemical Methods*. 2<sup>nd</sup> edn. New Age International (P) Limited, New Delhi-110 002, India. 1996; pp. 272.
  30. Singh D, Chhonkar PK, Pandey RN. *Soil, Plant and Water Analysis: A Method Manual*. IARI, New Delhi. India. 1999; pp. 134.
  31. Rahman MS, Scheffe C, Rajput S, Keizer D, Weatherley A. O-aryl and carbonyl carbon contents of food waste and biosolid predict P availability in an acidic soil. *Front Sustain Food Syst.* 2021; 4:609788. <https://doi.org/10.3389/fsufs.2020.609788>
  32. Meng X, Li B, Liu J, Tian S. Physiological responses and quality attributes of table grape fruit to chitosan preharvest spray and postharvest coating during storage. *Food Chem.* 2008; 106(2):501-508. <https://doi.org/10.1016/j.foodchem.2007.06.012>
  33. Chien PJ, Sheu F, Lin HR. Coating citrus (*Murcott tangor*) fruit with low molecular weight chitosan increases postharvest quality and shelf life. *Food Chem.* 2007; 100(3):1160-1164. <https://doi.org/10.1016/j.foodchem.2005.10.068>
  34. Liu J, Tian S, Meng X, Xu Y. Effects of chitosan on control of postharvest diseases and physiological responses of tomato fruit. *Postharvest Biol Technol.* 2007; 44(3):300-306. <https://doi.org/10.1016/j.postharvbio.2006.12.019>
  35. Romanazzi, G., Feliziani, E., Banos, S. B. and Sivakumar, D. (2017). Shelf life extension of fresh fruit and vegetables by chitosan treatment. *Crit Rev Food Sci Nutr.* 2017; 57: 579-601. <https://doi.org/10.3389/fmicb.2018.02745>
  36. Asgar A. Effect of storage temperature and type of packaging on physical and chemical quality of carrot. In: *IOP Conference Series: Earth and Environ Sci.* 2020; 443:012002. <https://doi.org/10.1088/1755-1315/443/1/012002>
  37. Devi KB, Bhadrachal B. Study of carrots for their storage environment and its effect on biochemical changes during pathogenesis. *Int J Sci Res.* 2015; 78:96-105.
  38. Sánchez C, Lidon FC, Vivas M, Ramos P, Santos M, Barreiro MG. Effect of chitosan coating on quality and nutritional value of fresh-cut 'Rocha'pear. *Emir J Food Agric.* 2015; 27(2):206-214. <https://doi.org/10.9755/ejfa.v27i2.19288>
  39. Lin B, Du Y, Liang X, Wang X, Wang X, Yang J. Effect of chitosan coating on respiratory behavior and quality of stored litchi under ambient temperature. *J Food Engg.* 2011; 102(1):94-99. <https://doi.org/10.1016/j.jfoodeng.2010.08.009>
  40. Coelho CM, de Mattos Bellato C, Santos JC, Ortega EM, Tsai SM. Effect of phytate and storage conditions on the development of the 'hard-to-cook'phenomenon in common beans. *J Sci Food Agric.* 2007; 87(7):1237-43. <https://doi.org/10.1002/jsfa.2822>
  41. Amanatidou A, Slump RA, Gorris LGM, Smid EJ. High oxygen and high carbon dioxide

- modified atmospheres for shelf-life extension of minimally processed carrots. *J Food Sci.* 2000; 65:61-66. <https://doi.org/10.1111/j.1365-2621.2000.tb15956.x>
42. Silva MO, Brigide P, Toledo NM, Canniatti-Brazaca SG. Phenolic compounds and antioxidant activity of two bean cultivars (*Phaseolus vulgaris* L.) submitted to cooking. *Braz J Food Technol.* 2017; 21:2016072. <https://doi.org/10.1590/1981-6723.7216>
  43. Beninger CW, Gu L, Prior RL, Junk DC, Vandenberg A, Bett KE. Changes in polyphenols of the seed coat during the after-darkening process in pinto beans (*Phaseolus vulgaris* L.). *J Agril Food Chem.* 2005; 53(20):7777-7782. <https://doi.org/10.1021/jf050051l>
  44. Leclerc J, Miller ML, Joliet E, Rocquelin G. Vitamin and mineral contents of carrot and celeriac grown under mineral or organic fertilization. *Biol Agric Hort.* 1991; 7(4):339-348. <https://doi.org/10.1080/01448765.1991.9754564>
  45. Khazaei H, Vandenberg A. Seed mineral composition and protein content of faba beans (*Vicia faba* L.) with contrasting tannin contents. *Agron.* 2020; 10(4):511. <https://doi.org/10.3390/agronomy10040511>