

# Effect of Organic Nutrients on Growth and Flowering Attributes of African Marigold (*Tagetes erecta* L.) under Terai Region of West Bengal

## ABSTRACT

A field experiment was conducted at Instructional farm, Department of Floriculture, Medicinal and Aromatic plants, Faculty of Horticulture, Uttar Banga Krishi Viswavidyalaya during 2019-2020 to study the "Effect of organic nutrients on growth and flowering attributes of African Marigold (*Tagetes erecta* L.) under Terai region of West Bengal". The experiment was laid out by following Factorial Randomized Complete Block Design, consists of two factors in which first factor was varieties (Pusa Bahar- $V_1$  and Pusa Narangi Gaiinda-  $V_2$ ) and second factor was ten treatments combination. Different Organic nutrients like Farm Yard Manure (FYM), Vermicompost, Poultry Manure, Sea Weed extract, Vermicompost wash and Mustard oil cake extract used in the experiments as basal application and foliar spray according to the treatments. The vegetative and floral attributes were recorded at peak flowering stage and the data was statistically analysed. The data was significantly differed among the varieties, treatments and different treatment combinations with varieties. Between the two varieties, var. Pusa Narangi Gaiinda ( $V_2$ ) showed early flowering compared to var. Pusa Bahar ( $V_1$ ) but the longer duration of flowering (49.41 days) was observed with the var. Pusa Bahar ( $V_1$ ). Among the ten treatments, the treatment  $T_{10}$  (Poultry Manure @ 2.5t/ha + Vermicompost wash @ 3%) resulted the best against the important flowering parameters like total number of flowers per plant (55.73) and weight of ten fresh flowers (103.08 g). The var. Pusa Narangi Gaiinda when treated with Poultry Manure @ 2.5t/ha + Vermicompost wash @ 3% ( $T_{10}$ ) showed better performance in respect of several morphological and quality parameters of African Marigold. It may be recommended that the application of Poultry Manure @ 2.5t/ha as basal with combination of Vermicompost wash @ 3% spraying at twice (30 and 45 days after transplanting) during the plant growth for quantitative and qualitative produces of African Marigold in the terai region of West Bengal.

*Key words: African Marigold, Varieties, Organic Nutrient Sources*

## INTRODUCTION

African Marigold (*Tagetes erecta* L.) belongs to the family Asteraceae bearing chromosome  $2n=24$ , which occupied an important place in Ornamental Horticulture. It is native of Central and Southern America especially in Mexico (Kaplan, 1960) [1]. African marigold is an annual, hardy and attains a height of 60-90 cm with crop duration of 3-4 months. African Marigold gained popularity among flower growers due to its easy cultivation, and year-round demand of consumers. It is mainly grown for loose flower which are being used in making garland, festival occasion, poojas, marriage ceremony, social functions etc. African Marigold is suitable for landscaping, garden display, for bedding, as an herbaceous border and it is also ideal for newly planted shrubberies to provide multi colours display, rockeries, edging and window boxes. For ecofriendly cultivation and quality produces of marigold, it is essential to change the traditional cultivation techniques in which organic manures and chemical fertilizers used. Organic

**Comment [H1]:** In the abstract section, state the purpose and background of the study.

**Comment [H2]:** What is the meaning of var?

farming is a holistic way of agriculture for conserving the soil physical and biological properties. It does not have any adverse effect on the ecosystem. Indian agriculture has a great opportunity to convert itself as organic cultivation by avoiding chemical fertilizers and pesticides in agriculture. Popularization of organic farming can effectively prevent the entry of pesticides and toxicants in the food chain and prevent soil and water pollution (Sreenivasaet al., 2009[2] and Natarajan, 2007) [3]. Hence, we are using various organic manures like FYM, Vermicompost, Poultry Manure, Sea Weed extracts as the sources of different nutrients. These are organic inputs can be applied in soil directly or we can spray to the standing crops also. Keeping these in view, the present investigation entitled “Effect of organic nutrients on growth and flowering attributes of African marigold (*Tagetes erecta*L.) under Terai region of West Bengal”.

**Comment [H3]:** If possible, providing information on the organic fertilizers used both strengthens the introduction and supports the purpose of the study.

#### MATERIALS AND METHODS

**Comment [H4]:** Explain the method used in the analysis of the data.

The present experiment was conducted at Instructional farm, Department of Floriculture, Medicinal and Aromatic plants, Faculty of Horticulture, Uttar BangaKrishiViswavidyalaya during 2019-2020. The experiment was laid out by following factorial Randomized Complete Block Design, consists of two varieties with ten treatments combination. The place is located in Terai Region of West Bengal at 26° 19 N latitude and 89° 23 E longitude. The site lies in the sub-Himalayan plains at an elevation of 43 metres above mean sea level. This zone is marked by a typical and sub-tropical climate with high relative humidity (86%), moderate temperature(8.30°C – 31.00°C), rainfall (2 mm) during the period of experiment and prominent winter continues from September to March. The treatments of the experiment designed statistically as two factors Randomized Complete Block Design. The factors were varietal reaction with organic nutrient sources used as singly or in combination which are given in the table no.1. The varieties namely PusaBahar (V<sub>1</sub>) and PusaNarangaiGaiinda (V<sub>2</sub>) collected from Indian Agricultural Research Institute (IARI), NewDelhi. Different Organic nutrients used in the experiments like Farm Yard Manure (FYM), Vermicompost, Poultry Manure, Sea Weed extract, Vermiwash and Mustard oil cake extract have been collected from adjoining areas of the farm and reputed shops of the district of Cooch Behar, West Bengal. Land was ploughed and pulverized properly with the help of Power tiller. The different organic nutrient sources applied as basal and spraying on foliage as per the treatments. The basal dose of organic manures viz., FYM, Vermi compost and Poultry manure were applied at the time of one week before transplanting of seedlings.

**Comment [H5]:** Express it as Table 1.

**Comment [H6]:** Does the term farmyard manure mean cattle manure?

**Comment [H7]:** When you use an abbreviation in the text, always use the same abbreviation in the rest of the text.

**Comment [H8]:** What is the meaning of viz?

**Comment [H9]:** Make sure to always use the terms in the same pattern.

**Table No-1: The treatment details of the experiment**

First factor	Varieties
Notation	Name of the variety
V <sub>1</sub>	PusaBahar
V <sub>2</sub>	PusaNarangaiGaiinda
Second factor	Organic manures application
Notation	Treatment details
T <sub>1</sub>	Control (without organic and inorganic fertilizers)
T <sub>2</sub>	Farm yard manure (FYM) @ 30 t/ha and Sea weed Extract spray @ 3%

T <sub>3</sub>	Farm yard manure (FYM) @ 30 t/ha and Mustard oil Cake @ 3%
T <sub>4</sub>	Farm yard manure (FYM) @ 30 t/ha and Vermiwash@3%
T <sub>5</sub>	Vermi compost @5t/ha and Sea weed Extract spray@3%
T <sub>6</sub>	Vermi compost @5t/ha and Mustard oil Cake @ 3%
T <sub>7</sub>	Vermi compost @5t/ha and vermi wash @3%
T <sub>8</sub>	Poultry Manure@2.5t/ha and Sea weed Extract spray@3%
T <sub>9</sub>	Poultry Manure@2.5t/ha and Mustard oil Cake @ 3%
T <sub>10</sub>	Poultry Manure@2.5t/ha and vermi wash @3%

Whereas other organic nutrient sources like fermented raw mustard oil cake and vermin wash (collected from vermi pit of the University) were diluted by water. Ten times dilution in volume with water was done for application as spray. The seedlings have been transplanted from nursery beds after 45 days from sowing. The seedlings transplanted in prepared plots (200 cm X 200 cm) with a spacing of 40 cm (row to row) X 40 cm (plant to plant). The planting was done at first week of November during morning hours and immediate after planting, light irrigation was applied to all the plots with the help of Rosecan. All the vegetative, floral and yield parameters were recorded at peak flowering stage according to the treatments.

**Comment [H10]:** Pay attention to text formatting (carriage returns, periods, commas and word spaces)

## RESULTS AND DISCUSSION

The experimental results obtained from the present investigation as well as discussion have been summarized under followings heads:

### Effect of Organic manures on vegetative parameters

The vegetative parameters like plant height and plant spread have been taken and statistically analysed which represented in the Table no.2. The plant height and plant spread significantly differed among the varieties as well as among the treatments. In the table no-2, between the two varieties (first factor), the maximum plant height was recorded in var. PusaNarangiGainda (74.60 cm) and the lowest was found in var. PusaBahar (62.14 cm). The highest plant height was observed in T<sub>10</sub> (74.98 cm) and it was statistically at par with T<sub>7</sub> (74.12 cm), T<sub>8</sub> (73.09 cm) and T<sub>9</sub> (74.09 cm). The lowest plant height was observed in T<sub>1</sub> (63.28 cm) among the ten treatments. Plant spread both (E-W and N-S) was found more in var. PusaBahar (53.43 cm, 54.33 cm respectively) compared to PusaNarangiGainda (48.46 cm, 49.92 cm respectively). Plant spread (E-W) was found maximum in T<sub>10</sub> (59.82 cm) and it was found that T<sub>9</sub> (58.84 cm) and T<sub>8</sub> (57.48 cm) were statistically at par with the treatment T<sub>10</sub> where as the lowest in T<sub>1</sub> (45.25 cm). Plant spread (N-S) was found maximum in T<sub>8</sub> (67.09 cm) which is statistically at par with T<sub>9</sub> (65.16 cm) and T<sub>10</sub> (66.06 cm) and minimum plant spread was found in T<sub>1</sub> (45.25 cm) among the ten treatments.

**Comment [H11]:** Express it as Table 2

**Comment [H12]:** It may be more appropriate to position your tables close to where they appear in the text.

The vegetative parameters like plant height and plant spread significantly differed among the different treatment combinations. The data presented in the table No.2 showed that the maximum plant

height (81.76 cm) was found in var. PusaNarangiGai ndawhen treated withPoultry Manure@2.5t/haper

Treatment details	Plant Height(cm)	Plant Spread(cm)	Days for first flower bud	Days for first flower full	Days for first flower wilting
-------------------	------------------	------------------	---------------------------	----------------------------	-------------------------------

hectare and in combination with vermi wash spraying @3% (T<sub>10</sub>) and the minimum plant height was found in var. PusaBaharwhen treated without organic and inorganic fertilizers (T<sub>1</sub>-Control). Plant spread {(E-W) (62.66 cm)} & {(N-S) (69.02 cm) respectively} was found maximum in var. PusaBahar (V<sub>1</sub>)when treated with Poultry Manure@2.5t and in combination with vermi wash spraying @3% (T<sub>10</sub>) and var. PusaNarangiGai nda (V<sub>2</sub>) when treated with Poultry Manure@2.5t/hectareand in combination with Sea weed Extract spraying@3%(T<sub>8</sub>)respectively whereas, the lowest Plant spread {(E-W) (42.21 cm)} & {(N-S) (43.46 cm)} was found in var. PusaNarangiGai nda when treated without organic and inorganic fertilizers (T<sub>1</sub>-Control)and in var. PusaBahar (V<sub>1</sub>) when treated without organic and inorganic fertilizers (T<sub>1</sub>-Control)respectively.

The plant height and plant spread were significantly influenced among the different treatment combinations as well as varieties. The vegetative attributes showed the best resultswhen application of poultry manures as basal followed by vermi wash spraying. It might be due to application of Poultry manure which is abundant source of major nutrients viz., NPK and other essential nutrients (Garg and Bahal, 2008) [4]. Nitrogen is the main constituent of chlorophyll, amino acids and protein which plays important role in cell division, protein synthesis and metabolite transport for building the plant tissues (Gupta and Prasad, 1991) [5]. Due to presence of higher amount of major essential nutrients and readily available form of nitrogen in uric acid and narrow C: N ratio than other nutrient facilitates increase in plant height, plant spread and number of leaves. The above findings were similar to the earlier study conducted by Malik *et al.*, (2021) [6], Swathi *et al.*, (2018) [7] in Marigold and Suseela *et al.*, (2016) [8] in Tuberose.

The flowering parameters like days taken to first flower bud initiation, days taken for first flower full blooming and days taken to first flower wilting were have been taken and statistically analysed and represented in the Table no.2. These flowering parameters significantly differed among the varieties as well as among the treatments. In the table no.2 between the two varieties (first factor) the maximum days taken to first flower bud initiation (42.07 days), maximum days taken for first flower full blooming (13.71 days) and maximum days taken to first flower wilting (10.75 days) were found in the var. Pusa

**Comment [H13]:** Pay attention to text formatting (carriage returns, periods, commas and word spaces)

<b>V<sub>1</sub></b>	62.14	53.43	54.33	42.07	13.71	10.75
<b>V<sub>2</sub></b>	74.60	48.46	49.92	39.17	13.40	9.96
<b>S. Em±</b>	0.60	0.45	0.35	0.18	0.18	0.11
<b>C.D. at 5%</b>	1.705	1.276	0.993	0.524	0.527	0.313
<b>Treatment details</b>						
<b>T<sub>1</sub></b>	63.28	45.64	45.25	46.37	16.62	7.78
<b>T<sub>2</sub></b>	69.86	51.89	53.94	38.43	12.15	10.97
<b>T<sub>3</sub></b>	68.13	51.44	54.40	39.07	13.70	10.38
<b>T<sub>4</sub></b>	69.93	52.92	53.51	39.43	14.00	10.77
<b>T<sub>5</sub></b>	68.92	55.82	55.52	41.13	9.63	13.30
<b>T<sub>6</sub></b>	71.03	54.96	55.25	41.83	10.70	13.23
<b>T<sub>7</sub></b>	74.12	56.03	56.93	41.17	10.67	13.60
<b>T<sub>8</sub></b>	73.09	57.48	67.09	40.30	12.83	10.40
<b>T<sub>9</sub></b>	74.08	58.84	65.16	40.33	12.62	11.07
<b>T<sub>10</sub></b>	74.98	59.82	66.06	40.53	13.13	10.67
<b>S. Em±</b>	1.33	1.00	0.78	0.41	0.41	0.24
<b>C.D. at 5%</b>	3.81	2.85	2.22	1.17	1.18	0.70
<b>V (Variety) X T (Treatments)</b>						
<b>V<sub>1</sub> x T<sub>1</sub></b>	55.10	49.06	47.03	47.07	16.13	8.13
<b>V<sub>1</sub> x T<sub>2</sub></b>	62.68	52.27	54.60	41.67	12.13	11.27
<b>V<sub>1</sub> x T<sub>3</sub></b>	63.35	53.53	56.59	40.40	15.33	10.37
<b>V<sub>1</sub> x T<sub>4</sub></b>	62.79	55.07	56.27	40.80	15.73	10.20
<b>V<sub>1</sub> x T<sub>5</sub></b>	66.68	57.23	57.15	40.40	9.20	13.80
<b>V<sub>1</sub> x T<sub>6</sub></b>	68.52	57.01	56.14	41.47	9.50	13.53
<b>V<sub>1</sub> x T<sub>7</sub></b>	67.87	53.99	57.15	41.33	8.93	13.67
<b>V<sub>1</sub> x T<sub>8</sub></b>	67.45	59.76	65.17	39.67	12.40	10.33
<b>V<sub>1</sub> x T<sub>9</sub></b>	69.27	61.24	65.50	39.00	11.10	11.93
<b>V<sub>1</sub> x T<sub>10</sub></b>	68.19	62.66	66.63	39.53	12.47	11.47
<b>V<sub>2</sub> x T<sub>1</sub></b>	71.39	42.21	43.46	45.67	17.10	7.43
<b>V<sub>2</sub> x T<sub>2</sub></b>	77.05	51.51	53.27	35.20	12.17	10.67
<b>V<sub>2</sub> x T<sub>3</sub></b>	72.90	49.35	52.20	37.73	12.07	10.40
<b>V<sub>2</sub> x T<sub>4</sub></b>	77.07	50.77	50.74	38.07	12.27	11.33
<b>V<sub>2</sub> x T<sub>5</sub></b>	71.16	54.42	53.88	41.87	10.07	12.80
<b>V<sub>2</sub> x T<sub>6</sub></b>	73.55	52.90	54.35	42.20	11.90	12.93
<b>V<sub>2</sub> x T<sub>7</sub></b>	80.37	58.07	56.70	41.00	12.40	13.53
<b>V<sub>2</sub> x T<sub>8</sub></b>	78.74	55.20	69.02	40.93	13.27	10.47
<b>V<sub>2</sub> x T<sub>9</sub></b>	78.89	56.45	64.81	41.67	14.13	10.20
<b>V<sub>2</sub> x T<sub>10</sub></b>	81.76	56.97	65.49	41.53	13.80	9.87
<b>S. Em±</b>	1.88	1.41	1.10	0.58	0.58	0.35
<b>C.D. at 5%</b>	5.39	4.04	3.14	1.66	1.67	0.99

Bahar. The minimum days taken to first flower bud initiation (39.17 days), minimum days taken for first flower full

blooming (13.40 days) and minimum days taken to first flower wilting (9.96 days) were found in the var. PusaNarangiGainda. Among the ten treatments, T<sub>1</sub> (Control (without organic and inorganic fertilizers) recorded maximum days taken to first flower bud initiation (46.83 days), days taken for first flower full blooming (16.62 days) and T<sub>7</sub> (Vermi compost @5t/ha and vermi wash @3%) recorded maximum days taken to first flower wilting (13.60 days). while it was minimum (38.43 days) days taken to first flower bud initiation in T<sub>2</sub>, minimum days taken for first flower full blooming (9.63days) was found in T<sub>5</sub> and T<sub>1</sub>(Control (without organic and inorganic fertilizers) recorded minimum days taken for first flower wilting (7.76 days).

The Flowering parameters significantly differed among the different treatment combinations. The maximum days taken to first flower bud initiation (47.07 days) was found in PusaBahar (V<sub>1</sub>) when treated without organic and inorganic fertilizers (T<sub>1</sub>-Control) and minimum days taken to first flower bud initiation (35.20 days) was recorded in var. PusaNarangiGainda (V<sub>2</sub>) when treated with Farm yard manure (FYM) @ 30 t/ha in combination with Sea weed Extract spray@3% (T<sub>2</sub>). The maximum days taken for first flower full blooming (17.10 days) was recorded in var. PusaNarangiGainda (V<sub>2</sub>) when treated without organic and inorganic fertilizers (T<sub>1</sub>-Control) and minimum days taken for first flower full blooming (8.93 days) recorded in var. PusaBahar (V<sub>1</sub>) when treated with Vermicompost @5t/ha and in combination with vermicompost spraying @3% (T<sub>7</sub>) whereas, maximum days taken to first flower wilting (13.67 days) was noticed in var. PusaBahar (V<sub>1</sub>) when treated with Vermicompost @5t/ha and in combination with vermicompost spraying @3% (T<sub>7</sub>). The minimum days taken to first flower wilting (7.43 days) was found in var. PusaNarangiGainda (V<sub>2</sub>) when treated without organic and inorganic fertilizers (T<sub>1</sub>-Control).

Plants treated with Farm Yard Manure (FYM) @ 30 t/ha + Sea Weed Extract spraying@ 3% recorded minimum days taken for first flower bud initiation. It may be due to amplification of nutrients mainly nitrogen, Phosphorous and potassium through FYM which paved the translocation of nutrients to phytohormones to the shoots resulting in early bud and flower initiation. Besides, increase in root density might be due to sea weed extract spray which promoted the more phosphorous absorption by roots and which helped in early bud initiation (Ali *et al* 2021) [9]. The above findings are in conformity with Singh and Vikas Kumar (2016) [10] in Marigold and Khanna *et al.*, (2016) [11] in China aster.

Earliness in first flower full blooming may be due to supplementing nutrients (macro & micro), growth regulators and other enzymes by vermicompost which helped in better translocation of nutrients. This earliness also due to gibberellins in vermicompost, which are directly associated with regulation of flowering. The above findings are in corroborating with the Bordoloi and Madhumita Choudhury Talukdar (2019) [12] in Chrysanthemum.

The flowering parameters like duration of flowering, flower diameter, weight of ten fresh flowers, total number of flowers per plant and shelf life of flowers have been taken and statistically analysed and represented in the Table no.3. These flowering parameters significantly differed among the varieties as well as among the treatments. In the table no.3 between the two varieties (first factor). The duration of flowering was recorded maximum (49.41 days) in the var. PusaBahar and minimum (49.35 days) was

found in var. PusaNarangiGainda. The maximum flower diameter (5.94 cm) was recorded in var. PusaNarangiGainda and minimum flower diameter was found in var. PusaBahar (5.37 cm). The maximum weight of ten fresh flowers (91.79 g.), total number of flowers per plant (53.49) and highest shelf life of flower (5.38 days) was recorded in var. PusaNarangiGainda whereas, minimum weight of ten fresh flowers (88.78 g.), total number of flowers per plant (37.65) and highest shelf life of flower (5.23 days) was found in var.PusaBahar.

Among Ten different treatments the maximum duration of flowering (61.57 days) and flower diameter (6.18) was recorded in T<sub>7</sub> and T<sub>9</sub> respectively. The maximum weight of ten flowers (103.08 g.), number of flowers per plant (55.73) and shelf life (7.28 days) was recorded in T<sub>10</sub> and T<sub>6</sub> respectively. It was found that duration of flowering (43.20 days), flower diameter (4.64 cm), weight of ten flowers (82.08 g.), number of flowers per plant (35.60) and shelf life (3.87 days) was found minimum in T<sub>1</sub>. The Flowering parameters significantly differed among the different treatment combinations. The data presented in Table No.-3 (Fig No.-1) revealed that duration of flowering (61.90 days) and flower diameter (6.63 cm) was found maximum in var.PusaBahar (V<sub>1</sub>) when treated with Vermi compost @5t/ha and in combination with vermi wash spraying @3% (T<sub>7</sub>) and in var. PusaNarangiGainda (V<sub>2</sub>) treated with Vermi compost @5t/ha and in combination with Mustard oil Cake spraying @ 3%(T<sub>6</sub>) respectively whereas, it was found duration of flowering (38.27 days) and flower diameter (4.51 cm) was found minimum in var. PusaBahar (V<sub>1</sub>) when treated without organic and inorganic fertilizers (T<sub>1</sub>-Control). The maximum weight of ten fresh flowers (116.67 g.) was found in var. PusaNarangiGainda(V<sub>2</sub>) when treated with Vermi compost @5t/ha and in combination with vermi wash spraying @3% (T<sub>7</sub>) and minimum weight of ten fresh flowers (80.50 g.) observed in var PusaBahar (V<sub>1</sub>) when treated without organic and inorganic fertilizers (T<sub>1</sub>-Control). Number of flowers per plant (63.27) and shelf life (8.60 days) was recorded highest in var. PusaNarangiGainda (V<sub>2</sub>) when treated with Poultry Manure@2.5t and in combination with vermi wash spraying @3% (T<sub>10</sub>) whereas, it was minimum in varPusaBahar (V<sub>1</sub>) when treated without organic and inorganic fertilizers (T<sub>1</sub>-Control).

The duration of flowering which deals for availability of flowers for longtime, is an important factor for more return from per unit area. Increase in duration of flowering might be due to mobilization of nutrients, increase in synthesis of proteins and degradation of chlorophyll prevention due to sufficient amount of nutrients supplied by the vermicompost and vermiwash. This finding is in conformity with Kumar *et al.*, 2020 [13], Kumar and Vishwanath (2020) [14], Mittal *et al.*, (2010) [15] in Marigold, Bordoloi and Madhumita Choudhury Talukdar (2019) in Chrysanthemum [12], Pandey *et al.* [16], (2017) in Dhalia. Application of Poultry manure@ 2.5t/ha in combination with Fermented Mustard oil Cake spraying @ 3% increased the flower diameter due to availability of sufficient macro and micro nutrients supplied by poultry manure which ultimately resulted in vigorous growth of plants. Increase in phosphorous is found to be responsible for the initiation of floral primordial formation resulting in the increase in flower diameter. The above finding is in conformity with Bohra *et al.*, 2014[16] in chrysanthemum.

**Fig.-1: Effect of different organic nutrient sources on flowering duration of African marigold.**

**Comment [H14]:** Check the compatibility of your references in the text with your references in the reference section.

**Comment [H15]:** Please note Pandey 16 reference

**Fig.-2: Effect of different organic nutrient sources on total number of flowers per plant of African marigold**

*Figure 3 : Effect of organic nutrient sources on floral and yield attributes of African Marigold*

Treatment details	Flowering duration(days)	Flower diameter(cm)	Weight of 10 fresh flowers (g)	Total number of flowers per plant	Shelf life of loose flower(days)
V <sub>1</sub>	49.41	5.37	88.78	37.65	5.23
V <sub>2</sub>	49.35	5.94	91.79	53.49	5.38
<b>S. Em±</b>	0.17	0.04	1.16	1.17	0.09
<b>C.D. at 5%</b>	0.478	0.127	3.313	3.363	0.268
<b>Treatment details</b>					
T <sub>1</sub>	43.20	4.64	82.08	35.60	3.87
T <sub>2</sub>	49.00	5.83	88.50	49.97	5.63
T <sub>3</sub>	49.80	6.05	96.78	46.73	5.88
T <sub>4</sub>	50.57	5.87	90.42	48.70	5.65
T <sub>5</sub>	59.57	6.13	90.50	47.93	6.67
T <sub>6</sub>	61.23	6.04	88.17	54.23	7.28
T <sub>7</sub>	61.57	5.94	98.71	48.63	6.80
T <sub>8</sub>	55.23	5.97	95.38	50.50	5.78
T <sub>9</sub>	52.67	6.18	100.21	49.93	5.83
T <sub>10</sub>	52.50	5.94	103.08	55.73	5.78
<b>S. Em±</b>	0.37	0.10	2.59	2.63	0.21
<b>C.D. at 5%</b>	1.07	0.28	7.41	7.52	0.60
<b>V (Variety) X T (Treatments)</b>					
V <sub>1</sub> x T <sub>1</sub>	38.27	4.51	80.50	33.67	4.20
V <sub>1</sub> x T <sub>2</sub>	48.40	5.35	86.50	50.00	5.20
V <sub>1</sub> x T <sub>3</sub>	49.87	5.71	93.88	38.27	5.57
V <sub>1</sub> x T <sub>4</sub>	51.20	5.45	87.50	42.00	5.57
V <sub>1</sub> x T <sub>5</sub>	59.33	5.84	95.50	39.80	5.60
V <sub>1</sub> x T <sub>6</sub>	61.40	5.45	85.00	47.73	5.97
V <sub>1</sub> x T <sub>7</sub>	61.93	5.39	80.75	47.33	5.73
V <sub>1</sub> x T <sub>8</sub>	57.07	5.55	85.25	51.40	6.23
V <sub>1</sub> x T <sub>9</sub>	53.33	5.77	99.75	51.07	6.07
V <sub>1</sub> x T <sub>10</sub>	52.07	5.85	104.33	61.47	6.10
V <sub>2</sub> x T <sub>1</sub>	48.13	4.77	83.67	37.53	3.53
V <sub>2</sub> x T <sub>2</sub>	49.60	6.31	90.50	36.67	6.07
V <sub>2</sub> x T <sub>3</sub>	49.73	6.39	99.67	55.20	6.20
V <sub>2</sub> x T <sub>4</sub>	49.93	6.29	93.33	55.40	5.73
V <sub>2</sub> x T <sub>5</sub>	59.80	6.41	85.50	56.07	7.73
V <sub>2</sub> x T <sub>6</sub>	61.07	6.63	91.33	60.73	5.47
V <sub>2</sub> x T <sub>7</sub>	61.20	6.48	116.67	49.93	7.87
V <sub>2</sub> x T <sub>8</sub>	53.40	6.38	105.50	49.60	5.33
V <sub>2</sub> x T <sub>9</sub>	52.00	6.59	100.67	48.80	5.60
V <sub>2</sub> x T <sub>10</sub>	52.93	6.02	101.83	63.27	8.60

<b>S. Em±</b>	0.53	0.14	3.66	3.71	0.30
<b>C.D. at 5%</b>	1.51	0.40	10.48	10.63	0.85

Plants treated with Vermicompost @ 5t/ha + Vermiwash @ 3% showed increased weight of flowers. It may be due to presence of existent growth promoting substances like essential lant nutrients, enzymes vitamins, and antibiotics in vermicompost and vermiwash. Present findings are in similar with reported by Kumar and Vishwanath (2020) [14], Sudhagaret *al.*, (2019) [18] in African marigold and Bohra *et al.*, (2014) [17]

The data presented in Table No.-3 (Fig No.-1) revealed that Poultry Manure@ 2.5t/ha + Vermiwash @ 3% application showed highest number of flowers per plant and shelf life of flowers. The maximum flower production could be due to poultry manure which increased the water holding capacity of soil by improving soil texture, also improves humus status of soil. It also helps in improving physico chemical properties of the soil viz., pH, EC, organic carbon, macro and micro nutrients. It may be due vermiwash which contains all secondary, macro, micro nutrients and presence of ethylene inhibitors or due to cytokinins may also enhanced the uptake of Ca and Mg. Similar result were also reported by Gargeet *al.*, (2020) [4], Paryaet *al.*, (2017) [19], Mittal *et al.*, (2010) [15] and Bohra *et al.*, (2014) [17].

Comment [H16]: Please note Garg 4 reference

## CONCLUSION

Based on the result of the present study It may be recommended that the application of Poultry Manure @ 2.5t/ha as basal with combination of Vermi wash @3% spraying at twice (30 and 45 days after transplanting) during the plant growth for quantitative and qualitative produces of African Marigold in the Terai Region of West Bengal.

## COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## REFERENCE

1. Kaplan (1960). Economic Botany, 14:200.
2. Sreenivasa, M. N., Nagaraj. M., Naik and Bhat, S. N. 2009. Beneficial traits of microbial isolates of organic liquid manures. First Asian PGPR Congress for sustainable agriculture, 21-24 June, ANGRAU, Hyderabad.

3. Natarajan, K., 2007. Panchagavya for plant. Proc. Nation. Conf. on Glory of Gomatha, Dec. 1- 3, S. V. Veterinary Uni, Tirupati, A. P., pp. 72-75.
4. Garge V C, Malik S, Awasthi M, Singh S P, Chaudhary M and Kumar A (2020) Effect of integrated nutrient management on flower quality of French Marigold (*Tagetespatula* L.) Cv.PusaArpita. *International Research Journal of Pure and Applied Chemistry* 21(24); 154-156.
5. Gupta R D and Prasad. 1991. Phosphobacterin is an ideal Phosphatic bacterial fertilizer. *Farmer's Digest* 5(4); 13-14.
6. Malik M, Kumar T, Jawla S K and Sahrawat A (2021) Economic analysis of Marigold production under the different application of organic manures. *The Pharma Innovation Journal* 10 (3);155-157.
7. Swathi K, Sarkar I, Maitra S and Sharma S (2018)Organic manures and bio-inoculants mediated influence on growth and flowering of African Marigold (*Tagetes erecta*L.) cv. PusaNarangiGainda. *International Journal of Bio-resource and Stress Management* 8(3); 429-432.
8. Suseela T, Chandrasekha R, Bhaskar V V, Suneetha D R S and Umakrishna K (2016) Effect of organic manures, inorganic fertilizers and micronutrients on vegetative and floral characters of tuberose (*Polianthes tuberosa* L.) cv. Suvasini. *International Journal of Scientific and Research Publications* 6(2); 170-173.
9. Ali, O.; Ramsubhag, A.;Jayaraman, J. Biostimulant Properties of Seaweed Extracts in Plants: Implications towards Sustainable Crop Production. *Plants* 2021, 10, 531
10. Singh U and Kumar V (2016) Studies on Efficacy of Integrated Nutrient Management on Growth and Yield of Marigold (*Tagetes erecta* L.) cv. PusaNarangiGainda. *Advances in Life Sciences* 5(20); 9252-9258.
11. Khanna R P, Bohra M, Parul P and Nautiyal B P (2016) Studies on the effect of organic manures and PSB on vegetative and floral parameters of China Aster[*Callistephus chinensis* (L.) Nees] cv. Kamini under mid hills region of Himalaya. *The Bioscan* 11(4); 2707-2710.
12. SupriyaBordoloi and Madhumita Choudhury Talukdar. 2019. Effect of Organic Inputs on Growth and Flowering Attributes of Chrysanthemum cv. Snowball. *International Journal of Current Microbiology and Applied Sciences*. 8(12); 2189-2196.
13. Kumar M, Vishnupriya J and Subramaniyan P (2020) Influence of organic manure bio-fertilizers for growth and yield in Chrysanthemum Cv. Poornima White. *International Journal of Current Research* 12(11); 14929-14933.
14. Kumar Y and Vishwanath (2020) Effect of integrated nutrients management with plant growth retardant on growth and yield of African Marigold (*Tagetes erecta* L.) Cv. PusaNarangiGainda. *Plant Archive* 20; 340-345.

**Comment [H17]:** Ensure that the references comply with the spelling rules

15. Mittal R, Patel H C, Nayee D D and Sitapara H H (2010) Effect of integrated nutrient management on flowering, yield and vase life of African Marigold (*Tagetes erecta* L.) cv. Local under middle Gujarat agro-climatic conditions. *Asian Sciences* 5 (1); 22-24.
16. Pandey S K, Kumari S, Singh D, Singh V K, Prasad V M (2017) Effect of bio –fertilizers and organic manures on plant growth, flowering and tuber production of a Dahila (*Dahlia variabilis* L.) cv. S P Kamala. *International Journal of Pure and Applied Bioscience* 5 (2); 549-555.
17. Bohra M and Kumar A (2014) Studies on effect of organic manures and bio inoculants on vegetative and floral attributes of Chrysanthemum cv. Little Darling. *The Bioscan*; 9(3); 1007-1010.
18. Sudhagar R, Alexander R, Pamela E B and Kamalakannan S (2019) Effect of integrated nutrient management on the flower yield of African Marigold (*Tagetes erecta* L.) cv Local Orange. *International Journal of Emerging Technologies and innovative Research* 6(5); 284-289.
19. Parya C (2017) Effect of integrated plant nutrient system for Gerbera flower production under protected cultivation. *Journal of Applied Horticulture* 19 (2); 139-142.