

“STANDARDIZATION OF ROSE APPLE (*Syzygium jambos*) SYRUP WITH DIFFERENT CONCENTRATIONS OF SUGAR ”

ABSTRACT

The experiment was conducted on “Standardization of rose apple (*Syzygium jambos*) syrup with different concentrations of sugar” during the period November 2021 to February 2022 at Post Harvest Technology Laboratory, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences Prayagraj (Uttar Pradesh). The experiment was planned in completely randomized block design, seven treatments with three replication. The observations were recorded on various characteristics like acidity, TSS, pH, ascorbic acid content and organoleptic qualities. The results obtained from the experiment shows that the treatment T₆ (Rose apple juice + 55% sugar syrup) was found best in terms of the quality attribute as compared to other treatments. It shows the highest mean score in sensory evaluation like colour and appearance (8.85), Taste (8.6), Aroma (8.45), overall acceptability (8.7) and the chemical parameters like titratable acidity (0.458), TSS (64.66°Brix), pH (4.427) and ascorbic acid (26.73%). Therefore on the basis of above findings it is concluded that the treatment T₆ (Rose apple juice + 55% sugar syrup) is the most promising and best syrup in terms of their physico-chemical and organoleptic test of rose apple syrup.

Key words: Rose apple, syrup, sugar, citric acid, Physico-Chemical

INTRODUCTION

The Rose Apple (*Syzygium jambos*) is a blooming plant of Myrtaceae family. The Rose apple is native to East India and Malaya region and is widely cultivated throughout Southeast Asian countries and some parts of tropical for different purposes. It is grown for its sweet watery fruits and for its very beautiful foliage. The fruit can range in colour from white to greenish white to pale yellow. The edible fruit has a guava-like shape with four fleshy calyx lobes at the apex. The rose apple is more often used in jellies and jams or preserved combinations with other fruits of more pronounced flavour. They are delicious and highly refreshing beverages recommended for the manufacturing of like RTS drink, nectar, squash, crush, syrup and other value added products like jam, jelly, chutney, butter, sauce, etc. Value added products of rose apple can

attract both national and international market because there is always a great demand from the consumers all over the world for the new food products which are nutritious, therapeutic, delicious, and attractive in colour and appearance. An infusion of the fruit act as a diuretic. According to Chattopadhyay and Ghosh (1992) rose apple is fair source of iron. It has many medicinal values. This fruit contains many minerals such as calcium, iron and magnesium. Fever is thought to be reduced by a sweetened floral mixture. Forty three volatile constituents were identified by Wong (1996) in rose apple. The Vitamin C, phenolic content, flavinoids and numerous antioxidants are abundant in the pulp. It can help to increase immunity, control diabetes, aid digestion, improve teeth and bone health, avoid muscle cramps, treat diarrhoea and constipation, and is also beneficial to pregnant women. This fruit is highly perishable in nature. To reduce the loss, the fruit is used to make juices, jams, pickles and wine. The juices are loved by everyone, irrespective of their age group. It can give you a nutritious and healthy life. The whole plant has a variety of medicinal values, ranging from dermatological, digestive, head and throat to endocrine remedies. In India, this fruit syrup is considered a tonic for the brain and liver because of its rich source of vitamin C, vitamin A, fibre, calcium and iron content.

Materials and Methods

The rose apple fruits (matured, fresh, ripen and free from insects and diseases), glass bottles, sugar and muslin cloth from the local market. Citric acid is used to avoid the crystallization of sugar syrup and it can also act as a preservative. The work was conducted in the Post harvest laboratory, Department of Horticulture, SHUATS, Prayagraj, during November 2021 – February 2022. The instruments viz. Knife, trays, grinder, sieve, electronic balance, refractometer, digital pH meter etc. were used.

Well, ripe, fresh, firm, healthy and disease-free rose apple fruits were washed thoroughly with running tap water and rinsed with distilled water. The pulp was extracted manually and juice has separated from it. The calculated amount of sugar syrup was made separately and added to rose apple juice according to different treatments viz. T₁ (Rose apple juice + 30% sugar syrup), T₂ (Rose apple juice + 35% sugar syrup), T₃ (Rose apple juice + 40% sugar syrup), T₄ (Rose apple juice + 45% sugar syrup), T₅ (Rose apple juice +50% sugar syrup), T₆ (Rose apple juice +55% sugar syrup), T₇ (Rose apple juice + 60% sugar syrup) and heated it for just 5 minutes over a gas burner for proper dilution. The citric acid was added at 0.1% to avoid crystallization. The rose apple syrup was filled in cleaned, sterilized bottles and sealed with caps. The bottles were then labeled and stored in cold storage condition (10°C) for up to 90 days and Physico-chemical parameters were analyzed. The rose apple syrup were analyzed for Titratable acidity by titrate known volume of aliquot against 0.1 N NaOH and expressed as mg/100g and TSS (° B) by using hand refractometer (Erma, japan). The pH was determined by using digital pH meter and Ascorbic acid was determined by 2, 6 - dichlorophenol indophenol titration method. Prepared rose apple syrup were evaluated for sensory attributed like colour and appearance, taste, aroma and overall acceptability using 9- point hedonic scale.

RESULTS AND DISCUSSION

The effect of sugar concentration in cold storage condition on the quality of rose apple syrup was determined by checking physico-chemical parameters viz. T.S.S., pH, titratable acidity, ascorbic acid content and organoleptic scores. The main objective of this experiment was to find out the suitable concentration of sugar syrup on rose apple syrup and the storage quality of rose apple syrup.

TOTAL SOLUBLE SOLIDS (T.S.S.) (° Brix)

TSS of rose apple syrup was recorded with a hand refractometer (Srivasthava and Kumar 2000).

The TSS of rose apple syrup found to be increased with the increase in the proportion of sugar syrup and storage period. The results were 57.66, 64.66 and 67.00 for the treatments T₅ (Rose apple juice + 50% sugar syrup), T₆ (Rose apple juice + 55% sugar syrup) and T₇ (Rose apple juice + 60% sugar syrup). The highest mean value was observed in T₇ (67.66° Brix) while minimum mean score recorded in T₁ (38.16° Brix). Similar observations were recorded by Marimuthu M. and Thirumaran A. S. (2000) and in jamun syrup by (Kannan and Thirumaran, 2001).

ASCORBIC ACID

A significant decrease of ascorbic acid was observed during 90 days of storage period. The highest mean value was observed in the treatment T₁ (rose apple juice + 30% sugar syrup) (37.64) which was superior to all other treatments, followed by the treatment T₂ (rose apple juice + 35% sugar syrup) (35.1) while the minimum was recorded in the treatment T₇ (rose apple juice + 60% sugar syrup) (24.74). Ascorbic acid is very sensitive to light, heat and oxygen. Hence, it might have been destroyed due to heat at the time of preparation and subsequently during storage period. Similar results of ascorbic acid content were found in rose apple squash (Basavaraja, P.S, 2005) and Pattar *et al.*, (2013) .

pH

The pH shows a slight increase with the increase during the storage period. The highest mean value was observed in treatment T₇ (Rose apple juice + 60% sugar syrup) (4.44) while the minimum value was observed in treatment T₁ (Rose apple juice + 30% sugar syrup) (4.29). Initially, the results were 4.373, 4.403 and 4.42 for the treatment T₅ (Rose apple juice + 50% sugar syrup), T₆ (Rose apple juice + 55% sugar syrup) and T₇ (Rose apple juice + 60% sugar syrup) which got increased during 90 days of storage period as 4.42, 4.46 and 4.48. Similar results were obtained on aonla syrup by Reddy and chhikksubbanna, (2009) and Basavaraja P.S, (2005) in rose apple syrup.

TITRATABLE ACIDITY

Acids present in different quantities can give desirable taste to the syrup. The titratable acidity of rose apple syrup varied significantly with different treatments and the storage period. Initially, the results were 0.597, 0.533 and 0.426 for the treatments T₅ (Rose apple juice + 50% sugar syrup), T₆ (Rose apple juice + 55% sugar syrup), T₇ (Rose apple juice + 60% sugar syrup) and it got decreased according to the storage period of 90 days as 0.426, 0.341 and 0.298. The highest mean value was observed in T₁ (Rose apple juice + 30% sugar syrup) (0.805) while the minimum was recorded in T₇ (0.399). Similar observations were found by Sravanthi *et al.*, (2014) studied on the custard apple squash and Nayak *et al.*, (2011) during the study of aonla syrup.

SENSORY EVALUATION

In terms of colour and appearance, The treatment T₆ (Rose apple juice + 55% sugar syrup) recorded the highest score (8.85) mean value, which was significantly superior to all treatments, followed by the treatment T₅ (Rose apple juice + 50% sugar syrup) (8.2) and treatment T₇ (Rose apple juice + 60% sugar syrup) (7.65). The lowest (7.0) mean value for colour and appearance was observed in treatment T₁ (Rose apple juice + 30% sugar syrup) followed by treatment T₂ (Rose apple juice + 35% sugar syrup). And it is also observed from the data that the colour of the syrup increases as the proportion of the sugar syrup increase in rose apple syrup. The decreasing trend in the organoleptic score of colour and appearance for rose apple syrup was noticed irrespective of the different recipes and the storage period.

In terms of taste of the syrup varies significantly during the storage irrespective of the treatments. The highest mean score (8.6) was observed in treatment T₆ (Rose apple juice + 55% sugar syrup) which was significantly superior to all the other treatments and followed by T₇ (Rose apple juice + 60% sugar syrup) (8.15) and treatment T₅ (Rose apple juice + 50% sugar syrup) (7.95). The lowest mean score of 6.2 was observed in treatment T₁ (Rose apple juice + 30% sugar syrup) and followed by treatment T₂ (Rose apple juice + 35% sugar syrup) at 6.9. It is also observed that the taste of the syrup got decreased during the storage from the 0th day to 90 days.

Terms of the aroma of the rose apple syrup revealed that the value for aroma is influenced significantly due to recipe treatments as well as the storage period. The aroma of the syrup varies significantly during the storage irrespective of the treatments. The highest mean score (8.2) was observed in treatment T₆ (Rose apple juice + 55% sugar syrup) which was significantly superior to all the other treatments and followed by treatment T₅ (7.40) (Rose apple juice + 50% sugar syrup) and T₇ (7.4) (Rose apple juice + 60% sugar syrup). The lowest mean score of 6.05 was observed in treatment T₁ (Rose apple juice + 30% sugar syrup) and followed by treatment T₂

(6.35) (Rose apple juice + 35% sugar syrup). It is also observed that the taste of the syrup got decreased during the storage of 90 days.

The organoleptic score for the overall acceptability of the syrup, prepared by seven different treatments was statistically significant. The treatment T₆ (Rose apple juice + 55% sugar syrup) recorded the highest mean (8.7) score for overall acceptability which was significantly superior to all the other treatments followed by T₅ (Rose apple juice+ 50% sugar syrup) with 7.85 and T₇ (Rose apple juice+ 60% sugar syrup) while the lowest score was achieved by treatment T₁ (Rose apple juice + 30% sugar syrup) 5.6 and T₂ (Rose apple juice + 35% sugar syrup) with the score 6.45. The organoleptic score for the overall acceptability of rose apple syrup was reduced during the storage periods (Kannan *et al.*, (2004)) and Roy and Singh (1979) in bael beverages.

Fig. 1 : Rose apple syrup during storage (immediately after preparation)

Fig. 2 : Rose apple syrup after 90 days of storage in refrigerator

Table 1: Effects of sugar level in cold storage condition on the colour and appearance, taste, aroma of rose apple syrup.

TREATMENTS	COLOUR AND APPEARENCE					TASTE				
	0 days	30 days	60 days	90 days	Mean	0 days	30 days	60 days	90 days	Mean
T ₁ (Rose apple juice +30% sugar syrup)	7.4	7.2	7.0	6.4	7.0	6.6	6.4	6.0	5.8	6.2
T ₂ (Rose apple juice +35% sugar syrup)	7.6	7.4	7.0	6.8	7.2	7.4	7.2	6.8	6.2	6.9
T ₃ (Rose apple juice +40% sugar syrup)	7.8	7.6	7.2	6.8	7.35	7.8	7.6	7.2	6.6	7.3
T ₄ (Rose apple juice +45% sugar syrup)	8.0	7.6	7.2	7.0	7.45	8.0	7.8	7.6	7.4	7.7
T ₅ (Rose apple juice +50% sugar syrup)	8.8	8.8	7.8	7.4	8.2	8.2	8.2	7.8	7.6	7.95
T ₆ (Rose apple juice +55% sugar syrup)	9.0	9.0	8.8	8.6	8.85	9.0	9.0	8.4	8.0	8.6
T ₇ (Rose apple juice +60% sugar syrup)	7.8	7.8	7.6	7.4	7.65	8.4	8.4	8.0	7.8	8.15
Mean	8.00	7.9	7.5	7.2	7.6	7.91	7.82	7.35	6.95	7.5
F ratio	S	S	S	S		S	S	S	S	
S.E.M	0.03	0.20	0.17	0.20		0.18	0.20	0.18	0.20	
CD 5%	0.11	0.59	0.50	0.60		0.53	0.59	0.53	0.59	

Graph 1: Graphical representation of colour and appearance , taste of rose apple syrup in cold storage condition

Table 2: Effects of sugar level in cold storage condition on the aroma and overall acceptability of rose apple syrup.

TREATMENTS	AROMA					OVERALL ACCEPTABILITY				
	0 days	30 days	60 days	90 days	Mean	0 days	30 days	60 days	90 days	Mean
T ₁ (Rose apple juice + 30% sugar syrup)	6.4	6.2	6.0	5.6	6.05	6.0	5.8	5.4	5.2	5.6
T ₂ (Rose apple juice + 35% sugar syrup)	6.8	6.4	6.2	6.0	6.35	6.8	6.6	6.4	6.0	6.45
T ₃ (Rose apple juice + 40% sugar syrup)	7.2	7.0	6.4	6.2	6.7	7.2	7.0	6.6	6.2	6.75
T ₄ (Rose apple juice + 45% sugar syrup)	6.8	6.6	6.4	6.4	6.55	7.6	7.4	6.8	6.4	7.05
T ₅ (Rose apple juice + 50% sugar syrup)	7.4	7.6	7.4	7.2	7.4	8.2	8.2	7.8	7.2	7.85
T ₆ (Rose apple juice + 55% sugar syrup)	8.6	8.6	8.4	8.2	8.45	9.0	8.8	8.6	8.4	8.7
T ₇ (Rose apple juice + 60% sugar syrup)	7.6	7.4	7.2	7.0	7.3	8.2	8.0	7.6	7.4	7.8
Mean	7.25	7.1	6.8	6.6	6.9	7.48	7.4	7.02	6.6	7.1
F ratio	S	S	S	S		S	S	S	S	
S.E.M	0.17	0.22	0.21	0.18		0.17	0.18	0.23	0.20	
CD 5%	0.69	0.63	0.61	0.53		0.51	0.53	0.67	0.59	

Graph 2: Graphical representation of aroma and overall acceptability of rose apple syrup in cold storage condition

Table 3: Effects of sugar level in cold storage condition on the ascorbic acid content and total soluble solids of rose apple syrup.

TREATMENT	ASCORBIC ACID					TOTAL SOLUBLE SOLIDS				
	0 days	30 days	60 days	90 days	Mean	0 days	30 days	60 days	90 days	Mean
T ₁ (Rose apple juice +30% sugar syrup)	41.68	38.19	36.62	34.10	37.64	37.66	38.00	38.00	39.00	38.16
T ₂ (Rose apple juice +35% sugar syrup)	39.48	35.29	33.67	31.96	35.1	40.33	40.66	41.33	42.00	41.08
T ₃ (Rose apple juice +40% sugar syrup)	37.27	33.62	32.28	31.06	33.55	45.00	45.33	45.66	46.00	45.49
T ₄ (Rose apple juice +45% sugar syrup)	35.68	31.96	30.21	28.67	31.63	47.00	47.33	50.66	51.00	48.99
T ₅ (Rose apple juice +50% sugar syrup)	33.62	30.06	28.96	27.78	30.10	57.66	58.00	59.33	59.66	58.99
T ₆ (Rose apple juice +55% sugar syrup)	30.58	27.88	25.28	23.21	26.73	64.66	65.00	65.33	66.00	65.24
T ₇ (Rose apple juice +60% sugar syrup)	27.62	25.28	23.78	22.29	24.74	67.00	67.33	68.00	68.33	67.66
Mean	34.99	31.89	30.11	28.43	31.3	51.33	51.85	51.37	51.85	52.3
F ratio	S	S	S	S		S	S	S	S	
S.E.M	0.003	0.002	0.002	0.003		0.25	0.25	0.30	0.21	
CD 5%	0.011	0.008	0.007	0.009		0.76	0.76	0.93	0.66	

Graph 3: Graphical representation of ascorbic acid content and total soluble solids of rose apple syrup in cold storage condition

Table 4: Effects of sugar level in cold storage condition on the pH and titratable acidity of rose apple syrup.

TREATMENT	pH					TITRATABLE ACIDITY				
	0 days	30 days	60 days	90 days	Mean	0 days	30 days	60 days	90 days	Mean
T ₁ (Rose apple juice +30% sugar syrup)	4.28	4.286	4.29	4.32	4.294	0.874	0.832	0.768	0.746	0.805
T ₂ (Rose apple juice +35% sugar syrup)	4.30	4.311	4.32	4.34	4.317	0.832	0.789	0.731	0.661	0.753
T ₃ (Rose apple juice +40% sugar syrup)	4.316	4.336	4.34	4.36	4.332	0.810	0.746	0.661	0.618	0.708
T ₄ (Rose apple juice +45% sugar syrup)	4.35	4.366	4.38	4.40	4.374	0.746	0.682	0.597	0.533	0.639
T ₅ (Rose apple juice +50% sugar syrup)	4.373	4.39	4.40	4.42	4.395	0.682	0.597	0.505	0.426	0.552
T ₆ (Rose apple juice +55% sugar syrup)	4.403	4.416	4.43	4.46	4.427	0.597	0.533	0.362	0.341	0.458
T ₇ (Rose apple juice +60% sugar syrup)	4.42	4.44	4.45	4.48	4.44	0.533	0.426	0.341	0.298	0.399
Mean	4.348	4.363	4.375	4.397	4.36	0.724	0.657	0.552	0.517	0.61
F ratio	S	S	S	S		S	S	S	S	
S.E.M	0.002	0.002	0.002	0.002		0.02	0.03	0.019	0.021	
CD 5%	0.008	0.008	0.006	0.006		0.07	0.091	0.059	0.064	

Graph 4: Graphical representation of pH and titratable acidity of rose apple syrup in cold storage condition

CONCLUSION

From the present investigation, it could be concluded that all the recipes of rose apple syrup were found to be organoleptically acceptable not only at the time of preparation but also throughout the storage period of 3 months under cold storage condition. The syrup prepared from rose apple fruit pulp was significantly superior concerning the overall acceptability of the product. The maximum overall acceptability at the day of preparation was shown by the treatment T₆ (9.0) and the minimum by the treatment T₁ (6.0). At the 90th day the maximum overall acceptability was observed in T₁ (5.2) and the maximum was recorded on T₆ (8.4). Therefore the syrup recipe treatment T₆ i.e. rose apple juice + 55% sugar syrup was found to be the best recipe for rose apple syrup with the highest organoleptic score for Colour, Taste, Aroma and Overall acceptability.

REFERENCES

- Basavraja, P.S. (2005)** Value added product from rose apple (*Syzygium jambos* Alston) M.Sc.(Hort.) Thesis, university of agriculture Science, Bengaluru.
- Bolarin, F. M. (2009)** Rose apple fruit: It prospects for juice and wine production , *European Journal of Educational and Development Psychology* Vol.4, No.2, pp.25-32, April 2016.
- Hamaran, M and Amutha S. (2007)** Effect of total soluble solids and CO₂ pressure on physico- chemical and sensory qualities of carbonated banana and sapota beverages. *Journal of Food Science and Technology*, **44**(2): 178-182.
- Jain, S.P., Tripathi, V.K. and Ram, H.B. 1984.** Effect of storage conditions on the keeping quality of fruit squash. I. Studies on storage behaviour of orange, lemon and bael squash. *Indian Food Packer* **38**(5): 33-39.
- Kannan, S. and Thirumaran, A.S. (2001)** Studies on storage life of jamun products. *Indian Food Packer* **55**(6): 125-127.
- Kannan and Thirumaran (2004)** Studies on storage life of jamun (*Syzygium cumini* Rom) fruit products. *J. Food Sci. Technol.*, **41**(2): 186-188.
- Khurdiya, D. S., Roy, S. K. (1985)** Processing of jamun(*syzygium cumini* Linn.) fruits into a ready to serve beverage.J, *Food Sci. Tech.* **22**(3): 27-30.

Korgaonkar M., Pujari, K.H. and Morgaonkar, V.R. (2015) Effect of different acidity levels of snap melons syrup on chemical and sensory properties. *IOSR Journal of Agriculture and Veterinary Science*, **8** (4):01-03.

Kotechua, P.M., Kadam, S.S. (2003) Preparation of RTS, syrup and Concentrate from tamarind. *J. Food sci. Tech.*, **40**(1) :76-79.

Marimuthu, M. and Thirumaran, A. S. (2000) . Utilization of jamun juice, squash and syrup. *Beverage and Food world*, **27**(8): 42-46.

Nayak P., Bhatt D.K., Shukla, D.K. and Tandon, D.K. (2011) Evaluation of aonla (*Phyllanthus emblica G.*) segments- insyrup prepared from stored fruits. *Research journal of agriculture science*, **43** (2): 252-257.

Pal R. and Sethi, V. 1993. Chemical Changes during storage of Kagzi lime syrup at different temperatures. *Maharashtra Journal of Horticulture* **7**(2): 83-88.

Pattar A., Kukanoor L., chavan M., Prabhuling G. (2013) Preparation and storage of tamarind syrup. *Beverage and food world*, **40** (8):61-63.

Prasad, R.N. and Mali, P.C. (2000) Changes in physico- chemical characteristics of pomegranate squash during storage. *Indian Journal of Horticulture*, **57**(1): 18-20.

Reddy, and Chikkasubbanna, v. (2009) studied the storage behaviour of amla syrup. Syrup prepared with 25 per cent pulp and 70 °Brix total soluble solids was found to be the best recipe. *The Asian Journal of Horticulture*. **4**(1) : 5-9

Roy, S.K. and Singh, R.N. (1979) Studies on utilization of bael fruit (*Aegle marmelos* Correa.) for processing III. Preparation and preservation of bael fruit. *Indian Food Packer* **33**(5): 9-14.

Salvi, M.J. (1991) Studies on processing qualities of mango (*Mangifera indica* L.) cashew (*Anacardium occidentale* L.) and karonda (*Carissa carandas* L.) fruits. M.Sc. (Agri.) thesis.

Singh, A.K. (1985) Studies on preparation and storage of guava (*Psidium guajava* L.) beverages M.Sc. Thesis, Narendra Deva University of Agriculture and Technology, Faizabad, U.P., India.

Sravanthi T., Jayasimha R.D., Waghrey k. (2014) studies on preparation and processing of custard apple (*Annona squamosa* L.) pulp, *International J. plant, animal and environmental sciences*, **4**(3):676-682.

Srinivas, K. N., Revanna, M. L. and Jamuna, K. V. (2007) Standardization of recipes for production of squash in pomegranate (*Punica granatum* L.) cv. Ganesh and Mridula. *J. Dairying Foods and Hort. Sci.*, **26**(2):133-136.

Sujana A. (2006) Studies on preservation of custard apple pulp for value addition. M.Sc (Agri.) Thesis, Dr. PDKV, Akola (M.S.), India.

Thakur N.S., Girish, D. S. and Joshi, V.K. (2013) Development of wild pomogranate aril- in-syrup and its quality evaluation during storage. *International journal food fermentation technology*, **3**(1):33-40.

Tiwari S., Chopra, C.S. and Kuari, S. (2005) Studies on preservation of aonla (*Emblica officinalis* Gaertn.) fruit pulp. *Beverage & Food World* **32**(12): 21-23.