

# Original Research Article

## Yield enhancement through a low-cost irrigation system in Groundnut

### ABSTRACT

The existing irrigation system namely drip and sprinkler adds cost to production. So, an alternate low-cost irrigation system (rain hose irrigation) was compared with other irrigation systems. The study was proposed to find out the optimum soil moisture by selecting a proper irrigation system that supports optimum soil moisture management, enhanced productivity, high water use efficiency, and also involves less cost on production. The treatment comprises different methods of irrigation (drip, sprinkler, rain hose, and BBF) and deficit irrigation (100%, 125%, and 75% of PET) was tested. The results have shown that higher pod yield (3086 kg/ha) and high water use efficiency (11.2 kg/ha.mm) were attained in drip + 100% of PET and drip + 75% of PET. The high B: C ratio was attained in rain hose + 100% of PET and hence, considered as the low-cost irrigation technology for higher profit.

**Keywords:** Groundnut, alternate irrigation system, rain hose irrigation, B: C ratio, low-cost irrigation system, higher profit

### I. INTRODUCTION

Groundnut (*Arachis hypogaea* L.) is an important legume crop that belongs to the family of Leguminosae. Groundnut is cultivated mainly in the *Kharif* season. In some states, it is also cultivated in the *rabi* and *spring* seasons as an irrigated crop. The crop is especially valued for its high amount of oil (43-55%) and protein (25-28%) content.

India has been the world's leading producer of groundnut in an area of 6.01 m ha, having an annual production of 10.2 m t with annual average productivity of 1703 kg ha<sup>-1</sup> (Indiastat 2020-21). It is the 6<sup>th</sup> most important oil seed crop after sunflower, sesame, etc. In Tamil Nadu, it is grown in an area of 4.09 lakh hectares, having an annual production of 1.023 m t with annual productivity of 2500 kg ha<sup>-1</sup> (Indiastat 2020-21). The water required by groundnut to complete its life cycle is around 500-700 mm (Ahmad *et al.*, 1998), but the needed quantity of water varies from crop's several stages. Among all the agronomic factors, water management stands second which contributes yield next to fertilizer application of around 27%.

Out of the total cost of production, more than 40% of the cost involved seed input and it is not reflected in yield. The seed cost has not contributed to the yield factor because of higher mortality by unscientific soil moisture maintenance. The irrigation technology which has to support optimum soil

**Comment [a1]:** Please refer to the unit kg grain/m or what means?

moisture maintenance for the better establishment of crops to maintain the optimum plant population and support for high productivity by reducing mortality is essential.

Adopting appropriate irrigation management technologies which significantly influences the water use efficiency and improves productivity in a cost-effective manner. Those irrigation management technologies maximize productivity by providing adequate moisture to the crop during the crop growth stages. (Soni *et al.*,2017).

In comparison with traditional irrigation practices, micro-irrigation systems play a crucial role in saving agricultural water use. The existing micro irrigation system namely drip and sprinkler adds cost to production and reduces profitability. The system which supports optimum soil moisture management and also involves less cost on production to be studied

With this background, an alternate method of irrigation is recently available in the market (rain hose system of irrigation) was considered as one of the treatments and compared with the existing micro-irrigation systems with the objective

- To find out the optimum soil moisture to avoid disease incidence
- To find out the best water management approach for higher productivity and higher profitability.

## II. MATERIALS AND METHODS

### A. Experimental site

The field experiment was carried out during *rab* season at Agricultural College and Research Institute, Madurai, Tamil Nadu, India. The initial soil sample has been taken for analyzing the nutrient content of the soil. The initial status of soil has low available nitrogen (246.4 kg/ha), medium available phosphorus (11.8 kg/ha), and high available potassium (478 kg/ha) respectively. The soil pH and EC values were 7.55 and 0.18 dsm<sup>1</sup>. The soil texture of the experimental field was sandy loam having a moderate infiltration rate of 2.25 cm/h.

Comment [a2]: format

### B. Selected variety

The variety selected for the experiment was VRI-8 (Virudhachalam 8), which was released by Regional Research Station (TNAU), Virudhachalam in the year 2016. It is a Spanish bunch type variety, with medium robust pods and rose-colored testa. It takes 105-110 days to reach full maturity.

### C. Design of experiment

The experimental design was a strip plot design which includes one main factor namely drip, sprinkler, rain hose, and broad bed and furrow (BBF) methods of irrigation and the sub-factor includes deficit irrigation at levels of 100%, 125%, and 75% of PET and it was replicated **thrice**.

Comment [a3]: three

#### D. Methodology involved

The irrigation was given to the crop once in three days intervals, and the quantity of water has been decided based on the pan evaporation method.

$$ET_O = E_{pan} \times K_p$$

Where,

$ET_O$  - Potential Evapotranspiration

$E_{pan}$  - Pan Evaporation (mm)

$K_p$  - Pan coefficient

$ET_O$  is the Potential Evapotranspiration; it was calculated by using the Class A pan evaporimeter.

The quantity of water that needs to be given was calculated at different levels of 100%, 125%, and 75% of PET by using the pan evaporation method, which was given to the crop through the drip, sprinkler, rain hose, and BBF method of irrigation systems. The application of the calculated quantity of water through different irrigation systems will be the same but the time taken to deliver the water has varied according to the irrigation system.

#### E. Water Use Efficiency

Water Use Efficiency (WUE) was a term that helps to evaluate deficit irrigation strategies. It is defined as the ratio of yield obtained to the volume of water utilized by the crop. (Kori *et al.*, 2017)

Yield ( $Y_a$ )

WP =

Volume of water ( $ET_a$ )

#### F. Economics

The market prevailing rates of varied inputs like seeds, fertilizers, and wages for the labourers and the available market rate of the produce at the local paid by the government have been considered while deriving the economics. The economics of the conventional method and micro irrigation systems also had been derived to calculate the net returns and B: C ratio.

### III. Results and discussion

#### Yield obtained

Among the irrigation combinations, the highest pod yield was obtained in drip irrigation + 100% of PET (3086 kg/ha) and was followed by rain hose irrigation + 100% of PET (2922 kg/ha), and it was on par with drip irrigation + 75% of PET (2856 kg/ha). The lowest yield was observed in the BBF + 125% of PET (1926 kg/ha). These findings were similar to the results given by Ranjitha *et al.*, 2018. The increased pod yield in drip irrigation + 100% of PET was due to the maintenance of optimum moisture throughout the growing season directly to the root zone which has a significant influence on the yield attributes like the number of pods/plant and providing adequate frequency of irrigation has kept the soil in optimum moisture condition with reduced disease incidence led to optimum plant population, when compared to other treatment combinations. These findings were similar to the results given by Ranjitha *et al.*, 2018. Hence, the contribution from an increased number of pods/plants, lower disease incidence, and sustained plant population boosted the drip + 100% PET to attain the highest yield when compared to all other treatments. The drip irrigation method had greatly influenced the yield parameters like pod and haulm by Mathuika *et al.*, (2019). The obtained results were similar to the reports given by (Sezen *et al.*, 2019).

**Table.1. Effect of different methods of irrigation and deficit irrigation levels on Pod yield**

Treatments	Deficit irrigation levels			Mean
	100%	125%	75%	
<b>Drip</b>	3086	2440	2856	<b>2794</b>
<b>Sprinkler</b>	2598	2293	2442	<b>2444</b>
<b>Rain hose</b>	2922	2346	2718	<b>2662</b>
<b>BBF</b>	2037	1926	1987	<b>1983</b>
<b>Mean</b>	<b>2582</b>	<b>2151</b>	<b>2435</b>	<b>2389</b>
	<b>I</b>	<b>S</b>	<b>I x S</b>	<b>S x I</b>
<b>S. Ed</b>	<b>31.92</b>	<b>29.98</b>	<b>46.97</b>	<b>43.54</b>
<b>CD (0.05)</b>	<b>65.18</b>	<b>62.07</b>	<b>121.92</b>	<b>118.14</b>

**Fig.1. Impact of different methods of irrigation and deficit irrigation levels on Pod yield**

#### **Water Use Efficiency**

Generally, getting high yield by lowering the quantity of water use would have higher water use efficiency, in that case drip irrigation had significantly increased the water use efficiency. Drip irrigation +75% of PET (11.2 kg/ha.mm) has been obtained with a result of the highest water use efficiency and which has been closely followed by rain hose + 75% of PET (10.7 kg/ha/mm) and the following treatment after that was sprinkler +75% of PET (9.6 kg/ha.mm) and the lower water use efficiency was attained by BBF + 125% of PET (4.5 kg/ha.mm). The obtained results were similar to the results by (Biswas *et al.*, 2016). While using drip irrigation, enhancement of water use efficiency has been attained by avoiding percolation, runoff, and soil evaporation(El-Habbashas *et al.*, 2015). Drip + 75% of PET has given the higher water use efficiency, as the quantity of water has been utilized effectively through direct root zone application to the crop with reduced evaporation losses decreased the water use which attained a high pod yield per mm of water applied(El-Habbashas *et al.*, 2015).

**Table.2. Effect of different methods of irrigation and deficit irrigation levels on**

Treatments	Deficit irrigation levels			Mean
	100%	125%	75%	
<b>Drip</b>	9.1	5.7	11.2	<b>8.7</b>
<b>Sprinkler</b>	7.6	5.4	9.6	<b>7.5</b>
<b>Rain hose</b>	8.6	5.5	10.7	<b>8.3</b>
<b>BBF</b>	6.0	4.5	7.8	<b>6.1</b>
<b>Mean</b>	<b>7.7</b>	<b>5.2</b>	<b>9.6</b>	<b>6.8</b>
	<b>I</b>	<b>S</b>	<b>I x S</b>	<b>S x I</b>
<b>S. Ed</b>	<b>0.046</b>	<b>0.058</b>	<b>0.050</b>	<b>0.047</b>
<b>CD (0.05)</b>	<b>0.25</b>	<b>0.27</b>	<b>0.39</b>	<b>0.35</b>

**Water Use Efficiency**

**Fig.2. Impact of different methods of irrigation and deficit irrigation levels on water use efficiency (WUE)**

## Economics

The cost of cultivation, gross return, net return, and benefit-cost ratio were briefly shown in the table. Among the irrigation management practices, drip irrigation + 100% of PET (Rs.2,25,429.79/ha) had the highest net return, which was closely followed by rain hose irrigation + 100% of PET (Rs.2,15,506.00/ha) and followed by drip irrigation at 75% of PET (Rs.2,03,390.29/ha), and the lower value of net return was attained in BBF + 125% of PET (Rs.1,26,087.50/ha). The highest net return in drip irrigation + 100% of PET was also attributed to the highest gross return (Rs. 2,85,043.50/ha). Though drip irrigation + 100% of PET attributed to the highest net return, rain hose irrigation + 100% of PET records the highest B: C ratio (5.0) as its cost of cultivation was low when compared to drip installation. The installment cost has been higher in drip irrigation system. The B:C ratio was closely followed by drip+100% of PET (4.8) and by rain hose irrigation +75% of PET (4.6) and the lowest net return and the B: C ratio were obtained in the BBF+ 125% of PET (3.4). The B: C ratio of rain hose + 100% of PET was high due to less installment cost when compared to other irrigation systems and the increased pod yield.

**Table.3. The Cost of Cultivation, Gross Return, Net Return and B: C ratio in the treatment combinations given below.**

Treatment Combinations	Cost of cultivation	Gross return	Net return	B: c ratio
DRIP + 100% of PET	59614	285043.50	225429.79	4.8
DRIP + 125% of PET	59614	226179.00	166565.29	3.8
DRIP + 75% of PET	59614	263004.00	203390.29	4.4
SPRINKLER +100% of PET	61253	240243.00	178990.00	3.9
SPRINKLER + 125% of PET	61253	213447.00	152194.00	3.5
SPRINKLER +75% of PET	61253	225279.00	164026.00	3.7
RAIN HOSE + 100% of PET	54353	269859.00	215506.00	5.0
RAIN HOSE + 125% of PET	54353	217735.50	163382.50	4.0
RAIN HOSE + 75% of PET	54353	250600.50	196247.50	4.6
BBF + 100% of PET	52603	189292.50	136689.50	3.6
BBF + 125% of PET	52603	178690.50	126087.50	3.4
BBF + 75% of PET	52603	183718.50	131115.50	3.5

**Fig.3. Impact of different methods of irrigation and deficit irrigation levels on Economics**

#### **IV. CONCLUSION**

The higher productivity (3086 kg/ha), high water use efficiency(11.2 kg/ha.mm) was achieved in drip + 100% of PET and drip + 75% of PET. However, the higher pod yield has not resulted in higher profitability.Hence, the low-cost irrigation technology (rain hose system of irrigation) with optimum moisture level has attained the highest B: C ratio (5.0). Thus,the experimental result has been concluded that the low-cost irrigation technology i.e., rain hose + 100% of PET has given a better performance when compared to all other irrigation systems and farmer's practice.

## V. REFERENCE

1. Ahmad A, Kirda C, Moutonnet P, Hera C, and DR Nielsen. 1998. "Yield response of groundnut grown under rainfed and irrigated conditions." *Developments in Plant and Soil Sciences* 84:128-135.
2. Biswas, SK, AR Akanda, MS Rahman, and MA Hossain. 2016. "Effect of drip irrigation and mulching on yield, water-use efficiency and economics of tomato." *Plant, Soil and Environment* 61 (3):97-102.
3. El-Habbashas F, Okasha EM, Abdelraouf RE, and ASH Mohammed. 2015. "Effect of pressured irrigation systems, deficit irrigation and fertigation rates on yield, quality and water use efficiency of groundnut." *International Journal of chemtech Research* 7 (4):1751-1764.
4. Kori, SM, Soothar C, Qureshi AL, and Shaharyar Hassan Ashrafani. 2017. "Effect of different irrigation methods on water use efficiency and yield of maize crop." *Pakistan Journal of Agriculture, Agricultural Engineering and Veterinary Sciences* 33 (1):54-65.
5. Mathukia RK, Chhodavadia SK, Sagarka BK, Bhalu VB, "Summer Groundnut (*Arachis hypogaea* L.) Response to Micro-irrigation and Transient Water Stress ". *Research and Review Journal of Crop Science and Technology*. 2019; 8(2): 16-20p.
6. Ranjitha, P Sri, Ramulu V, Jayasree G, and Narender Reddy S. 2018. "Growth, yield and water use efficiency of groundnut under drip and surface furrow irrigation." *International Journal of Current Microbiology and Applied Sciences* 7 (09):1371-1376.
7. Sezen, S Metin, Seral Yucel, Servet Tekin, and Mehmet Yildiz. 2019. "Determination of optimum irrigation and effect of deficit irrigation strategies on yield and disease rate of peanut irrigated with drip system in Eastern Mediterranean." *Agricultural Water Management* 221:211-219.
8. Soni, Jeetendra Kumar, Asoka Raja N, and Ajaykumar R. 2017. "RESPONSE OF PHYSIOLOGICAL AND YIELD ATTRIBUTING PARAMETERS OF GROUNDNUT UNDER DRIP AND MICROSPRINKLER FERTIGATION." *An International Quarterly Journal of Life Sciences* 12:(1) 373-376.