

Plexiform neurofibroma of arm: a case report

Abstract

Plexiform neurofibroma (PNF) is a rare benign nerve tumor. It is usually considered pathognomonic of neurofibromatosis type 1 (NF1 or Von Recklinghausen disease). MRI is the examination of choice in the exploration of this pathology. It provides decisive arguments for the positive diagnosis, the evaluation of the prognosis and the follow-up of the lesions.

Histological confirmation is sometimes necessary, especially in cases other than NF1. We report the observation of a 22-year-old woman who consulted for a mass of the left arm.

Key words: Neurofibromatosis, plexiform, MRI, anatomy pathology

Introduction

Neurofibromas are benign tumors that develop from the roots and plexuses of spinal nerves. They can be uni or bilateral, sometimes stepped, superficial or deep. There are 3 types of neurofibromas: plexiform neurofibromas (nodular form - diffuse form), cutaneous and nodular. Plexiform neurofibromas correspond morphologically to a more or less long segment of tortuous dilatation of a nerve and its branches, producing the appearance of a bundle of string. Plexiform neurofibromas are pathognomonic of NF1. They are generally slow growing tumors. Their symptomatology is variable depending on their topography. We report the case of a 22-year-old woman who consulted for a mass of the left arm.

CASE REPORT

She was a 22-year-old woman with a father suffering from neurofibromatosis type I. Since childhood, she presented with a mass in the upper 1/3 of the anterolateral aspect of the left arm, which was painless, soft, and gradually increasing in size. For some time, the patient reported a sensation of heaviness with paresthesia of the left upper limb. Clinical examination revealed a mass on the anterolateral aspect of the arm with a soft, painless consistency, well limited and without inflammatory signs, measuring 10 cm in length (figure 1). Skin examination revealed the presence of multiple café au lait spots on the trunk and limbs (figure 2). In view of this very suggestive clinical picture and the patient's past history, the diagnosis of neurofibromatosis type I was evoked. An MRI of the left arm was performed in order to characterize the mass described above. It revealed the presence of an oblong tissue mass in the subcutaneous soft tissues of the upper-external aspect of the left arm, with regular contours, in T1 hypersignal (Figure 3) and T2 heterogeneous hypersignal (Figure 4), discreetly enhanced after injection of gadolinium (Figure 5), measuring 68x15mm extended over 114mm, respecting the muscular and vascular

structures and the bone opposite. The diagnosis of plexiform neurofibroma of the left arm was retained.



Figure 1: Profile view of the mass on the left arm.



Figure 2: Café au lait stains on the trunk.




Figure 3: MRI of the left arm, axial and coronal section in T1-weighted sequence without injection of gadolinium: lesion process of the superior-external aspect of the left arm well limited, with regular contours in T1 hypo signal.




Figure 4: MRI of the left arm, axial section in T2-weighted FSTSE sequence: lesion process of the superior-external aspect of the arm with regular T2 hypersignal contours.




Figure 5: MRI of the left arm, coronal section in T1-weighted sequence after gadolinium injection: the mass is discreetly enhanced after gadolinium injection.

Discussion

Plexiform neurofibromas are pathognomonic of NF1, affecting between 27 and 40% of NF1 subjects. They are recognized as congenital, slow growing and often remain asymptomatic. NPs can develop from superficial or deep nerve plexuses, essentially in the trunk and extremities, all spinal segments are affected. They are most often multiple, multi-radicular and tend to expand along the nerve sheaths. Often asymptomatic (40-60% of cases), they may reveal themselves by a varied clinical symptomatology, pain and/or radicular irritation, paresthesias, sensory-motor deficits, and even signs of spinal cord compression, which may vary according to the affected plexus or nerve trunk [1,2]. Clinically, they present as diffuse, patchy, bumpy, irregular tumors, covered with skin that is sometimes atrophic, flabby, wrinkled, and of the color of normal skin or sometimes bluish and pigmented. Their palpation is usually painless and gives a sensation of "bundle of strings" and inside the soft tumor, hard fibrous cords can be found on palpation [1,4].

Imaging is of great help in the diagnosis, allowing the characterization of lesions for a positive diagnosis, the search for possible associated lesions, the evaluation of the prognosis and the follow-up [2].

On ultrasonography, NPs appear as tortuous lobulated masses with a clustered, hypoechoic appearance, with well-defined contours oriented along the axis of the nerve trunk on longitudinal sections and characterized by the target sign on transverse sections with an echogenic center and a hypoechoic periphery, which correlates with the ultrastructure of neurofibromas with a fibrocollagenous center surrounded by a myxomatous periphery. Cystic formations can be seen within the mass as well as posterior enhancement encountered in 70% of cases. On color Doppler, there may be moderate, central or predominantly peripheral vascularization. Some may be poorly vascularized [3,8].

On CT, plexiform neurofibromas appear spontaneously isodense compared to muscle. They take up contrast only slightly, less than muscle, which gives them a hypodense appearance, differentiating them from schwannomas and neurofibrosarcomas.

The NP appears on magnetic resonance imaging sequences, which is the reference radiological examination, by a relative T1 hypo signal in relation to the muscle, T2 hyper signal and when it is voluminous, it may contain a central hypo signal producing a characteristic cocoon aspect. The enhancement is variable: central, diffuse, peripheral, in target [2,5,9].

The diagnosis of NFP is essentially anatomopathological, especially outside the context of neurofibromatosis type I. The risk of sarcomatous degeneration justifies, whenever technically possible, the removal of the lesion as completely as possible. Finally, these patients should be monitored clinically and radiologically by a multidisciplinary team, at least once a year and then on a regular basis, in order to assess any recurrence or malignant transformation [5].

Conclusion

Plexiform neurofibromas are rare benign tumors of the peripheral nerves. They are pathognomonic tumors of NF 1 which is a multi-systemic pathology with clinical and radiological polymorphism. MRI is the examination of choice in the exploration of this pathology. It provides decisive arguments for the positive diagnosis, the evaluation of the prognosis and the follow-up of the lesions.

References

1. Aloï FG, Massobrio R. Solitary plexiform neurofibroma. *Dermatologica*.1989; 179(2): 84-6.
2. Jacques, C., & Dietemann, J. L. (2005). *Imagerie de la neurofibromatose de type 1. Journal of Neuroradiology, 32(3), 180–197*.doi:10.1016/s0150-9861(05)83136-0
3. Lange F, Herlin C, Frison L et al. Prise en charge du neurofibrome plexiforme isolé de l'enfant: à propos de quatre observations. *Ann Chir Plast Esthet*. 2013; 58(6): 694-9.
4. Benaïssa L, El Kharras A, Bassou D et al. Masse cervicale gauche. *Feuil Radiol*. 2009; 49(3): 207-10.
5. Plexiform neurofibroma with neurofibromatosis type I/ von Recklinghausen's disease: A rare case report Pooja Poswal 1, Namita Bhutani 2, Sunil Arora 1, Raj Kumar 1
6. La neurofibromatose 1 (NF1) ou maladie de Von S.PinsonabdP.Wolkensteincd La Revue de Médecine Interne Volume 26, Issue 3, March 2005, Pages 196-2 Apr-Jun 2006;48(2):155-8.

7. Plexiform neurofibroma of the tongue: a case report of a child Enis Alpin Güneri¹, Ertap Akođlu, Semih Sütay, Kerim Ceryan, Ozgül Sađol, Uđur Pabuçcuođlu
Apr-Jun 2006;48(2):155-8.
8. Cross Lin J, Martel W. Cross-sectional imaging of peripheral nerve sheath tumors: charecteristics signs on CT, MRI and sonography. AJR. 2001; 176(1): 7-82.
9. IMAGERIE DE LA NEUROFIBROMATOSE DE TYPE 1 : 21/03/08-C. JACQUES,J.L. DIETEMANN Service de Radiologie 2, Hôpitaux Universitaires – Hôpital de Hautepierre Avenue Molière, 67098 Strasbourg Cedex.