

Original Research Article

“Effect of Spacing and Boron levels on growth and yield of Black gram (*Vigna Mungo* L.)”

Abstract

A field experiment was conducted aiming to study the influence of Spacing and Boron levels on growth and yield of black gram during **Zaid season 2022**. The treatments comprised of row spacing (25, 30, 35 cm) and boron at (1.0 ,1.5, 2.0 kg/ha). The results show that application of boron 2.0 kg/ha at 35 ×10 cm spacing has recorded significantly higher plant height (44.53 cm), number of branches per plant (9.27), number of nodules (35.53) **and** dry weight (10.44 g). However **higher** number of pods per plant (30.73), seeds per pod (6.40), test weight (42.48 g), seed yield (930.33 kg/ha) and stover yield (2040.33 kg/ha) gross return (Rs 58,610.79/ha), net return (Rs 34,311/ha) and B:C ratio (1.41) were recorded with the application of boron 2.0 kg/ha at 25×10 cm spacing. Therefore, the spacing of 25×10 cm and application of Boron @2.0 kg/ha could be a promising option for the yield enhancement in black gram.

Key words: *Boron, Black gram, Economics, Spacing, Yield.*

Introduction

Black gram (*Vigna Mungo* L.) is one of **the** important pulse **crops**. It is cultivated in many tropical and sub-tropical countries of **the** world. It is grown throughout India. Black gram is **a** widely grown grain legume and belongs to the family Fabaceae and assumes considerable importance from the point of food and nutritional security in the world. It is also known as urad bean, urad dal, **and** urad. It also acts **as a** cover crop and its deep root system protects the soil from erosion. Besides, this green fodder of **the** urad bean is very nutritive and is especially useful for mulch cattle. **U**rad bean being leguminous **can** fix atmospheric nitrogen and thus helps in restoring soil fertility (Shamim, M Z; Pandey, A 2017).

Black gram is short duration pulse crop which is grown in India area of 50.31 lakh hectare having the production of 32.84 lakh tons with productivity 652 kg/ha. (Ministry of Agriculture and Farmer Welfare, 2017-18). It accounts for 10 per cent of country's total pulse production (Gowda *et al.*, 2013). Black gram seed contain 55-60% carbohydrate, 22-24% protein and 1.0-1.3% of fat besides, phosphoric acid (H₃PO₄), being 5-10 times more than other pulses (Ali *et al.*, 2002) and sulphur containing amino acid (methionine

and cysteine). Dried black gram contains about 9.7% water, 23.4% protein, 1.0% fat, 57.3% carbohydrate and 3.8% fibre along with 154 mg Calcium, 9.1 mg Iron, 0.37 g riboflavin and 0.42 g Thiamine in each gram of black gram (Verma *et al.*, 2011) Due to cheaper protein source, it is designated as “poor man’s meat” (Aslam *et al.*, 2010). Because of their vital role in nutritional protection and soil development, pulses have been an integral part of sustainable agriculture since ancient times. (Tomar *et al.*, 2011).

The continuous cultivation of traditional low potential cultivars, the use of low seed rate, and improper agronomic practises were important reasons for the low average yield of black gram in farmers' fields. Among the various crop production constraints, the most important are appropriate varieties and crop spacing, both of which contribute significantly to black gram seed yield. Many research studies have revealed that most of the growth and yield contributing attributes were significantly and positively correlated with the seed yield of crop plants viz., black gram (Siddique *et al.*, 2006), soybean (Malik *et al.*, 2007) and sunflower (Vahedi *et al.*, 2010).

Plant density can have a major effect on the final yield of most of the legumes and the general response of yield to increasing population is well documented. To realize the maximum yield potential of black gram during the summer and rainy season, maintenance of optimum space made available to the individual plant is of prime importance. Row and plant spacing must be worked out to get desired spacing. The spacing requirement depends upon the growth behaviour of the genotype. So, it is required to maintain spacing for obtaining higher yield (P Veeramani, 2019). It is prime necessity to maintain optimum plant population by maintaining inter and intra row spacing properly. Maximum or minimum plant density may minimize yield of black gram causing physiological change in plant. Hence appropriate fertilizer dose with adequate plant population may increase crop yield of black gram. Similar results were noted by (Choudhary *et al.*, 2017).

Boron (B) is an essential non-metal micronutrient plays a decisive role in maintaining cell wall structure, cell division, membrane stability, nitrogen assimilation, sugar translocation, K⁺ transport, protein and sucrose synthesis, phenol, carbohydrate, nucleic acid and IAA metabolism (Singh *et al.*, 2014). Calcareous soils and those with low organic matter content are more prone to B deficiency (Dwivedi *et al.*, 1990 and Niaz *et al.*, 2013). Boron is retained in soils by adsorption on to minerals and humic particles and by forming insoluble precipitates (Goldberg and Glaubig, 1985). 4-5 lines related to boron importance for pulse crops. The potential B deficiency in soils of India was estimated as 33 per cent (Alloway 2008). It has been reported that boron deficiency limits reproductive growth. Havlin *et al.* (1999) documented that due to shortage of boron, flowering and fruit development were highly restricted. In recent times, management of boron is a challenging aspect as the optimum level of B application range is narrow S Sharmila (2020). With these considerations in mind, the current research aimed to examine the effect of spacing and boron on growth parameters and black gram yield.

MATERIALS AND METHODS

The experiment was conducted during the *Zaid* season 2022, at the CRF (Crop Research Farm), Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj, Uttar Pradesh. The CRF is located at 25° 30' 42''N latitude, 81° 60' 56'' E longitude and 98 m altitude above the mean sea level (MSL). This area is situated on the right side of the river Yamuna and by the opposite side of Prayagraj city. The soil of the experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 6.9), low in organic carbon (0.112 %), available N (278.93 kg/ha), available P (10.8 kg/ha) and available K (206.4 kg/ha).

The experiment was laid out in Randomized Block Design with nine treatments each replicated thrice. The crop was sown on 28th February 2022 using Shekhar-2 variety. The experiment was laid out in Randomized Block Design with nine treatments each replicated thrice viz **T₁** (25 ×10 cm + B 1.0 kg/ha), **T₂** (25 ×10 cm + B 1.5 kg/ha), **T₃** (25 ×10 cm + B 2.0 kg/ha), **T₄** (30 ×10 cm + B 1.0 kg/ha), **T₅** (30 ×10 cm + B 1.5 kg/ha), **T₆** (30 ×10 cm + B 2.0 kg/ha), **T₇** (35 ×10 cm + B 1.0 kg/ha), **T₈** (35 ×10 cm + B 1.5 kg/ha), **T₉** (35 ×10 cm + B 2.0 kg/ha).

Fertilizers were applied as band placement, for which 4-5cm deep furrows were made along the seed rows with a hand hoe. The nutrient sources were Urea, SSP and MOP to fulfil the requirement of nitrogen, phosphorous and potassium. The recommended dose of 20 kg/ha nitrogen, 40 kg/ha phosphorous and 20 kg/ha potassium and Boron 1.0, 1.5, 2.0 kg/ha were applied according to the treatments. The sources used for applying N P K and Boron were Urea (N 46%), Single Super Phosphate (P 16%) and Muricate of Potash (K 60%), Boron (Di Sodium Tetra Borate Penta Hydrate) (B 14.5%) respectively.

Seeds were sown in line manually on 28 Feb 2022 at a depth of 4-5 cm in furrows with seed rate of 15-20 kg/ha. Entire doses of NPK and Boron were applied as basal for respective plots. Seeds are then covered with soil immediately after sowing the seeds. The row-row spacing between crop 25, 30, 35 cm and plant to plant 10 cm were according to the treatment details. The growth parameters were recorded at periodical intervals of 15, 30, 45, 60 DAS and at harvest stage from the randomly selected five plants in each treatment. Statistically analysis was done for all the parameters in one-way Anova and mean compared at 5% probability level of significant results.

Results and Discussion

Effect of Spacing and Boron levels on growth parameters are given in table 1.

GROWTH ATTRIBUTES

Plant Height (cm)

At harvest, significantly highest plant height (44.53 cm) was recorded with the application of (2.0 kg/ha boron at 35 × 10 cm spacing) T₉ and T₈ (35 × 10 cm + B 1.5 kg/ha) (44.00 cm) was statistically at par with T₉. The spacing practices had significant effects on plant height (cm); however, an increasing trend with optimum geometry level could be noticed. This may be due to the competition between the inter and intra plant roots for sunlight, water, nutrients and space at closer spacing. Whereas, optimum spacing helped in significantly highest plant height. Significant results were obtained due to the optimum spacing of 35×10 cm and similar results were obtained by Singh *et al.* (2009). The increase in plant height may be due to an appropriate dose of boron. Because B plays important role in various enzymatic and other biochemical reactions. Similar results were discussed by Zahoor *et al.* (2011), Gitte *et al.* (2005).

No. of Branches/plant

At harvest, significantly highest No. of branches (9.27) was recorded with the application of (2.0 kg/ha boron at 35 × 10 cm spacing) T₉ and T₈ (35 × 10 cm + B 1.5 kg/ha) (9.20) was statistically at par with T₉. The optimum plant spacing between plants resulted in enhanced space, sunlight, nutrients and soil moisture for increased photosynthesis, metabolic activities, growth, and development which resulted in higher number of branches. The results were following Amruta *et al.*, (2015) and Jiotode *et al.*, (2017). The significantly maximum number of branches/plants was due to the application of boron. This might be due to the reason for the increase in this yield attribute that the boron plays important role in plant metabolism and translocation of photosynthates from source to sink. These are similar findings obtained by (Janaki *et al.*, 2018).

No. of nodules/plant

At harvest, significantly highest No. of nodules (6.33) was recorded with the application of (2.0 kg/ha boron at 35 × 10 cm spacing) T₉ and T₈ (35 × 10 cm + B 1.5 kg/ha) (6.13) was statistically at par with T₉. The optimum spacing resulted in the increase of nodulation, root development and growth, such increase in nodulation, root and growth might be due to an increase in the number of nodules which might have supplied sufficient nitrogen by nitrogen fixation and finally enhance the productivity of Black gram. Prasad *et al.*, (2014).

Dry Matter accumulation

At harvest, significantly highest dry weight (10.44) was recorded with the application of (2.0 kg/ha boron at 35 × 10 cm spacing) T₉ however T₈ (35 × 10 cm + B 1.5 kg/ha) (10.39 g) which was statistically at par to T₉. Higher dry matter production is observed in 40 × 10 cm² spacing due to better photosynthetic activity because of greater exposure to light and increased availability of nutrients to plants have also resulted in higher root dry weight on the plants results reported by Khan *et al.*, (2017).

Effect of Spacing and Boron levels on yield parameters of Black gram are given in table 2.

YIELD ATTRIBUTES AND YIELD

Pods/plant (No's)

At harvest, the treatment T₃ (25 × 10 cm + B 2.0 kg/ha) recorded a significantly highest number of pods/plant (30.73). However, T₂ (25 × 10 cm + B 1.5 kg/ha) (30.33) was found to be statistically at par with T₃. The higher number of pods/plants might be possible due to more vigour and strength attained by the plants because of better photosynthetic activities with sufficient availability of light, and supply of nutrients in the balanced quantity of the plants at growing stages. Jitendra kumar *et al.* (2015) observed similar results.

Seeds/pod (No's)

At harvest, the treatment T₃ (25 × 10 cm + B 2.0 kg/ha) recorded significantly highest number of Seeds/Pod (6.40). However, T₂ (25 × 10 cm + B 1.5 kg/ha) (6.13) was found to be statistically at par with T₃. The increase in the number of seeds per pod might be due to an increase in translocation of assimilates from source to sink. Shivay and Shekawat, (2008).

Test weight(g)

At harvest, significantly highest Test weight (42.48 g) was recorded with T₃ (25 × 10 cm + B 2.0 kg/ha) and T₂ (25 × 10 cm + B 1.5 kg/ha) (42.14 g) was statistically at par with T₃. Test weight of black gram seeds was increased due to the role of boron in increasing pollen viability and stigmatic receptivity which brings in increased seed set & increased translocation of photosynthesis to sink which increases test weight Prasad *et al.* (2015). Better availability of moisture and moderation of soil temperature led to greater uptake of nutrients and reduced number of days taken to meet the required heat units for proper growth and development of plants and ultimately the yield attributes. The results were recorded similar to Anand *et al.* (2020).

Seed Yield (Kg/ha)

At harvest, the treatment T₃ (25 × 10 cm + B 2.0 kg/ha) recorded significantly higher seed yield (930.33 kg/ha) over rest of the treatments. The B application improved the seed yield because it maintains a good balance between photosynthesis and respiration. Boron removal alters the cell wall structure, with a transitory decrease in elasticity modulus, followed by a secondary hardening and reduction in the incidence of plasma membrane bound reductase activity for better translocation to sink Shekawat and Shivay (2008). Similar results were obtained by Yu *et al.*, (2002).

Stover Yield (Kg/ha)

At harvest, the treatment T₃ (25 × 10 cm + B 2.0 kg/ha) stover yield (2045.33 kg/ha) was found significantly higher. However, T₂ (25 × 10 cm + B 1.5 kg/ha) (2040.33 kg/ha) was found to be statistically at par with T₃. The stover yield was significantly influenced by different levels of boron application M.A. Khan (2020). Higher stover yield of black gram at closed row spacing might be due to more population per unit area which contributed to more biomass and hence higher stover yield. Sathyamoorthi K. *et al.*, (2008). Similar results were obtained by Kailash Raman Bhatt (2020).

Harvest index (%)

Significantly highest harvest index (31.32) was observed in T₃ (25 × 10 cm + B 2.0 kg/ha) and is superior to rest of all treatments. "Highest harvest index was observed due to cell activities, enhanced cell multiplication and enlargement and luxuriant growth and yield attributes of the crops probably due to more absorption and utilization of available nutrients leading to overall improvement of crop growth reflected to source-sink relationship, which in turn enhanced the yield attributes that ultimately more yield" Kokani *et al.*, (2010).

Conclusion

Based on the findings of the experiment it can be concluded that maintaining a spacing of 25 × 10 cm and application of Boron @ 2.0 kg/ha in black gram during *zaid* season was found to be more productive as well as economically viable.

Acknowledgements

I express my sincere gratitude to my advisor **Dr. (Mrs.) Shikha Singh** for her constant support, guidance and valuable suggestions for improving the quality of this work. I am indebted to all the faculty members of Department of Agronomy, SHUATS, Prayagraj, Uttar Pradesh (U.P), India for their cooperation, encouragement and for providing indispensable facilities.

References

- Alloway BJ. 2008. Micronutrients and crop production: An introduction. In Micronutrient deficiencies in global crop production. *Springer, Dordrecht* 1-39.
- Ali A, Malik MA, Nadeem MA, Tahir M, Sohail R. 2002. Production potential of mashbean genotype in response to phosphorus application. *International Journal of Agriculture and Biology* 4:355-356

- Anand DD, Sumarjit Singh M, JamkhoginLhungdim N, Devi S, Gopimohan Singh.2020. N. Effect of Mulching on Growth and Yield of Soybean (*Glycine max* L.) Merrill). *International Journal of Current Microbiology and Applied Sciences***9**(9):3313-3318.
- Amruta, N., Maruthi, J.B., Sarika, G. and Deepika, C.2015. Effect of Integrated Nutrient Management and Spacing on Growth and Yield Parameters of Black Gram CV.LBG 625(Rashmi). *An International Quality Journal of Life Sciences***10**(1): 193-198.
- Aslam M, Hussain N, Zubair M, Hussain SB, Baloch MS. 2010. Integration of organic and inorganic sources of phosphorus for increased productivity of mung bean (*Vigna radiata* L). *Pakistan Journal of Agriculture science***47**(2):111 -114.
- Dwivedi, B.S., Ram, M., Singh, B.P., Das, M. and Prasad,R.N. 1990. Differential response of crops to boron fertilization in acid Alfisols. *Indian Journal of Agricultural Sciences***60**:122-127.
- Gitte AN, Patil SR, Tike TK .2005. Influence of zinc and boron on biochemical and yield characteristics of sunflower. *Ind. J. Plant Academia Journal of Agricultural Research Saeed* **10**: 400-403.
- Goldberg, S.; Glaubig, R.A. 1986. Boron adsorption on California soils. *Soil Science Society of America Journal* **50**:1173-1176.
- Gowda CL, Samineni S, Gaur PM, Saxena KB.2013. Enhancing the productivity and production of pulses in India. 145-159.
- Havlin, L.J., D.J. Beaton, L.S. Tisdale and L.W. Nelson. 1999. Soil fertility and fertilizers. *Prentice Hall of Indian*. 6th ed., Pp: 220, 227, 228, 319-346.
- Janaki, A.M., Parmar, K.B. and Vekaria, L.C. 2018. Effect of boron and molybdenum on yield and yield attributes of summer green gram (*Vigna radiata* L.) under medium black calcareous soils. *International Journal of Chemical Studies***6**(1): 321-323.

- Jiotode, D. J., Sonune, D. G., Mohod, A. R., Parlawar, N. D., Khawale, V. S. Studies on effect of weather parameters on *Kharif* green gram (*Vigna radiata* L.) varieties under different sowing date. *Association of Soils and Crops Research Scientists***27**(2): 185-191
- Jitendra Kumar, Sarvesh Kumar, Brajesh Prajapati, Amar Kant Verma and Awadhesh Kumar. 2016. Yield and yield attributes of plant geometry of Gram (*Cicer arietinum* L.) under rainfed conditions in Uttar Pradesh. *Research Environmental Life Science***9**(9):1087- 1089.
- Kailash Raman Bhatt.2020. "Effect of Plant Spacing on Yield and Yield Contributing Traits of Black gram (*Vigna mungo* L. Hepper) During Autumn Season at Gokuleshwor, Baitadi, Nepal". *Acta Scientific Agriculture***4**(3):01-04.
- Khan MA., Jahiruddin M., Bodruzzaman. M. 1996. Response of wheat to boron in relation to different varieties and sowing dates. *Bangladesh Journal of Agricultural Science***23**(2) 27-32.
- Khan K, Prakash V.2014. Effect of rhizobial inoculation on growth, yield, **nutrient**, and economics of summer urdbean (*Vigna mungo* L. Hepper) in relation to zinc and molybdenum. *International Journal of Advanced Research in Chemistry and Chemical Engineering***1**(1):1-10.
- Khan, M.M.S., Singh, V. P., & Kumar, A. 2017. Studies On Effect of Plant Densities on Growth and Yield of Kharif Mungbean (*Vigna radiata* L.) **Wilczek**. *Bulletin of Environment, Pharmacology and Life Sciences***6**(1): 291-295.
- Kokani JM, Shah KA, Tandel BM, Bhimani GJ. 2010.Effect of FYM, phosphorus, and sulphur on yield ofsummer black gramand post-harvest nutrient status of soil. *TheBioscan an International Quarterly Journal of Life Sciences***10**(1):379-383
- Loomis, W.D., Durst, R.W., 1992. Chemistry and biology of boron.*BioFactors***3**:229–239.
- Malik M F, Ashraf A, Qureshi S and Ghafoor A. 2007. Assessment of genetic variability, correlation and path analyses for yield and its components in soybean. *Pakistan J Bot***39**: 405-413.
- M.A. Khan, F.Naznin, M.A. Hossain, M.A.Islam and A.K.M.H. Rahman.2020.Effect of Boron on Growth, Yield and Nutrient Accumulation in Black gram.*A Scientific Journal of Krishi Foundation the Agriculturists* **18**(2): 34-43.
- Niaz, A., Ahmed, W., Zia, M.H. and Malhi, S.S. 2013.Relationship of soil extractable andfertilizer boron tosome soil properties, crop yields, and total boron incotton and wheat plants on selected soils of Punjab,*Pakistan. Journal of Plant Nutrition***36**:343–356.

- Prasad, S. K., Singh, M. K. and Singh, J. 2014. Response of rhizobium inoculation and phosphorus levels on mungbean (*Vigna radiata* L.) under guava-based agri-horti system. *The Bioscan* **9**: 557-560.
- Pushkar Choudhary, Gajendra Singh, GunapatiLakshma Reddy, Bhanwar Lal Jat.2017. Effect of Bio - fertilizer on Different Varieties of Black Gram (*Vigna mungo* L.) *Int. J. Curr. Microbiol. App. Sci***6**(2):302 -316.
- Sathyamoorthi K., *et al.* 2008.“Physiological paramters and yield of greengram (*vigna radiata* L.) Wilczek as influenced by increased plant density and fertilizer levels”. *Indian Journal of crop science***3**(1):115-122.
- Shamim, M.Z. and Pandey, A. 2017. Effects of arsenic toxicity on morphological characters in black gram (*Vigna mungo* L.) during early growth stage. *Cellular and Molecular Biology***63**(6): 38–43.
- Shekhawat K, Shivay YS. 2008. Effect of nitrogen source, sulphur and boron levels on productivity, nutrient uptake and quality of sunflower (*Helianthus annuus* L.). *Indian Journal of Agronomy***53**:129-134.
- Siddique M, Malik M F A and Awan S I. 2006. Genetic divergence, association and performance evaluation of different genotypes of mungbean (*Vigna radiata*). *Int J Agric Biol***8**: 793-795.
- Singh, H., Elamathi, S., and Anandhi, P.2009.Effect of row spacing and dates of sowing on growth and yield of lentil (*Lens culinaris*) under northern eastern region of U.P. *Legume Research***32**(4): 307-308.
- Singh AK, Khan MA, Arun S.2014. Effect of boron and molybdenum application on seed yield of mungbean. *Asian Journal of Bio Science* **9**(2):169-72.
- S Sharmila, P Saravana Pandian, R Indirani and E Subramanian.2020.Influence of boron on growth attributes, yield of black gram and available boron status in Typic chromustert.*International Journal of Chemical Studies***8**(6): 2291-2294.
- Tomar GS, Taunk SK, Chaudhary JL.2011. Black gram. In: Science of Crop production -part I (Kharif crops). Kushal Publications and Distributers, Maidagin, Varanasi -221 001 (U.P).
- Vahedi B, Gholipouri A and Sedghi M. 2010.Effect of planting pattern on radiation use efficiency, yield and yield components of sunflower.
- Veeramani P. 2019.Effect of plantspacing on the growth and yield of Black gram (*vigna mungo*).*Journal of Krishi Vignan***8** (1):101-104.

Verma CK, Yadav DD, Singh V.2011. Effect on yield and quality of green gram (*Vigna radiata* L.) varieties by foliar spray of urea and seed rates. *Plant Archives* **11**(1):289 -291.

Yu Q., Hlavacka A., Matoh T., Volkmann D., Menzel D., Goldbach H.E., and Baluska F., 2002. Short-term boron deprivation inhibits endocytosis of cell wall pectins in meristematic cells of maize and wheat root apices. *Plant Physiology***130**(1): 415–421.

Zahoor RSM, Basra A, Munir H, Ndeem MA, Yousaf S.2011. Role of boron in improving assimilates partitioning and yield in sunflower *J. Agric. Soc. Sci***7**:49-55.

Table 1 Effect of Spacing and Boron levels on growth attributes of Black gram.

S.No.	Treatments	At Harvest			At 60-75 DAS		
		Plant height(cm)	No. of Branches	Nodules/plant	Plant dry weight (g/plant)	Crop growth rate (g/m ² /day)	Relative growth rate (g/g/day)
1	25 ×10 cm + B 1.0 kg/ha	39.67	7.60	4.40	9.55	3.64	0.02
2	25 ×10 cm + B 1.5 kg/ha	39.90	7.80	4.67	9.65	2.26	0.02
3	25 ×10 cm + B 2.0 kg/ha	40.93	8.00	4.93	9.76	4.50	0.02
4	30 ×10 cm + B 1.0 kg/ha	41.57	8.20	5.20	9.84	3.27	0.02
5	30 ×10 cm + B 1.5 kg/ha	41.90	8.40	5.40	9.95	3.37	0.02
6	30 ×10 cm + B 2.0 kg/ha	42.70	8.60	5.67	10.16	4.51	0.02
7	35 ×10 cm + B 1.0 kg/ha	43.33	8.80	5.87	10.24	3.28	0.02
8	35 ×10 cm + B 1.5 kg/ha	44.00	9.20	6.13	10.39	3.45	0.02
9	35 ×10 cm + B 2.0 kg/ha	44.53	9.27	6.33	10.44	3.50	0.02
	F test	S	S	S	S	NS	NS
	S. EM (±)	0.18	0.07	0.07	0.02	0.59	0.01
	CD (P=0.05)	0.53	0.20	0.20	0.05	1.76	0.04

Table 2 Effect of Spacing and Boron levels on yield attributes of Black gram.

Treatments	At harvest					
	Pods/plant (No's)	Seed/pod (No')	Test weight (g)	Seed yield (kg/ha)	Stover yield (kg/ha)	Harvest index (%)
1. 25 ×10 cm + B 1.0 kg/ha	29.67	5.87	40.48	828.33	2032.67	28.95
2. 25 ×10 cm + B 1.5 kg/ha	30.33	6.13	42.14	862.67	2045.33	29.66
3. 25 ×10 cm + B 2.0 kg/ha	30.73	6.40	42.48	930.33	2040.33	31.32
4. 30 ×10 cm + B 1.0 kg/ha	28.80	5.20	37.63	739.67	1981.67	27.17
5. 30 ×10 cm + B 1.5 kg/ha	29.20	5.40	38.53	772.00	1989.00	27.95
6. 30 ×10 cm + B 2.0 kg/ha	29.40	5.60	39.36	793.67	2002.00	28.39
7. 35 ×10 cm + B 1.0 kg/ha	28.00	4.33	34.40	651.00	1957.00	24.96
8. 35 ×10 cm + B 1.5 kg/ha	28.40	4.60	35.78	685.00	1968.00	25.81
9. 35 ×10 cm + B 2.0 kg/ha	28.60	4.93	36.27	713.33	1977.00	26.51
F test	S	S	S	S	S	S
S. EM (±)	0.16	0.09	0.14	4.47	3.56	0.12
CD (P = 0.05)	0.47	0.27	0.41	13.41	10.66	0.35

