

## EFFECT OF ORGANIC, INORGANIC AND BIO-FERTILIZERS ON GROWTH, YIELD AND QUALITY OF TURNIP(*Brassica rapa* L.)

### ABSTRACT

The experiment was carried out with title “Effect of organic, inorganic and bio-fertilizers on growth, yield and quality of Turnip (*Brassica rapa* L.)cv. Purple Top White Globe” at the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Science, Prayagraj (U.P) During October 2021 to December 2021. The experiment was laid out in RBD with 9 treatments with 3 replications. The results revealed that Treatment T<sub>8</sub> (75% RDF through chemical fertilizers + 25% through vermicompost + PSB + Azotobacter) performed the best in terms of days to Plant Height (44.44 cm), No. of Leaves (22.26), Leaf length (30.08 cm), shoot weight (23.93 g), root length (8.19 cm), root diameter (6.62 cm), root yield per plant (117.25 g), root yield per hectare (234.51 q/ha), TSS °Brix (5.5), B:C Ratio (4.46:1). Therefore, the Treatment T<sub>8</sub> (75% RDF through chemical fertilizers + 25% through vermicompost + PSB + Azotobacter) is the best when compared to other treatments. As the highest benefit cost ratio was observed in treatment T<sub>8</sub> (75% RDF through chemical fertilizers + 25% through vermicompost + PSB + Azotobacter) i.e., (4.46:1) which states that it is economically profitable compared to all other treatments.

**Key words:-** Turnip, PSB, Azotobacter, growth, Yield, Quality

### 1. INTRODUCTION

Turnip (*Brassica rapa* L.) is a member of the cruciferous family of vegetables. *Brassica rapa* L. commonly known as field mustard or turnip mustard is a plant widely cultivated as a leaf vegetable, a root vegetable, and an oilseed. (Duke and Ayensu 1985). It has a crisp white flesh and a zesty mustard-like flavor and a rapidly maturing crop (Choudhary 2015). Its root is called underground modified root, which is napiform in shape. The stem is short at vegetative stage but elongated at the reproductive stage. A decoction of the leaves or stems is used in the treatment of cancer (Duke and Ayensu 1985). The powdered seed is said to be a folk remedy for cancer (Duke 1983). The crushed ripe seeds are used as a poultice on burns (Foster and Duke 1990). The root when boiled with lard is used for breast tumors (Duke 1983). A salve derived from the flowers is said to help skin cancer (Duke 1983). Moreover, turnip extract is also useful for lowering uric acid and extracting renal stones. It increases

visual keenness and is used to treat night blindness. Turnip syrup strengthens the memory (**Khashayar 2007**). Turnip root peelings contain a natural insecticide (**Allardice 1993**). In India Turnip cultivation is mostly confined to north-western parts of the country like Punjab, Haryana, Rajasthan and Western U. P. as an early winter season crop. Green Top, Purple Top and 'Kenshin-Kaba' are important varieties of Turnip for fodder production. Although many agronomic factors affect its production but the optimum date of sowing and nitrogen dose are the most important factors affecting the production and productivity of fodder Turnip of its short duration and its effect on the crop growth and green fodder and root yield. (**Turk et al., 2009, Sarhan et al., 2003, Smart et al., 2004**). There is a great scope to increase the production and productivity of fodder Turnip by selecting the optimum date of sowing and nitrogen requirement for getting higher yield, higher benefit cost ratio and better quality of fodder Turnip.

Organic manures comprise of a variety of plant-derived materials that range from fresh or dried plant material to animal manures and litters to agricultural by-products. The nutrient content of organic fertilizers varies greatly among source materials, and readily biodegradable materials make better nutrient sources. Nitrogen and Phosphorus content is lower, often substantially lower, in organic manures compared to chemical fertilizers. Moisture content is another factor that reduces or dilutes the nitrogen and phosphorus concentrations of organic fertilizers. Thus, it can be cost ineffective to transport high moisture organic manures long distances. The bio-fertilizers are organic in origin and thus are absolutely safe. Therefore, it is essential to adopt a strategy of integrated nutrient management using combination of chemical fertilizers, organic manures and bio-fertilizers so as to minimize the cost of production and to maintain biological productivity of soils. The interactive advantages of combining inorganic and organic sources of nutrient generally provide superior status to use of each component separately. The judicious application of organic and inorganic fertilizers have maintained long term soil fertility and sustained higher level of productivity.

This study aims to determine the effect of organic, inorganic and bio-fertilizers on growth, yield and quality of Turnip.

## 2. MATERIALS AND METHODS

The field experiment entitled "Effect of organic, inorganic and bio-fertilizers on growth, yield and quality of Turnip (*Brassica rapa* L.)" was adapted randomized block design consisting of 8 treatments and three replications during October 2021 to December 2021. Horticulture Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.). The experiment includes the following treatments: T<sub>0</sub> RDF (Control 60:40:40 NPK + FYM 25 t/ha), T<sub>1</sub> 75% RDF Through chemical fertilizer + 25% through FYM, T<sub>2</sub> 75% RDF Through chemical fertilizer + 25% through vermicompost, T<sub>3</sub> 75% RDF Through chemical fertilizers + 25% through FYM + PSB, T<sub>4</sub> 75% RDF Through chemical fertilizer + 25% through Vermicompost + PSB, T<sub>5</sub> 75% RDF Through chemical fertilizer + 25% through FYM +

Azotobacter, T<sub>6</sub> 75% RDF Through chemical fertilizer + 25% through vermicompost + Azotobacter, T<sub>7</sub> 75% RDF Through chemical fertilizer + 25% through FYM + PSB + Azotobacter, T<sub>8</sub> 75% RDF Through chemical fertilizer + 25% through Vermicompost + PSB + Azotobacter. The mean (maximum and minimum) temperature was 35.77°C and 8.28°C respectively, mean (maximum and minimum) relative humidity was 93.5% and 43.71% during the crop growing season. The experimental soil was sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.318%), medium in available N (87 Kg/ha), medium available P (375 Kg/ha) and medium available K (50 Kg/ha). Fertilizers were applied in the form of urea, single super phosphate and murate of potassium, respectively.

### 3. RESULTS AND DISCUSSION

#### 3.1 Growth parameters

The statistical data on growth parameters are recorded in table 1. The maximum Plant Height (44.44 cm), plant height (22.26), Leaf length (30.08 cm) of Turnip was found in T<sub>8</sub> (75% RDF through chemical fertilizers + 25% through vermicompost + PSB + Azotobacter). The mean value of plant height was found significant at different levels of NPK, Vermicompost, PSB and Azotobacter. Increase in plant height is due to adequate supply of nutrients which turns helps in vigorous vegetative growth of plants and subsequently increase the plant through cell elongation, cell division, photosynthesis and turbidity of plant cells. The increase in nodulation and nitrogen fixation leads to more plant height. Similar findings were reported by **Chavan *et al.*, (2015)**, **Deshmukh *et al.*, (2014)**, **Ayubet *et al.*, (2012)**, **Sajid *et al.*, (2009)**. The bio-fertilizers lead to good root development, better water absorption and high uptake of nutrients from the soil body, which ultimately enhance number of leaves, leaf length per plant similar observation have been reported by **Chattoo *et al.*, (2005)**.

#### 3.2 Yield and quality parameters

The statistical data on growth parameters are recorded in table 2. The maximum Shoot Weight of Turnip (23.93 g), Root Length (8.19 cm), Root Diameter (6.62 cm), root yield plant (117.25 g), root yield per hectare (234.15 q/ha) was found in T<sub>8</sub> (75% RDF through chemical fertilizers + 25% through vermicompost + PSB + Azotobacter). Organic, inorganic and bio-fertilizers played a pivotal role in improving shoot weight of Turnip. Similar findings have been reported by **Pimpini *et al.*, (1992)**. Increase in root length, The higher root weight and root yield may be due to more organic matter added in the soil would be helped better penetration roots in the soil and thereby better root length, root weight and root yield Similar reports have been given **Abbey *et al.*, (2003)**. The root diameter of Turnip is increase with the increase the photosynthetic activity and higher nutrients uptake that results the increasing the root diameter similar finding have been reported by **Manivannan *et al.*, (2004)**. Significantly the maximum TSS (°Brix) of Turnip was found in T<sub>8</sub> (75% RDF through chemical fertilizers + 25% through vermicompost + PSB + Azotobacter) which was 5.5 in 50 DAS. **Manoj *et al.*, (2011)** observed that treatment with high as well as low content of total nutrient resulted in significantly lower TSS and soluble carbohydrates. Whereas higher sugar content was obtained with application of higher dose of nutrient and vermicompost. The

maximum BC ratio (4.46) was also recorded maximum under the treatment T<sub>8</sub> which is shown in table 2.

#### **4. CONCLUSION**

The present investigation entitled “Effect of organic, inorganic and bio-fertilizers on growth, yield and quality of Turnip (*Brassica rapa* L.)” it was concluded that the treatment T<sub>8</sub> (75% RDF through chemical fertilizers + 25% through vermicompost + PSB + Azotobactor) was found best in plant height, number of leaves, leaf length, shoot weight, root weight, root length, root diameter, root yield, TSS, net return of 364052 Rs/ha with cost benefit ratio of 4.46 .

**Table 1. Effect of organic, inorganic and bio-fertilizers on growth parameters of Turnip.**

Treatments	Treatment combination	Plant height			Number of leaves			Leaf length
		15 DAS	30 DAS	At Harvest	15 DAS	30 DAS	50 DAS	
T <sub>0</sub>	RDF (C ontrol 60:40:40 NPK + FYM 25 t/ha)	7.93	18.03	36.63	3.47	6.78	13.89	20.94
T <sub>1</sub>	75% RDF + 25% FYM	8.85	18.89	37.72	4.17	8.12	17.70	23.50
T <sub>2</sub>	75% RDF + 25% vermicompost	9.42	19.49	38.14	4.49	8.26	18.57	24.49
T <sub>3</sub>	75% RDF + 25% FYM + PSB	8.37	18.54	37.13	3.88	7.35	16.60	21.49
T <sub>4</sub>	75% RDF + 25% Vermicompost + PSB	9.93	19.32	39.32	4.84	8.78	19.55	25.05
T <sub>5</sub>	75% RDF + 25% FYM + Azotobacter	11.97	21.44	41.15	5.59	9.68	20.75	27.40
T <sub>6</sub>	75%RDF + 25% vermicompost + Azotobacter	10.96	20.83	40.34	5.09	9.27	20.01	26.42
T <sub>7</sub>	75% RDF + 25% FYM + PSB + Azotobacter	12.70	23.21	41.33	6.02	10.56	21.69	28.81
T <sub>8</sub>	75% RDF + 25% Vermicompost + PSB + Azotobacter	13.46	24.23	44.44	6.52	11.51	22.26	30.08
	<b>F-Test</b>	S	S	S	S	S	S	S
	<b>SEd(±)</b>	1.19	0.17	0.23	6.77	5.64	4.34	3.31
	<b>C.D. 0.05%</b>	0.57	0.52	0.68	0.19	0.29	0.48	0.48
	<b>CV</b>	3.18	1.48	0.99	0.57	0.87	1.43	1.45

**Table 2. Effect of organic ,inorganic and bio-fertilizers on yield and quality parameters on Turnip**

Treatments	Treatment combination	Shoot weight(g)	Root length(cm)	Root diameter(cm)	Root weight(g)	Root weight(q/ha)	TSS (°Brix)	B:C Ratio
T <sub>0</sub>	RDF (control 60:40:40 NPK+FYM 25 t/ha)	18.96	5.66	4.74	84.42	168.85	4.8	1.44
T <sub>1</sub>	75% RDF + 25% FYM	20.94	6.17	5.31	105.28	210.56	5.1	3.43
T <sub>2</sub>	75% RDF + 25% vermicompost	21.23	6.58	5.52	111.83	223.65	5.1	4.36
T <sub>3</sub>	75% RDF + 25% FYM+PSB	20.00	6.10	5.06	96.65	193.31	4.9	3.14
T <sub>4</sub>	75% RDF + 25% Vermicompost + PSB	22.30	6.94	5.63	113.16	226.31	5.2	4.34
T <sub>5</sub>	75% RDF + 25% FYM + Azotobacter	23.36	7.52	6.11	115.31	230.62	5.3	3.73
T <sub>6</sub>	75% RDF + 25% vermicompost + Azotobacter	22.84	7.10	6.04	114.76	229.51	5.3	4.39
T <sub>7</sub>	75% RDF + 25% FYM + PSB + Azotobacter	23.54	7.79	6.30	116.10	232.21	5.4	3.74
T <sub>8</sub>	75% RDF + 25% Vermicompost + PSB + Azotobacter	23.93	8.19	6.62	117.25	234.51	5.5	4.46
	<b>F-Test</b>	S	S	S	S	S	S	
	<b>SE (d)±</b>	3.87	6.44	5.91	1.77	1.77	1.81	
	<b>C.D. 05</b>	0.49	0.26	0.19	1.11	2.22	0.05	
	<b>CV</b>	1.47	0.77	0.58	3.32	6.65	0.16	

**REFERENCES**

- Abbey, L.,(2000)** Effect of poultry manure and post-production application of fungicide on the shelf life of Onion cv Bawku Red *Crop Research*,Hissar **20**(1): 87-92.
- Allardice, P., (1993)** A - Z of Companion Planting. *Cassell Publishers Ltd.* 1993 ISBN 0-304-34324-2 A well produced and very readable book.
- Chattoo, M.A., Gandroo, MY., Zargeer, MY.,(1997)** Effect of Azospirillum and Azotobacter on growth, yield and quality of knol-khol (*Brassica oleracea* var. *gongylodes*). *Vegetable Science* **24**(1):16-19.
- Chavan,P.J., Syed,I., Rudraksh, G.B., Malewar,G.V.,and Baig,M.I.,(1997)** Effect of various nitrogen levels through FYM and urea on yield, uptake of nutrients and ascorbic acid content in chilli *Journal Indian Soil Science* **4**(4): 833-835.
- Duke, J.A. and Ayensu,E.S., (1985)** Medicinal Plants of China *Reference Publications, Inc.* ISBN 0-917256-20-4.
- Duke,J., (1983)** Handbook of energycrops published only on the internet excellent information on a wide range of plants.
- Manivannan, M. I., Singh, J.P., (2004)** Effect of bio-fertilizers on the growth and yield of sprouting broccoli (*Brassica oleracea* var. *Italica Plenck*) under Allahabad agro-climatic conditions. *Bioved* **15**(1/2): 33-36.
- Manoj, Das, Prasad, K.K., Prasad and Kumar (2011)** Effect of integrated nutrient management on quality of broccoli (*Brassica oleraceae*)cv.fiesta under jharkand conditions. *The Asian journal of horticulture* **6** (2): 388-392.
- Pimpini, F., Giardini ,L., Borin ,M., Gianquinto, G., (1992)** Effect of poultry manure and mineral fertilizers on the quality of crops *Journal of agriculture sciences* **118** (2): 215- 221.