

## **Original Research Article**

# **Carriage of HBs antigen and hepatitis C virus in people infected with human immunodeficiency virus at the Grand-Yoff General Hospital in Dakar, Senegal**

### **Abstract**

**Objective :** To investigate the carriage of HBsAg and hepatitis C virus in people with human immunodeficiency virus (HIV) at Grand-Yoff General Hospital.-

**Materials and methods :** Sera from patients living with HIV and monitored at Grand-Yoff General Hospital have benefited from research for HBs Ag, HBe Ag, anti-HBe antibodies and on these co-infected patients, a test for hepatitis C virus was also carried out. **Results :** This study included 147 HIV-infected patients and including 95 women (64.63%) and 52 men (35, 37%) with a median age of 46 years (16-78 years). The sex ratio is M/F was 0.54 and the age group most represented at inclusion was 30-39 years old with 31.97% of patients, 12, 25% of patients were co-infected HIV and hepatitis B including 11 women (61.11%) and 7 men (38.89%). The median age was 39.5 years (33-60 years), women were the affected by co-infection and 44, 44% of them were married. Chronic hepatitis B and hepatitis C markers were tested for in HIV / HBV coinfecting patients using a microparticulate chemiluminescence immunoassay technic. Investigation of hepatitis B chronicity markers indicated that all patient, except one, underwent anti-HBe seroconversion (94,4%) and no case of co-infection with HCV had been noted. During follow-up, all patients were under antiretroviral therapy with mainly tenofovir (TDF) which was the most used molecule with 83.33%, followed by Lamivudine (3TC) 16.67%.-

**Conclusion :** HIV infection is a major public health problem when it is associated with hepatitis B and C virus.-

**Keywords :** Co-infection, Hepatitis, HIV, HCV

### **1. Introduction**

Human immunodeficiency virus (HIV) infection remains a major public health problem with 36.9 million people infected worldwide, 25.7 million of them live in the African region [12]. Sharing transmission routes with hepatitis B and C viruses, co-infections with these viruses have become real factors of comorbidity and mortality [4]. HBV and HCV are the two most dangerous liver viruses as they become chronic lifelong diseases in most people. Hepatitis B is the leading cause of acute and

Style Definition ... [1]

Formatted: Font: 10 pt, Not Bold, Font color: Auto

Formatted: Body Text, Left, Line spacing: single

Formatted: Title, Left, None

Formatted ... [2]

Formatted: Font: Arial, 10 pt, Italic, Font color: Auto

Formatted: Body Text, Left, Line spacing: single

Formatted: Indent: Left: 0.27", Right: 0.74", Space Before: 4.25 pt

Formatted ... [3]

Formatted: Font: 17.5 pt

Formatted: Body Text, Left, Space Before: 0.55 pt, Line spacing: single

Formatted: Heading 1, Left, Line spacing: single

Formatted: Body Text, Left, Indent: Left: 0.15", Space Before: 6.85 pt

Formatted ... [4]

Formatted ... [5]

Formatted: Body Text, Left, Indent: Left: 0.15", Right: 0.55"

Formatted ... [6]

Formatted ... [7]

chronic liver disease in the world with 257 million people living with the [12]. The prevalence of hepatitis B surface antigen (HBsAg) carriage varies widely by geographical area, with sub-Saharan Africa and the WHO Western Pacific regions being areas of high prevalence, with 6.1% and 6.2% of the adult population infected respectively [12]. For hepatitis C virus, approximately 2.3 million people (6.2%) are co-infected with HCV worldwide with only 2.4% among people living with HIV. In Senegal, 85% of the general population have at least one HBV marker [15] [11] and the prevalence of HBsAg assessed in several population groups of interest ranged from 7.35% in blood donors to 14% in prisoners [8].

HIV/HBV co-infection as well as HIV/HCV co-infection are very frequent in the world and Africa remains the most affected continent. In Senegal, there is little data on the prevalence of these co-infections. Hospital studies have shown HBV prevalence among PLHIV to be 16.8% and 16.98% respectively [5] and 16.98% [1]. However, no data on HIV/HCV co-infection is available.

Human immunodeficiency virus (HIV) infection remains a major public health problem with 36.9 million people infected worldwide, 25.7 million of them live in the African region [12]. Sharing transmission routes with hepatitis B and C viruses, co-infections with

these viruses have become real factors of comorbidity and mortality [4]. HBV and HCV are the two most dangerous liver viruses as they become chronic lifelong diseases in most people. Hepatitis B is the leading cause of acute and chronic liver disease in the world with 257 million people living with the [12]. The prevalence of hepatitis B surface antigen (HBsAg) carriage varies widely by geographical area, with sub-Saharan Africa and the WHO Western Pacific regions being areas of high prevalence, with 6.1% and 6.2% of the adult population infected respectively [12]. For hepatitis C virus, approximately 2.3 million people (6.2%) are co-infected with HCV worldwide with only 2.4% among people living with HIV.

In Senegal, 85% of the general population have at least one HBV marker [15] [11] and the prevalence of HBsAg assessed in several population groups of interest ranged from 7.35% in blood donors to 14% in prisoners [8].

HIV/HBV co-infection as well as HIV/HCV co-infection are very frequent in the world and Africa remains the most affected continent. In Senegal, there is little data on the prevalence of these co-infections. Hospital studies have shown HBV prevalence among PLHIV to be 16.8% and 16.98% respectively [5] and 16.98% [1]. However, no data on HIV/HCV co-infection is available.

It is in this perspective that we were interested in the HIV/HBV/HCV morbid association, by studying the prevalence of HBsAg and hepatitis C virus carriage among PLWHA in the active file of the Grand-Yoff General Hospital.

## 2. Matériel and méthodes

### 2.1. Type and période of study

This was a retrospective study of 147 HIV-infected patients followed at Grand-Yoff General Hospital during the period July to December 2017.

### 2.2. Study population

Our study involved all HIV-infected patients in the active file who came for consultation during the study period at the Grand Yoff General Hospital and who met the inclusion criteria.

### 2.3. Inclusion and exclusion criteria

All patients living with HIV were included in the study, excluding those lost to follow-up or with incomplete medical records.

### 2.4. Méthodology

Formatted: Font: 12 pt

Formatted: Body Text, Indent: Left: 0.15", Right: 0.6"

Formatted ... [8]

Formatted: Font: 12 pt, Font color: Auto, Pattern: Clear

Formatted: Heading 1, Justified, Indent: Hanging: 0.25", Space Before: 0.1 pt, After: 0 pt, Line spacing: single, Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.15" + Indent at: 0.4", Tab stops: 0.4", Left

Formatted ... [9]

Formatted ... [10]

Formatted: Body Text, Indent: Left: 0.15", Right: 0.6", Space Before: 7 pt

Formatted: Body Text, Left, Space Before: 0.5 pt, Line spacing: single

Formatted: Heading 1, Justified, Indent: Hanging: 0.53", Line spacing: single, Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.15" + Indent at: 0.68", Tab stops: 0.68", Left

Formatted ... [11]

Formatted: Body Text, Indent: Left: 0.15", Right: 0.6", Space Before: 6.95 pt

Formatted ... [12]

Formatted: Heading 1, Justified, Indent: Hanging: 0.53", Line spacing: Exactly 13.75 pt, Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.15" + Indent at: 0.68", Tab stops: 0.68", Left

Formatted ... [13]

Formatted: Body Text, Indent: Left: 0.15", Right: 0.6", Space Before: 6.85 pt

Formatted ... [14]

Formatted: Heading 1, Justified, Indent: Hanging: 0.53", Line spacing: single, Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.15" + Indent at: 0.68", Tab stops: 0.68", Left

A blood sample on EDTA tube was taken from each patient and sent to the laboratory within two hours. After centrifugation, HIV 1/2 antibodies were tested using a chromatographic screening test (Determine® HIV 1/2 rapid test) and a confirmatory discriminatory test (*ImmunoComb® II HIV 1 & 2 Bispot*). HBsAg testing was performed using a rapid immunochromatographic test (ABON® HBs-Ag; Alere, USA). For the determination of hepatitis B markers and anti-HCV antibodies, the Architect automated immunoassay platform was used to perform immunoassays using microparticle chemiluminescence assay technology.

## 2.5. -Data analysis-

The data were collected on consultation forms. Data entry was performed using Microsoft Office Access (version 2007) database management software. Sociodemographic, clinical and —biological data were collected. Statistical analysis was performed using Excel 2016.

## 3. Results

### 3.1. Socio-demographic characteristics of the active file

The characteristics of our study population are resumed in **Table I**. A total of 147 patients were included with a sex ratio M/F of 0.54 and a median age of 46 years. The majority of the HIV-infected patients were married (n=73; 49.66%). Regarding the age groups most affected by HIV infection, those aged 30-39 and 40-49 were more representative with prevalences of 31.97% (n=47) and 29.93% (n=44) respectively (**Table II**).

**Table I: Socio-demographic characteristics of patients living with HIV in the active file**

	Number	Percentage
<b>Number of patients</b>	147	100
<b>Gender</b>		
Male	52	35,37

Formatted [15]

Formatted: Font: 12 pt

Formatted [16]

Formatted: Body Text, Indent: Left: 0.15", Right: 0.6"

Formatted: Heading 1, Justified, Indent: Hanging: 0.53", Line spacing: single, Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.15" + Indent at: 0.68", Tab stops: 0.68", Left

Formatted: Body Text, Indent: Left: 0.15", Right: 0.59", Space Before: 6.85 pt

Formatted [17]

Formatted: Heading 1, Justified, Indent: Hanging: 0.25", Line spacing: single, Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.15" + Indent at: 0.4", Tab stops: 0.4", Left

Formatted: Font: 12 pt, Font color: Auto, Pattern: Clear

Formatted [18]

Formatted: Body Text, Indent: Left: 0.15", Right: 0.59", Space Before: 6.85 pt

Formatted [19]

Formatted: Font: 13 pt

Formatted: Body Text, Left, Line spacing: single

Formatted: Font: 12 pt, Character scale: 105%, Condensed by 0.05 pt

Formatted: Heading 1, Left, Indent: Left: 0.23", Hanging: 2.95", Right: 0.55", Space Before: 10.4 pt

Formatted [20]

Female	95	64,63
<b>Median age at diagnosis [Extreme]</b>	41 ans	*
	[6 – 75 years]	
<b>Current median age [Extreme]</b>	46 ans	*

Formatted: Font: 12 pt, Bold

**Table II : Main socio-demographic characteristics of co-infected active file**

	<u>Number</u>	<u>Percentage</u>
<b>HIV/HBV co-infected active file</b>	18 (12,25%)	100

Number of patients	Number	Percentage
	147	100
<b>Gender</b>		
Male	52	35,37
Female	95	64,63
Female	11	61,11
<b>Median age at diagnosis [VHB [Extreme]]</b>		
	41 ans	*
	[6 – 75] years	[30 – 55 years]
<b>Current median age [Extreme]</b>		
	46 ans	*
	[16 – 78] years	[33 – 60 years]
<b>Marital status</b>		
Singles	5	27,78
<b>Residence</b>		
Senegal	130	88,44
Foreigner	12	8,16
Not specified	5	3,40
Married	2	11,11
Divorced	2	11,11
Widow (er)	1	5,56
<b>Marital status</b>		
Single	12	8,17
Divorced	9	6,12
Married	73	49,66
Widow (er)	14	9,52
Not specified	39	26,53
<b>Age groups (years) at inclusion</b>		
<20	2	1,36
20-29	12	8,16
30-39	47	31,97
40-49	44	29,93
50-59	25	17,01
60-69	13	8,85
≥70	4	2,72
Total	147	100
30-50	16	88,9
>50	1	5,6

- Formatted ... [21]
- Formatted ... [22]
- Formatted ... [30]
- Formatted Table ... [23]
- Formatted ... [26]
- Formatted ... [27]
- Formatted ... [31]
- Formatted ... [25]
- Formatted ... [24]
- Formatted ... [28]
- Formatted ... [29]
- Formatted ... [32]
- Formatted ... [33]
- Formatted ... [34]
- Formatted Table ... [36]
- Formatted ... [37]
- Formatted ... [40]
- Formatted ... [41]
- Formatted ... [35]
- Formatted ... [38]
- Formatted ... [39]
- Formatted ... [42]
- Formatted ... [43]
- Formatted ... [44]
- Formatted ... [47]
- Formatted ... [48]
- Formatted ... [45]
- Formatted ... [46]
- Formatted Table ... [49]
- Formatted ... [54]
- Formatted ... [50]
- Formatted ... [51]
- Formatted ... [52]
- Formatted ... [53]
- Formatted ... [55]
- Formatted ... [56]
- Formatted Table ... [57]
- Formatted ... [58]
- Formatted ... [59]
- Formatted ... [60]
- Formatted ... [61]
- Formatted ... [62]
- Formatted ... [63]
- Formatted ... [64]
- Formatted ... [65]
- Formatted ... [66]
- Formatted ... [67]
- Formatted ... [68]
- Formatted ... [71]
- Formatted ... [72]
- Formatted ... [73]
- Formatted ... [74]
- Formatted ... [69]
- Formatted ... [70]

Total	18	100
HIV/HCV co-infection	0	0

**Table II : Main socio-demographic characteristics of co-infected active file**

HIV/HBV co-infected active file	Number	Percentage
	18 (12,25%)	100
<b>Gender-</b>		
Male	7	38,89
Female	11	61,11
<b>Median age at diagnosis VHB [Extreme]</b>	39 years [30—55 years]	
<b>Current median age [Extreme]</b>	39,5 years [33—60 years]	±
<b>Marital status</b>		
Singles	5	27,78
Married	8	44,44
Divorced	2	11,11
Widow (er)	1	5,56
Not specified	2	11,11
<b>Age groups at inclusion</b>		
<30	1	5,6
30-50	16	88,9
≥50	1	5,6
Total	18	100
HIV/HCV co-infection	0	0

Formatted: Font: 10 pt

Formatted: Body Text, Left, Line spacing: single

### 3.2. Characteristics of HIV/HBV and HIV/HCV co-infected patients

Hepatitis B surface antigen (HBsAg) was found in 12.25% (n=18) of the patients living with HIV who were predominantly female (n=11; 61.11%) with a sex ratio of 0.63 M/F. Their median age was 39.5 years (33-60 years). None of patients living with HIV was infected with HCV.

### 3.3. Treatment regimen

All patients were on ARV treatment. The most commonly used treatment was the combination Tenofovir+Lamivudine+Efavirenz (77.77%; n=14), followed by the combination Zidovudine+Lamivudine+Efavirenz or Nevirapine with 11.11% (n=3). Only

one patient (5.56%) is on second line treatment with Tenofovir+Lamivudine+Lopinavir/Ritonavir (Figure 1). For treatment against hepatitis B virus, Tenofovir (TDF) remains the most used molecule with 83.33%, followed by Lamivudine (AZT) 16.67%. Concerning the occurrence of opportunistic infections, 16.67% presented opportunistic infections such as gastroenteritis (n=1), tuberculosis (n=2).

- Formatted: Font: 12 pt
- Formatted: Font: 12 pt, Expanded by 0.05 pt
- Formatted: Font: 12 pt, Expanded by 0.05 pt
- Formatted: Font: 12 pt, Expanded by 0.05 pt
- Formatted: Font: 12 pt, Expanded by 0.05 pt
- Formatted: Body Text, Indent: Left: 0.15", Right: 0.59"
- Formatted: Font: 12 pt, Expanded by 0.05 pt
- Formatted: Font: 12 pt, Expanded by 0.05 pt
- Formatted: Font: 12 pt
- Formatted: Font: 12 pt, Expanded by 2 pt
- Formatted: Font: 12 pt, Expanded by 1.75 pt
- Formatted: Font: 12 pt, Expanded by 1.9 pt
- Formatted: Font: 12 pt, Expanded by 2.05 pt
- Formatted: Font: 12 pt, Expanded by 2.1 pt
- Formatted: Font: 12 pt, Expanded by 2 pt
- Formatted: Font: 12 pt, Condensed by 3 pt
- Formatted: Font: 12 pt, Not Expanded by / Condensed by
- Formatted: Font: 12 pt, Expanded by 0.05 pt
- Formatted: Font: 12 pt, Character scale: 105%
- Formatted: Font: 12 pt, Character scale: 105%, Condensed by 3 pt
- Formatted: Font: 12 pt, Condensed by 0.05 pt
- Formatted: Font: 12 pt, Condensed by 0.1 pt
- Formatted: Font: 12 pt, Expanded by 0.1 pt
- Formatted: Font: 12 pt, Expanded by 0.05 pt
- Formatted: Font: 12 pt, Condensed by 0.15 pt
- Formatted: Font: 12 pt, Expanded by 0.2 pt
- Formatted: Font: 12 pt, Condensed by 0.05 pt
- Formatted: Font: 12 pt, Character scale: 100%

### Figure 1 : Antiretroviral treatment regimen for co-infected HIV/HBV patients

#### 4. Discussion

This study made it possible to estimate the prevalence of HIV/HBV, HIV/HCV and HIV/HBV/HCV co-infection in the active file, at the Grand-Yoff General Hospital due to the high prevalence of HIV and viral hepatitis in sub-Saharan Africa, which constitute major public health problems.

Our study still showed the feminization of HIV infection (64.63%) as demonstrated in other studies notably in Nigeria (65.2%) [9], Ethiopia (69.5%) [16] and Ghana (71%) [14].

This female predominance was also found among HIV/HBV co-infected patients (61%). Similar results were found in Burkina Faso, Nigeria and Ethiopia. In contrast, a Nigerian study reported a predominance of HIV-HBV co-infection in men [3]. The median age at diagnosis of the active file in our study population was 41 years with extremes of 6 and 75 years and that of the HIV/HBV co-infected active file was 39 years with extremes of 33 and 60 years. Wondimeneh in Ethiopia [16] and Jacquet in Senegal [8] respectively found median ages of 32 and 30 years, lower than ours for patients living with HIV. Concerning co-infected patients, a study carried out in Burkina Faso showed a median age of 38 years [2].

HBsAg carriage among PLHIV was 12.25% (n=18). Similar studies have been observed previously in Dakar [5] but also in other countries such as Benin [6] and Ghana [7] and results were respectively 16.8%, 16.9% and 16.7% higher than ours. On the other hand, a Botswana study conducted in 2011 reported a prevalence of co-infection of 5.3% [13].

Our study still showed the feminization of HIV infection (64.63%) as demonstrated in other studies notably in Nigeria (65.2%) [9], Ethiopia (69.5%) [16] and Ghana (71%) [14].

This female predominance was also found among HIV/HBV co-infected patients (61%). Similar results were found in Burkina Faso, Nigeria and Ethiopia. In contrast, a Nigerian study reported a predominance of HIV-HBV co-infection in men [3]. The median age at diagnosis of the active file in our study population was 41 years with extremes of 6 and 75 years and that of the HIV/HBV co-infected active file was 39 years with extremes of 33 and 60 years. Wondimeneh in Ethiopia [16] and Jacquet in Senegal [8] respectively found

Formatted: Condensed by 0.1 pt
Formatted: Heading 1, Left, Indent: Left: 0.53"
Formatted: Condensed by 0.15 pt
Formatted: Condensed by 0.1 pt
Formatted: Condensed by 0.1 pt
Formatted: Condensed by 0.15 pt
Formatted: Condensed by 0.1 pt
Formatted: Condensed by 0.15 pt
Formatted: Condensed by 0.1 pt
Formatted: Condensed by 0.15 pt
Formatted: Font: 12 pt
Formatted: Justified, Indent: Hanging: 0.25", Space Before: 10.05 pt, Line spacing: single, Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.15" + Indent at: 0.4", Tab stops: 0.4", Left
Formatted: Font: 12 pt, Character scale: 100%
Formatted: Body Text, Indent: Left: 0.15", Right: 0.6", Space Before: 6.85 pt
Formatted: Font: 12 pt, Character scale: 100%, Expanded by 0.05 pt
Formatted: Font: 12 pt, Character scale: 100%
Formatted: Font: 12 pt, Character scale: 100%, Expanded by 0.05 pt
Formatted ... [75]
Formatted ... [76]
Formatted ... [77]
Formatted ... [78]
Formatted ... [79]
Formatted ... [80]
Formatted ... [81]
Formatted ... [82]
Formatted ... [83]
Formatted ... [84]
Formatted ... [85]
Formatted ... [86]
Formatted ... [87]
Formatted ... [88]
Formatted ... [89]
Formatted ... [90]
Formatted ... [91]
Formatted ... [92]
Formatted: Font: 12 pt, Expanded by 0.05 pt
Formatted: Font: 12 pt, Condensed by 0.1 pt
Formatted ... [93]
Formatted ... [94]
Formatted ... [95]

median ages of 32 and 30 years, lower than ours for patients living with HIV. Concerning co-infected patients, a study carried out in Burkina Faso showed a median age of 38 years [2].

HBsAg carriage among PLHIV was 12.25% (n=18). Similar studies have been observed previously in Dakar [5] but also in other countries such as Benin [6] and Ghana [7] and results were respectively 16.8%, 16.9% and 16.7% higher than ours. On the other hand, a Botswana study conducted in 2011 reported a prevalence of co-infection of 5.3% [13].

In our study, no HIV/HBV/HCV co-infection was observed.-

For the treatment regimen according to WHO recommendations, which recommends a regimen containing Tenofovir (TDF) and Lamivudine (3TC) or Emtricitabine (FTC), the-

- Formatted: Font: 12 pt, Condensed by 0.15 pt
- Formatted: Font: 12 pt, Condensed by 0.1 pt
- Formatted: Font: 12 pt, Condensed by 0.15 pt
- Formatted: Font: 12 pt, Condensed by 0.1 pt
- Formatted: Font: 12 pt, Condensed by 0.15 pt
- Formatted: Font: 12 pt, Expanded by 0.15 pt
- Formatted: Font: 12 pt, Condensed by 0.15 pt
- Formatted: Font: 13 pt, Font color: Auto
- Formatted: Body Text, Left, Line spacing: single
- Formatted: Font color: Auto
- Formatted: Font: 12 pt, Expanded by 1.8 pt
- Formatted: Font: 12 pt, Expanded by 1.75 pt
- Formatted: Font: 12 pt, Expanded by 1.75 pt
- Formatted: Font: 12 pt, Expanded by 1.9 pt
- Formatted: Font: 12 pt, Expanded by 1.9 pt
- Formatted: Font: 12 pt, Expanded by 1.7 pt
- Formatted: Font: 12 pt, Expanded by 1.95 pt
- Formatted: Font: 12 pt, Expanded by 1.8 pt
- Formatted: Font: 12 pt, Expanded by 1.75 pt
- Formatted: Font: 12 pt, Expanded by 1.95 pt
- Formatted: Font: 12 pt, Condensed by 3 pt
- Formatted: Font: 12 pt, Expanded by 0.75 pt
- Formatted: Font: 12 pt, Expanded by 0.75 pt
- Formatted: Font: 12 pt, Expanded by 0.7 pt
- Formatted: Font: 12 pt, Expanded by 0.65 pt
- Formatted: Font: 12 pt, Expanded by 0.7 pt
- Formatted: Font: 12 pt, Expanded by 0.75 pt
- Formatted: Font: 12 pt, Expanded by 0.65 pt
- Formatted: Font: 12 pt, Expanded by 0.6 pt
- Formatted: Font: 12 pt, Expanded by 0.75 pt
- Formatted: Font: 12 pt, Expanded by 0.8 pt

most commonly used combination in our study was Tenofovir+Lamivudine + Efavirenz (77.77%) and 83% of the HIV/HBV co- infected patients in the active file were on treatment with this combination that is active against both HIV and HBV. The remaining 17% did not receive the combination recommended by the WHO, they were on treatment with only one molecule (lamivudine) which was also active against both viruses. The others had their treatment adapted as soon as the presence of HBsAg was discovered. Bado in Burkina Faso found that 68.5% of patients were on a single drug regimen which was either lamivudine or emtricitabine with 30.7% of patients not on treatment. [2]-[2]. This disparity is due to the fact that in most ARV protocols in Sub-Saharan african countries, lamivudine is the only active molecule available against HBV. Matthews G showed that after four (4) years of triple ARV therapy with a single molecule active against HBV, such as Lamivudine, 94% had resistance to the molecule [10]-as Lamivudine, 94% had resistance to the molecule Error! Bookmark not defined.. These results, which differ from ours, can be explained by the fact that in Senegal, since 2006, Tenofovir, Lamivudine and Emtricitabine are part of the treatment of any patient co-infected with HIV- HBV in accordance with WHO recommendations.

#### 5. Limitations of our study-

The diagnosis of HBV infection was made by testing for HBsAg. This method carries a risk of minimising prevalence because of the occult hepatitis B increasingly encountered in patients living with HIV. Detection of HBV viral load or PCR would have been the best options to take into account occult hepatitis B

#### 6. Conclusion

In Senegal, co-infection with the hepatitis B virus among patients living with HIV is not an uncommon phenomenon, since 12.25% of patients in our study were concerned. However, no co-infection with HCV was observed in these patients. Thus, the systematisation of HBV and HCV screening should allow for the adaptation of therapeutic regimens, especially in populations at risk.

#### Bibliographic References

1. Ba P S, Niang A, Fall A K, et al. B 01 : La co infection VIH/VHB/VHC à l'hôpital principal de Dakar. Médecine et Maladies Infectieuses. 2014;44:29.-
2. Bado G, Penot P, N'Diaye MD, et al. Hepatitis B seroprevalence in HIV infected patients consulting in a public day care unit in Bobo Dioulasso, Burkina Faso. Médecine et Maladies Infectieuses. 2013;43(5):202-7.-

Formatted: Font: 12 pt

Formatted: Body Text, Left, Indent: Left: 0.15", Right: 0.55", Space Before: 3.8 pt

Formatted [96]

Formatted [97]

Formatted: Body Text, Indent: Left: 0.15", Right: 0.6", Space Before: 6.85 pt

Formatted [98]

Formatted: Font: 12 pt, Font color: Auto

Formatted: Heading 1, Justified, Indent: Hanging: 0.25", Line spacing: single, Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.15" + Indent at: 0.4", Tab stops: 0.4", Left

Formatted: Font: 12 pt, Character scale: 100%

Formatted: Body Text, Indent: Left: 0.15", Right: 0.6", Space Before: 6.95 pt

Formatted [99]


Formatted: Font: 13 pt, Not Bold, Italic, Font color: Auto

Formatted: Body Text

Formatted: Heading 1, Space Before: 8.9 pt

Formatted [100]

3. Balogun TM, Emmanuel S, Ojerinde EF. HIV, Hepatitis B and C viruses' coinfection among patients in a Nigerian tertiary hospital. *Pan Afr Med J.* 2012;12(1)
4. Bourlière M, Fontaine H, Yazdanpanah Y, et al. Co-infection avec les hépatites virales. *J Afr Hepato Gastroenterol.* 2007;1:38-50.
5. Diop Ndiaye H, Touré Kane C, Etard JF, et al. Hepatitis B, C seroprevalence and delta viruses in HIV-1 Senegalese patients at HAART initiation (retrospective study). *J of Med Virol.* 2008;80(8):1332-1336.
6. Dovonou CA, Amidou SA, Kpangon AA, et al. Prévalence de l'hépatite B chez les personnes infectées par le VIH à Parakou au Bénin. *Pan African Med J.* 2015;20
7. Geretti AM, Patel M, Sarfo FS, et al. Detection of Highly Prevalent Hepatitis B Virus Coinfection among HIV Seropositive Persons in Ghana. *J of Clin Microbiol. American Society for Microbiology Journals.* 2010;48(9):3223-3230.
8. Jacquet, A., Wandeler, G., Nouaman, M., et al. HIV and Liver Fibrosis among Prison Inmates. 2015.
9. Ladep NG, Agaba PA, Agbaji O, et al. Rates and impact of hepatitis on human immunodeficiency virus infection in a large African cohort. *World J Gastroenterol.* 2013;19(10):1602-1610.
10. Matthews GV, Bartholomeusz A, Locarnini S, et al. Characteristics of drug-resistant HBV in an international collaborative study of HIV HBV infected individuals on extended lamivudine therapy. *AIDS.* 2006;20(6):863-870.
11. Modi AA, Feld JJ. Viral hepatitis and HIV in Africa. *AIDS Rev.* 2007;9(1):25-39.
12. OMS. Hépatite B : principaux repères. World Health Organ. 2018. Consulté le 29/4/18 Disponible <http://www.who.int/fr/news-room/fact-sheets/detail/hiv-aids>.
13. Patel P, Davis S, Tolle M, et al. Prevalence of Hepatitis B and Hepatitis C Coinfections in an Adult HIV Centre Population in Gaborone, Botswana. *The Am J of Trop Med and Hyg.* 2011;85(2):390-394.
14. Sagoe KWC, Agyei AA, Ziga F, et al. Prevalence and impact of hepatitis B and C virus co-infections in antiretroviral treatment naïve patients with HIV infection at a major treatment center in Ghana. *J of Med Virol.* 2012;84(1):6-10.
15. Sall Diallo A, Sarr M, Fall Y, et al. Étude de l'infection par le virus de l'hépatite virale B dans la population infantile au Sénégal. *Dakar Med.* 2004;136-42.
16. Wondimeneh Y, Alem M, Asfaw F, et al. HBV and HCV seroprevalence and their correlation with CD4 cells and liver enzymes among HIV positive individuals at University of Gondar Teaching Hospital, Northwest Ethiopia. *Virol J.* 2013;10(1):1-8.



[3. Balogun TM, Emmanuel S, Ojerinde EF. HIV, Hepatitis B and C viruses' coinfection among patients in a Nigerian tertiary hospital. Pan Afr Med J. 2012;12\(1\)](#)

[4. Bourlière M, Fontaine H, Yazdanpanah Y, et al. Co-infection avec les hépatites virales. J Afr Hepato Gastroenterol. 2007;1:38-50.](#)

[5. Diop-Ndiaye H, Touré-Kane C, Etard JF, et al. Hepatitis B, C seroprevalence and delta](#)

[viruses in HIV-1 Senegalese patients at HAART initiation \(retrospective study\). J of Med Virol. 2008;80\(8\):1332-1336.](#)

[6. Dovonou CA, Amidou SA, Kpangon AA, et al. Prévalence de l'hépatite B chez les personnes infectées par le VIH à Parakou au Bénin. Pan African Med J. 2015;20](#)

[7. Geretti AM, Patel M, Sarfo FS, et al. Detection of Highly Prevalent Hepatitis B Virus Coinfection among HIV-Seropositive Persons in Ghana. J of Clin Microbiol. American Society for Microbiology Journals. 2010;48\(9\):3223-3230.](#)

[8. Jacquet, A., Wandeler, G., Nouaman, M., et al. HIV and Liver Fibrosis among Prison Inmates. 2015.](#)

[9. Ladep NG, Agaba PA, Agbaji O, et al. Rates and impact of hepatitis on human immunodeficiency virus infection in a large African cohort. World J Gastroenterol. 2013;19\(10\):1602-1610.](#)

[10. Matthews GV, Bartholomeusz A, Locarnini S, et al. Characteristics of drug resistant HBV in an international collaborative study of HIV-HBV-infected individuals on extended lamivudine therapy. AIDS. 2006;20\(6\):863-870.](#)

[11. Modi AA, Feld JJ. Viral hepatitis and HIV in Africa. AIDS Rev. 2007;9\(1\):25-39.](#)

[12. OMS. Hépatite B : principaux repères. World Health Organ. 2018. Consulté le 29/4/18 Disponible <http://www.who.int/fr/news-room/fact-sheets/detail/hiv-aids>.](#)

[13. Patel P, Davis S, Tolle M, et al. Prevalence of Hepatitis B and Hepatitis C Coinfections in an Adult HIV Centre Population in Gaborone, Botswana. The Am J of Trop Med and Hyg. 2011;85\(2\):390-394.](#)

[14. Sagoe KWC, Agyei AA, Ziga F, et al. Prevalence and impact of hepatitis B and C virus co-infections in antiretroviral treatment naïve patients with HIV infection at a major treatment center in Ghana. J of Med Virol. 2012;84\(1\):6-10.](#)

Formatted: Font: 11 pt, Not Italic, Font color: Auto

15. Sall Diallo A, Sarr M, Fall Y, et al. Étude de l'infection par le virus de l'hépatite virale B dans la population infantile au Sénégal. *Dakar Med.* 2004;136-42.

16. Wondimeneh Y, Alem M, Asfaw F, et al. HBV and HCV seroprevalence and their correlation with CD4 cells and liver enzymes among HIV positive individuals at University of Gondar Teaching Hospital, Northwest Ethiopia. *Viro J.* 2013;10(1):1-8.

Formatted: Font: 11 pt, Not Italic, Font color: Auto

Formatted: Font: 12 pt

Formatted: Left: 0.83", Right: 0.35", Top: 1.11", Bottom: 0.6", Header distance from edge: 0", Footer distance from edge: 0.47"

Formatted: Font: 8.5 pt, Bold

Formatted: Body Text, Space Before: 0.2 pt

Formatted: Right: 0"