

Original Research Article

Evaluation of Phytochemical and Antibacterial Potential of *Vernonia amygdalina* and *Elaeis guineensis* on some Bacteria Associated with Diarrhea

ABSTRACT

Background: Bitter leaf and palm oil combination in treating diarrhoea is common in Africa. The aim of this study is to determine the synergistic and antimicrobial potential of *Vernonia amygdalina* and *Elaeis guineensis* on bacteria isolates from diarrheic stool.

Comment [u1]: Unbold

Methodology: One hundred and fifty (150) diarrheic stool samples from the Diagnostic Laboratory in Rivers State University Teaching Hospital, Port Harcourt, were inoculated in Selenite F broth, MacConkey, Nutrient, and Salmonella-Shigella agar plates using standard methods. Fresh and dry bitter leaf extracts were prepared by mashing, weighing and dissolving 100g and 150grams(g) concentrations of each in 100milliliters(mls) of ethanol and sterile distilled water respectively. Undiluted mixture of bitter leaf and palm oil were prepared by mixing 2mls of each in a sterile beaker. Zero-point one milliliter(0.1ml) of each undiluted extracts, their mixture, ethanol and palm oil were separately dispensed into appropriately labeled wells using sterile pipette for sensitivity test on the isolated bacteria. The pH, temperatures and the phytochemical contents of the bitter extracts and palm oil were determined using pH meter, thermometer and spectrophotometric method.

Comment [u2]: Remove comma

Comment [u3]: Space between 2 and mis

Results: *Escherichia coli*, *Staphylococcus aureus* and *Salmonella sp.* were isolated. The pH and temperature of bitter leaf extract and palm oil were 7.2 and 26°C and 6.7 and 26°C respectively while the pH and temperature for bitter leaf and the palm oil mixture was 6.9 and 26°C. The bitter leaf and palm oil extracts, their combination, ethanol and distilled water used on the isolates as antibiotics did not show any sensitivity. The phytochemicals identified in the bitter leaf were saponins, alkaloids and tannins while carotene was detected in the palm oil.

Comment [u4]: Remove comma

Comment [u5]: 26°C

Conclusion: It can be concluded that the bitter leaf extract, palm oil and their combination did not inhibit the growth of the bacteria as a result of lack of bioactive components from the bitter leaf. The nature of the diluents used for the extraction may not have been the right one as it would have caused the low concentration, dilution and disfiguring of the molecules of the bioactive compounds that prevented the proper reaction between the bacteria and the extracts used.

Keywords: Bacteria, Extracts, Diarrhoea, Bitter leaf, Palm oil

1. INTRODUCTION

Plant extracts are continuously being sought for as effective and cheaper alternative sources of medication all over the world. The use of plants extracts in the treatment of human ailment is an ancient art, a practice that has passed on from generation to generation. Research into traditional plants received further boost due to the increasing resistance to many orthodox drugs and thus, a search for new organic molecules from plants with antimicrobial properties [1]. Bitter leaf is a small tree or a shrub belonging to the family of *Asteraceae*, and it is a popular vegetable in Africa which grows in several parts of subtropical and tropical Africa [2]. It is known as “Ewuro” in Yoruba, “Onugbu” in Igbo, “Chusar-doki” in Hausa, “Oriwo” in Edo, “Etidot” in Ibibo, “Ityuna” in Tiv, and ‘Grawa” in Amharic. In other parts of Africa, it is also known as *Ndole* or *Muop* (Cameroon), *Tuntwono* (Tanzania), and *Mululuza* (Uganda) [3]. Palm oil is always extracted from two types of oil palm fruit: *Elaeis guineensis*, which is common in African and *Elaeisoleifera*, which is found in South America. Historical accounts suggest that palm oil was a part of the diet of indigenous populations. Presently, it has become the second most traded oil crop in the world, after soy, with Malaysia and Indonesia as its main producers. Palm oil is not to be confused with palm kernel oil. Both are obtained from the fruit, but the palm kernel oil is gotten from the seeds. Palm kernel oil has a higher amount of saturated fat, which makes it ideal for cooking [4]. Palm oil, also called red oil, contains high amount of saturated fat, antioxidants and vitamins, and is used as cooking oil, its stability makes it easier to be stored at room temperature for many months. It is also used as an ingredient in soups and sauces, dietary supplement, personal care and household products, skin moisturizer, sun block and biofuel. Diarrhea is an abnormal increase in the frequency of emptying waste matter from the bowel. It is characterized by loose, watery stools, and usually last a few days [5]. Signs of dehydration often begin with loss of the normal stretchiness of the skin and irritable behavior. This can progress to decreased urination, loss of skin colour, fast heart rate, and a decrease in responsiveness as it becomes more severe. Diarrhea can be caused by viruses, bacteria or parasite, a condition also known as gastroenteritis. *Campylobacter sp* are common cause of bacterial diarrhea, but infections by *Escherichia coli*, *Salmonella sp*, *Shigella sp* and *Staphylococcus aureus* are also a frequent cause. These infections are often acquired from water or food that has been contaminated by faeces. The presence of resistant genes which results in multi drug resistance in diarrhoea treatment calls [6] for other treatment options of diarrhoea caused by microorganisms. Phytochemicals are the active biological component of plants which include alkaloids, tannins, terpenoids and flavonoids[8]. Some plant extracts inhibit peptidoglycan synthesis, damage microbial membrane structures, modify surface hydrophobicity and modulate quorum-sensing [8]. This research is designed to determine the synergistic and antibacterial effect of bitter leaf (*Veronica amygdalina delile*) and palm oil (*Elaeis guineensis* Jacq) on some organisms associated with diarrhea

2. MATERIALS AND METHOD

2.1 Source of Plant Materials

Fresh leaf sample of bitter leaf and palm fruits were purchased from Mile 3 General Market, Diobu in Port Harcourt, Nigeria. [The plant and palm fruits were identified by a Plant Taxonomist

in the Department of Plant Science and Biotechnology, Faculty of Science, Rivers State University, Port Harcourt.

2.2 Preparation of Plant Extracts

2.2.1 Fresh Leaves extract

The fresh bitter leaf plant was used to prepare three extracts. The leaf sample was rinsed three times with clean water. The fresh leaves were then mashed using sterile mortar and pestle in Medical Microbiology Laboratory, Faculty of Science, Rivers State University, Port Harcourt. Some portions of the leaf were shade-dried for few days and ground with a sterile surface grinder into powdery form and stored in clean tight-capped bottle labeled and dated [9]. One hundred gram and one hundred and fifty grams (100g and 150g) of each of the ground fresh leaves were weighed in three batches. The first batch was soaked in [100ml] of ethanol, stoppered with cotton wool, covered with aluminum foil and left to stand for 24hours in the refrigerator. This was then filtered using Whatman's No 1 filter paper and the extract stored in the refrigerator. The second batch of 100g and 150g of ground fresh bitter leaf were soaked in 100mls of water, stoppered with cotton wool, covered with aluminum foil and left to stand for 24hrs in the refrigerator. This was filtered using Whatman's No 1 filter and stored in the refrigerator. The third batch of the 100g and 150g of the ground fresh bitter leaf were placed in a sterile muslin cloth respectively and squeezed out the direct liquid from them individually into separate clean sterile containers. These were filtered with Whatman's No 1 filter paper and decanted into a sterile bottle that were properly labelled and stored in the refrigerator for use [9].

Comment [u7]: space

Comment [u8]: space and consistency- Use either hours or hrs

2.2.2 Dry Leaves Extracts

One hundred grams (100g) and one hundred and fifty grams (150g) separately of ground powder of the dry bitter leaf were each weighed and placed into four (4) separate sterile conical flasks of [250ml] capacity to be used for ethanol and water extractions separately. One hundred ([100ml]) milliliter of absolute ethanol and water were separately measured and added to each conical flask that has been properly labeled respectively. These were shaken properly as the mixtures were left to stand at room temperature for [24hrs]. Each was then filtered by passing it through a muslin cloth and then filtered with Whatman's No 1 filter paper and each filtrate was stored in the refrigerator for further use [9].

Comment [u9]: space

Comment [u10]: space

Comment [u11]: space and be consistent- hours or hrs?

2.2.3 Palm Oil Extraction

The Palm fruits were washed thoroughly in clean water and boiled to soften the skin. It was then mashed in a mortar to separate the pulp from the nuts. The whole mass was immersed in water, stirred and the palm oil was skimmed off. The fiber was sifted out of the water and the nuts were collected and separated from the remaining fiber. The palm oil obtained was then boiled and the palm oil was decanted into a sterilize reagent bottle and kept in the refrigerator for further use [10].

2.3 Collection and processing of Samples

Stool samples were collected from patients presenting with gastroenteritis at Rivers State University Teaching Hospital (RSUTH), Port Harcourt. A total of one hundred and fifty samples were processed for pathogenic bacteria, using standard protocols of stool bacteriology as given by Chesbrough, [10]. All identified bacteria were tested for antimicrobial susceptibility using the different obtained extracts.

Comment [u12]: remove comma

2.4 Bacterial Sensitivity Testing Using Agar Well Diffusion Technique

The fresh and dried leaf extracts of bitter leaf were subjected to bacterial sensitivity test. Two dried agar plates that were properly labelled were used for the testing, one for the prepared fresh leaves and the other for the prepared dried leaves. One milliliter (1.0ml) of each was dispensed onto the dried surface of a sterile nutrient agar plates. They were evenly distributed and allowed to dry but properly covered to avoid atmospheric contamination. The plates were bored with a sterile cork borer of about 8mm. Zero-point one milliliter (0.1ml) of each extract from the fresh and dried bitter leaf of different concentration were aseptically dispensed into the wells of about 8mm deep including both positive and negative control (synthetic agent and ethanol) [12]. The wells were labelled as follows Well 1 – Ethanol alone, Well 2 – Palm oil alone, Well 3 – Water extract of fresh bitter leaf, Well 4 - Water extract of dry bitter leaf, Well 5 – Fresh bitter leaf liquid, Well 6 – Ethanol extract of fresh bitter leaf, Well 7- Ethanol extract of dry bitter leaf and Well 8 – mixture of fresh bitter leaf liquid and palm oil

In well 8, 2mls of the bitter leaf crude extract and palm oil were mixed together and zero-point one milliliter (0.1ml) of the mixture was dispensed into the well. Ciprofloxacin, an orthodox antimicrobial agent was used as positive control. The plates were then incubated for 24hours at 37°C. The sensitivity of the extracts and the control were observed [12].

Comment [u13]: space between 8 and mm

Comment [u14]: space

Comment [u15]: 34 hrs or 24 hours?? Which?

2.5 Phytochemical Screening

Phytochemical screening was performed on the crude extract of bitter leaf for detection of alkaloids (Dragendorff test), flavonoids (Shinoda's test), tannins (lead acetate test), saponins (Rothing test), glycosides, (Liebermann-Burchard test) and carotenes. These were carried out using standard methods as described by Ehiowenwengeon *et al.*, [13].

Comment [u16]: Delete comma

2.6 Determination for pH and Temperature of Bitter leaf and Palm Oil and their Mixture

2.6.1 pH

With the aid of small pHep pocket sized pH meter (Made by Hanna Instrument). A standard buffer solution of known pH of 4 and 7 was used to calibrate the instrument after which the pH of the sample was taken.

2.6.2 Temperature

A mercury thermometer was placed into the sample and allowed to stay for five minutes for the stability of the temperature after which the reading was taken.

Comment [u17]: of

3. RESULTS

The inhibitory effect of ethanol and water extracts of bitterleaf isolated from diarrheic stool samples is represented in table 1 while table 2 shows the inhibitory effect of direct bitter leaf liquid and palm oil on bacteria isolated from diarrheic stool samples.

Table:1 The Inhibitory Effect of Ethanol and Water Extracts of Bitter Leaf on Bacteria Isolated from Diarrheic Stool Samples

Concentration (g/ml)

Extracts /Antibiotic	Bacterial Isolates	Concentration (g/ml)	
		100g/100ml	150g/100ml
Ethanol (fresh leaves)	<i>Escherichia coli</i>	Nil	Nil
	<i>Salmonella sp</i>	Nil	Nil
	<i>Staphylococcus aureus</i>	Nil	Nil
Water (fresh leaves)	<i>Escherichia coli</i>	Nil	Nil
	<i>Salmonella sp</i>	Nil	Nil
	<i>Staphylococcus aureus</i>	Nil	Nil
Ethanol (dry leaves)	<i>Escherichia coli</i>	Nil	Nil
	<i>Salmonella sp</i>	Nil	Nil
	<i>Staphylococcus aureus</i>	Nil	Nil
Water (dry leaves)	<i>Escherichia coli</i>	Nil	Nil
	<i>Salmonella sp</i>	Nil	Nil
	<i>Staphylococcus aureus</i>	Nil	Nil
Ciprofloxacin (control)	<i>Escherichia coli</i>	S	S
	<i>Salmonella sp</i>	S	S
	<i>Staphylococcus aureus</i>	S	S

Nil = No Inhibition, S = Sensitive

Table: 2 The Inhibitory Effect of Direct Bitter Leaf Liquid and Palm Oil on Bacteria Isolated from Diarrheic Stool Samples

Extracts /Antibiotic	Bacterial Isolates	Concentration (g/ml)
		Neat

Fresh Palm Oil	<i>Escherichia coli</i>	Nil
	<i>Salmonella sp</i>	Nil
	<i>Staphylococcus aureus</i>	Nil
Fresh Bitter Leaf and Palm oil Combination	<i>Escherichia coli</i>	Nil
	<i>Salmonella sp</i>	Nil
	<i>Staphylococcus aureus</i>	Nil
Raw Bitter Leaf Liquid	<i>Escherichia coli</i>	Nil
	<i>Salmonella sp</i>	Nil
	<i>Staphylococcus aureus</i>	Nil
Ciprofloxacin (Control)	<i>Escherichia coli</i>	S
	<i>Salmonella sp</i>	S
	<i>Staphylococcus aureus</i>	S

Key:

Nil=No inhibition, S=Sensitive

Table 3 on the other hand is used to represent the quantitative screening of active phytochemical compounds of bitter leaf extract and palm oil and table 4 represents the mean value of pH and temperature of bitter leaf, palm oil and their mixture.

Table: 3 Qualitative Screening of Active Phytochemical Components of Bitter Leaf Extract and Palm oil

Active Components

Extracts	Saponins	Alkaloid	Tannins	Steroids	Flavonoid	Phenols	Carotenes
Ethanol (fresh leaves)	+	+	+	Nil	Nil	Nil	Nil
Water (fresh leaves)	+	+	++	Nil	Nil	Nil	Nil
Ethanol (dry leaves)	Nil	+	+	Nil	Nil	Nil	Nil
Water (dry leaves)	Nil	+	+	Nil	Nil	Nil	Nil
Raw Bitter leaf(fresh)	+	+	+	Nil	Nil	Nil	Nil
Raw Bitter leaf (dry)	+	+	Nil	Nil	Nil	Nil	Nil
Palm Oil	Nil	Nil	Nil	Nil	Nil	Nil	+
Control							
Ethanol	Nil	Nil	Nil	Nil	Nil	Nil	Nil
Water	Nil	Nil	Nil	Nil	Nil	Nil	Nil

Key

Nil = Not detected, + = Moderately detected, ++ = Absolutely detected

Table: 4 Mean Value of pH and Temperature of Bitter Leaf, Palm Oil and their Mixture

S/No	Sample	Parameters	
		pH	Temperature (°C)
1	Butterleaf	7.2	26
2.	Palm oil	6.7	26
3.	Butterleaf and palm oil mixture	6.9	26

4. DISCUSSION

The effect of bitter leaf and palm oil on *Escherichia coli*, *Salmonella sp.*, and *Staphylococcus aureus* were investigated. Out of the one hundred and fifty (150) diarrheic stool samples obtained from Rivers State University Teaching Hospital (RSUTH), thirty (30) bacteria were isolated. Eleven (11) samples yielded *Escherichia coli*, whereaseight (8) samples yielded *Salmonella sp* while *Staphylococcus aureus* were isolated from eleven (11) samples. The research work revealed that *Escherichia coli*, *Salmonella sp* and *Staphylococcus aureus* were not susceptible to either of the extracts as shown in table 1. The result obtained in the work agrees with the observation of Eloff [14], who reported that the aqueous extracts of bitter leaf showed no inhibition on *Escherichia coli*, *Salmonella sp* and *Staphylococcus aureus*. He also reported that the most active ingredients in bitter leaf are not water soluble. Eloff [14] examined a variety of extracts for the ability to solubilize antimicrobial agents from plants and observed that the most commonly used solvent (ethanol and methanol) may not demonstrate the greater sensitivity in yielding antimicrobial chemicals on an initial screening.

According to Ibrahim *et al.*, [15], the antibacterial susceptibility test of *Vernonia amygdalina* and *Elaeis guineensis* showed that the aqueous extract of the plants showed low inhibition on food borne pathogens such as *Escherichia coli*, *Bacillus cereus*, *Shigella dysenteriae* and *Salmonella typhimurium* as shown in table 2. Ali *et al.*, [16] revealed that the ethanolic and aqueous extract of bitter leaf were inactive against diarrhea causing organism such as *Escherichia coli*, *Salmonellae sp* and *Shigella sp*. Evbuomwan *et al.*, [17] investigated the antibacterial activity of *Vernonia amygdalina* leaf extracts against multidrug resistant bacterial isolates. He reported that *Escherichia coli* was observed to be completely resistance to all concentration of the plant extracts.

Uzoigwe and Agwa [18] carried out research on the antimicrobial activity of *Vernonia amygdalina* on selected urinary tract pathogen. The isolates were *Klebsiellasp*, *Escherichia coli* and *Staphylococcus sp*. The bacterial isolate that showed different levels of susceptibility to both leaf and stem extracts of *Vernonia amygdalina* was *Klebsiella sp* while *Staphylococcus sp* and *Escherichia coli* were not susceptible to either of the extracts. According to Simplice *et al.*, [19], three Gram negative bacteria; *Escherichia coli*, *Klebsiella pneumonia* and *Acinetobactersp* were tested for their susceptibility to some antibiotics, the highest resistance rates were recorded with *Escherichia coli* been the highest at 76.64%.The obtained values from the determination of the pH and temperature of the extracts showed that the pH ranged from 6.9 to 7.2 whereas the temperature of the extracts was 26° as is indicated on table 4. These are not extreme pH that

Comment [u18]: delete comma

Comment [u19]: 76.6 %

might prevent the growth of bacteria. This is because the isolated bacteria can grow at that pH since they are neutrophiles. The temperature of the different extracts was 26°C and the incubation temperature for these bacteria was 37°C. It could be that the temperature at which these extracts were introduced to the bacteria could have brought a sudden shock on the bacteria thereby making the bacteria cells to shrink and caused no reaction between the cell and the plant extracts. The determination of phytochemicals agrees with the work of Ojinnaka *et al.*, 2016 that supported the presence of saponins, tannins, and alkaloids to be present in *Vernonia amygdalina*. The method used in the extraction of the bioactive ingredient was mashing and extracting with water and ethanol which did not produce enough of the ingredients which agrees with the work of Ojinnaka *et al.*, [20], which stated that blanching and abrasive washing reduces the level of nutrients in the plant extract and low amounts of these nutrients, such as tannins, saponins and alkaloids were observed. In conclusion the study has shown that the aqueous, ethanol and direct bitter leaf liquid and palm oil were not effective on the bacterial isolates used. The bacterial isolates were susceptible to the ciprofloxacin antibiotics disc compare to the extract that showed no effectiveness against *Escherichia coli*, *Salmonella Sp* and *Staphylococcus aureus*.

Comment [u20]: Delete. You are using Vancouver referencing

5. CONCLUSION:

It can be concluded that the bitter leaf extract, the palm oil extract and their combination did not inhibit the growth of the bacteria used due to low level of extracted bioactive components from the bitter leaf. However, the nature of the diluents used for the extraction could have caused the low concentration, dilution and disfiguring of the molecules of the bioactive compounds that prevented the proper reaction between the bacteria and the extracts used.

Comment [u21]: In your discussion, you include the public health implications of your results to make the discussion complete

Comment [u22]: Is this not conclusion? But you still had another conclusion

Comment [u23]: Delete the colon

Comment [u24]: This is another conclusion. It seems you are still discussing your results here again which been captured under discussion

REFERENCE

- [1] Sofowora, E.A. (1993). *Medicinal Plants and Medicine in Africa* (2nd eds.) John Wiley and sons, New York, p. 6.
- [2] Erasto, P., Grierson, D.S., & Afolayan, A.J. Bioactive Sesquiterpene lactones from the leaves of *Vernonia amygdalina*. *Journal of Ethnopharmacology*. 2006; 106(2), 117-120.
- [3] Owolabi, M.A., Jaja, S.L., Oyekanmi, O.O. & Olatunji, O. J. Evaluation of the antioxidant activity and lipid peroxidation of the leaves. Of *Vernonia amygdalina*. *Journal of Complementary Integrative Medicine*. 2008; 5(1), 1550-13840.
- [4] Poku, K. Small- Scale Palm Processing in Africa. Food and Agriculture organization of the UN. Rome. 2002.
- [5] Bibitha, B; Jisha, V.K; Salitha, C.V; Mohan, S. and Valsa, A.K. Antibacterial activity of different plant extracts. *Indian Journal of Microbiology*. 2002; 42; 361-363.
- [6] Agi, V. N., Abbey, S. D., Wachukwu, C. C. K., Nwokah, E. G., & Ollor, O. A. Some resistant genes associated with Diarrhoea in Rivers State. *European Journal of Pharmaceutical and Medical Research*. 2007; 4(10), 116-121.

Comment [u25]: Remove from here. You are not using APA

Comment [u26]: Is this supposed to be comma or colon? Be consistent with the journal format pls

Comment [u27]: Cross check this

Comment [u28]: Microbiology Why are some journal names in italics and some unitalicized? Conform to the journal guidelines and correct appropriately

Comment [u29]: Colon or comma??

- [7] Kenny, C.R. & Furey, A. A post-antibiotic era looms: can plant natural product Research fill the void. *British Journal of Biomedical Science*. 2015;72(4), 191-200.
- [8] Cushnie, T.P., Tim, C., Benjamart, L. & Andrew, J. Alkaloids: an overview of their antibacterial, antibiotic- enhancing and ant virulence activities. *International Journal of Antimicrobial Agents*. 2014; 44(5), 377-386.
- [9] Adetunji, C. O., Olaniyi, O.O. & Ogunkunle, A.T.J. Bacterial activity of crude extracts of *Veronica amygdalina* on clinical isolates. *Journal of Microbiology and Antimicrobials*. 2013; 5(6), 60-64.
- [10] Ekwenye, U.N. & Ijeoma, A. "Antimicrobial effects of palm kernel oil and palm oil", *KMITEL Sci., J. Vol. 5(2)*, pp. 502-505, 2005
- [11] Cheesbrough, M. *District Laboratory practice in Tropical Countries*, Part 11, (2nd edition), Cambridge, UK. Cambridge University Press. 2000; 67-70.
- [12] Oshilim, A.O. Comparative study of the effect of bitter lead extracts and antibiotics (gentamycin and amoxicillin) on bacterial species isolated from wound. *Journal of Bacteriology & Pharmaceutical Research*. 2017; 6(1), 009-012.
- [13] Ehiowenguan, G., Emoghene, A. O. & Inetianbor, J.E. Antibacterial and photochemical analysis of Peeledfruit. *International Organization of Scientific Research*. 2014; 4(8) 18-25.
- [14] Eloff, J.N. A sensitive and quick microplate method to determine the minimum inhibitory concentration of plant extracts for bacteria. *Plant Medicine*. 1998; 60(9), 1-8.
- [15] Ibrahim, T.A., Ajala, L., Adetuyi, F.O. Ojei, J.B. Assessment of the Antibacterial activity of *Vernonia amygdalina* and *Occimum gratissimum* leaves in selected food borne pathogens. *International Journal of Third World Medicine*. 2009; 8(2), 1539-1545
- [16] Ali, M., Diso, S. U., Waiya, S. A., & Abdallah, M. S. Phytochemical screening and antibacterial activity of bitter leaf (*Vernonia amygdalina*). *Annals Microbiol. Infectious Dis*. 2019; 2(4), 01-07.
- [17] Evbuomwan, L., Chukwuka, E. P., Obazenu, E. I., & Ilevbare, L. Antibacterial activity of *Vernonia amygdalina* leaf extracts against multidrug resistant bacterial isolates. *Journal of Applied Sciences and Environmental Management*. 2018; 22(1), 17-21.
- [18] Uzoigwe, C.I. & Agwa O.K. The antimicrobial activity of *Vernonia amygdalina* on selected urinary tract pathogens. *African Journal of Microbiology Research*. 2011; 5(12), 1467-1472.
- [19] Simplice, D.K., Llnoudo, D.P., Wendyame, M.C., Nadembega, Y.A., Ajeneba, O., Silvatore, P., Virginio, P., Alfred, S.T., & Comlar de Souza, Jacques, S. Antimicrobial resistance of major urinary tract pathogen. *Pakistan Journal of Biology of Science*. 2000; 12(9), 712-712.

Comment [u30]: Correct all these

Comment [u31]: Comma or colon missing??

Comment [u32]: Insert full stop

Comment [u33]: Cross check this

[20] Ojinnaka, M. C., Odimegwu, E. N., & Chidiebere, F. E. Comparative study on the nutrient and antinutrient composition of the seeds and leaves of Uziza (*Piper guineense*). *IOSR Journal of Environmental Science, Toxicology and Food Technology*. 2016; 10(8), 42-48.

Comment [u34]: What is this???

UNDER PEER REVIEW