

## Original Research Article

# Optimization of Thermally Processed Bambara Groundnut Flour Nutrients Using Response Surface Methodology

### ABSTRACT

White/cream colour bambara groundnut (BGN) seed (*Vigna subterranean L.*) Was grouped into 12 portions and subjected to different processing methods by soaking in water at room temperature (28°C) for 12- 48h and further boiling for different times (30 – 60min). The soaked-boiled seeds were dried and milled into flours. Proximate composition and functional properties of these processed flours were determined and significant ( $P = .05$ ) differences among the samples recorded. Combination of soaking and boiling of the seeds for different times resulted to irregular functional properties and nutrient losses of the BGN. Optimization of the responses showed that the seed soaked for 12h and further boiled for 46.26min would yield flour with optimum nutrient properties, while soaking for 12h before boiling for 9.96 min would produce flour with optimum functional properties.

*Keywords: Proximate, Functional properties, Soaking, Boiling, Bambara groundnut flour*

### 1. INTRODUCTION

The cultivated Bambara groundnut (*Vigna subterranean L.*) is a pulse with subterranean fruit set, and is currently being cultivated by small farm holders [1], [2]. The Bambara groundnut (*Vigna subterranean L.*) belongs to the plantae of the family of *fabaceae* and subfamily of *faboidea*, [3]. Because of its composition, BGN produces almost a balanced diet with a reported good balance of essential amino acid; its lysine (6.8%) and methionine (1.3%), [4]; [5], [6] and [7]. Due to its high protein score of 80% and predicted protein energy ratio (P-PER) of above 2.00. BGN is regarded as of high protein quality, [8]. Its other nutrients include 60-65% carbohydrate, [7] 6.34% to 6.85% crude fibre [3], [9] and is also credited with good quantities of micronutrient and vitamins. BGN has also historically been part of inexpensive meals throughout the world [10].

Bambara groundnut is regarded as the third most important legume crop after groundnuts *Arachis hypogea* and cowpea, but it is still one of the lesser utilized and underexploited legume. Bambara groundnut has attracted very little scientific research and industrial utility, despite its nutritional and other reported potentials [11], therefore the objective of this study is to determine combination the better combination of two process variables: soaking time ( $X_1$ ) and boiling time ( $X_2$ ) that will give optimum quality of instant BGN flour, in terms of nutrition and functionality. The information generated from this study is expected prediction on its possible use in industrial processes. This will lead to diversification and utilization in addition to provision of enhanced food security.

Modeling and optimization are vital in product development. Also Response Surface Methodology (RSM) has been frequently used in optimization of food process variables [12]. Optimization of process conditions is one of the most critical stages in the development of any efficient and economic bioprocess, [13]. RSM is a mathematical and statistical technique for designing experiments, building models, evaluating the relative significance of several independent variables, in order to determine the optimum conditions for desirable responses, [14]. Therefore, the objective of this study is to determine through RSM the best combination of soaking time ( $X_1$ ) and boiling time ( $X_2$ ) that will give optimum quality of bambara groundnut (BGN) flour for nutrition and other processes.

### 2. MATERIAL AND METHODS

#### Source of Materials

White/cream colour dry BGN seeds were procured from the "Akwata" section of Ogbete market, Enugu, Enugu State, Nigeria.

**Comment [A1]:** Does the materials used only the BGN seed only? Since in the research also doing many analysis, supposedly the materials for analysis should be mentioned here include the materials supplier should be mentioned.

## 2.1 Experimental design

A Response Surface Methodology (RSM) involving two process variables at three levels was applied to study the effect of soaking times ( $X_1$ ) and boiling time ( $X_2$ ) on the resultant quality of the BGN flours. The proximate and the functional properties of the flours were the response variables measured.

Table 1 shows 2-factor-3- levels Face Centered Central Composite design (FCCCD) experimental matrix of interaction effect of soaking time (h) and boiling time (min) employed to produce and evaluate the proximate and functional properties of Bambara ground nut flour. Table 2 shows the design key used to generate Table 1

**Table 1: Experimental Design with both coded and actual values**

Run	Soaking time (h)	Boiling time (min)
	A	B
1	12 (-1.00)	30 (-1.00)
2	12 (-1.00)	45 (0.00)
3	12 (-1.00)	45 (0.00)
4	12 (-1.00)	60 (1.00)
5	30 (0.00)	30 (-1.00)
6	30 (0.00)	45 (0.00)
7	30 (0.00)	45 (0.00)
8	30 (0.00)	60 (1.00)
9	48 (1.00)	30 (-1.00)
10	48 (1.00)	30 (-1.00)
11	48 (1.00)	45 (0.00)
12	48 (1.00)	60 (1.00)

*Values in the brackets are the codes while those outside are the actual*

**Table 2: Codes and actual levels of the independent variable (or conditions) and their corresponding levels and times**

Independent Variables	Symbols	Coded levels		
		-1	0	+1
Soaking time	$X_1$	12h	30h	48h
Boiling time	$X_2$	30min	45min	60min

### Preparation of raw materials

The BGN seeds were cleaned through dry cleaning process to remove extraneous materials such as pieces of stones, strings, pieces of metals and spoilt seeds. The BGN seeds were washed with potable water and then divided into twelve portions. Each portion was subjected to soaking and

boiling as shown in Table 1. The categories of soaked and boiled samples were dried and then hammer milled to produce the BGN flours.

These 12 flour samples were later analyzed for proximate composition and functional properties. The results were presented in Tables 3 - 6 respectively.

### 2.1.1 ANALYSIS

#### Proximate Composition

The official method of [15] was used to determine the moisture, crude protein, lipid, ash and crude fibre contents of the Bambara groundnut flours. The carbohydrate was obtained by subtracting the summation of protein, fat, ash, crude fibre and moisture from 100

#### Functional Properties

Water absorption and oil absorption capacities; foam capacity and stability, gelation temperature, solubility and swelling power of the flours were determined using [16]

#### Statistical analysis

To determine the mean differences between treatments, the triplicate data were subjected to One Way Analysis of Variance (ANOVA). Duncan's multiple range tests was used to separate the means where differences existed using IBM SPSS, version 22. Design Expert version 12 was used to design the experiments and replications covering the full design of two factors were used for building quadratic models, which produced reasonable amount of information for testing the conditions for model adequacy and well suited for process optimization, (Myers *et al.*, 2009). Polynomial equations were determined and simplified based on the influence of the factors on the final response. The experimental data obtained from the model experiments can be represented in the form of the equation

$$Y = a + b_1X_1 + b_2X_2 + b_{12}X_1X_2 + b_1X_1^2 + b_2X_2^2 + e \dots \dots \dots (1)$$

A quadratic polynomial regression model was assumed for predicting various responses Montgomery, (2013). The model proposed for each response Y was represented as

$$Y = \beta_0 \sum_{i=1}^3 + \beta_{i \times 1} + \sum_{i=1}^3 \beta_2 X_i^2 + \sum_{i=1}^2 \sum_{j=i+1}^3 \beta_{ij} \times X_i \dots \dots \dots (2)$$

Regression models and response surface plots were developed after examining the ANOVA of various response variables to select the terms whose p-value are < 0.05, significant model, high R<sup>2</sup>adj and non-significant lack of fit. The coefficient of determinations of responses was used to fit the models for only the responses that qualified. Table 3 shows the proximate composition of the samples

## 3. RESULTS AND DISCUSSION

**Comment [A2]:** In the result and discussion, please add some introduction paragraph before showing the results.

**Table 3: Experimental responses of the samples under different soaking and boiling times**

Run	Factors		RESPONSES					
	X <sub>1</sub>	X <sub>2</sub>	%Pro	%Ash	%Fat	%C. fibre	%CHO	%Moisture
1	12 (-1.00)	30 (-1.00)	7.88±0.13 <sup>c</sup>	3.18±0.03 <sup>b</sup>	3.70±0.03 <sup>f</sup>	0.32±0.04 <sup>g</sup>	77.94±0.19 <sup>c</sup>	6.97±0.02 <sup>abc</sup>
2	12 (-1.00)	45 (0.00)	8.87±1.11 <sup>b</sup>	3.43±0.28 <sup>a</sup>	4.74±0.58 <sup>de</sup>	0.31±0.01 <sup>g</sup>	75.18±0.57 <sup>e</sup>	7.22±0.03 <sup>ab</sup>
3	12 (-1.00)	45 (0.00)	11.36±1.22 <sup>a</sup>	3.00±0.01 <sup>c</sup>	4.12±0.06 <sup>ef</sup>	1.13±0.12 <sup>bc</sup>	73.47±1.27 <sup>f</sup>	6.93±0.02 <sup>abc</sup>
4	12 (-1.00)	60 (1.00)	7.65±0.02 <sup>c</sup>	2.68±0.02 <sup>d</sup>	5.14±0.11 <sup>d</sup>	1.01±0.08 <sup>ode</sup>	76.36±0.03 <sup>d</sup>	7.31±0.25 <sup>a</sup>
5	30 (0.00)	30 (-1.00)	6.07±0.36 <sup>d</sup>	2.08±0.02 <sup>f</sup>	5.29±0.12 <sup>d</sup>	0.51±0.08 <sup>fg</sup>	79.02±0.54 <sup>c</sup>	7.01±0.11 <sup>abc</sup>
6	30 (0.00)	45 (0.00)	2.76±0.41 <sup>g</sup>	2.18±0.02 <sup>ef</sup>	5.06±0.02 <sup>de</sup>	0.74±0.21 <sup>ef</sup>	82.60±0.5 <sup>a</sup>	6.67±0.05 <sup>c</sup>
7	30 (0.00)	45 (0.00)	1.94±0.26 <sup>gh</sup>	2.29±0.03 <sup>e</sup>	7.37±0.17 <sup>c</sup>	1.09±0.12 <sup>cd</sup>	80.37±0.95 <sup>b</sup>	6.94±0.09 <sup>ab</sup>
8	30 (0.00)	60 (1.00)	1.97±0.12 <sup>gh</sup>	2.24±0.02 <sup>e</sup>	5.51±0.41 <sup>d</sup>	1.19±0.11 <sup>bc</sup>	82.48±0.99 <sup>a</sup>	6.63±0.56 <sup>c</sup>

9	48 (1.00)	30 (-1.00)	1.84±0.13 <sup>gh</sup>	2.18±0.00 <sup>ef</sup>	10.75±0.79 <sup>a</sup>	1.02±0.15 <sup>cde</sup>	76.80±1.10 <sup>d</sup>	5.15±0.16 <sup>d</sup>
10	48 (1.00)	30 (-1.00)	1.71±0.22 <sup>h</sup>	1.82±0.01 <sup>g</sup>	8.20±0.77 <sup>c</sup>	0.81±0.09 <sup>de</sup>	82.30±0.24 <sup>a</sup>	5.16±0.16 <sup>d</sup>
11	48 (1.00)	45 (0.00)	4.27±0.27 <sup>i</sup>	1.84±0.02 <sup>g</sup>	9.54±0.93 <sup>b</sup>	0.96±0.06 <sup>cde</sup>	76.60±0.04 <sup>d</sup>	6.80±0.62 <sup>bc</sup>
12	48 (1.00)	60 (1.00)	5.49±0.46 <sup>de</sup>	2.70±0.02 <sup>d</sup>	7.64±1.13 <sup>c</sup>	1.39±0.42 <sup>a</sup>	76.28±0.44 <sup>d</sup>	6.83±0.10 <sup>abc</sup>

Values are mean ± standard deviation of replicated determinations. Mean values in the same column followed by different letters are significantly different (P=0.05)

X<sub>1</sub> = Soaking time; X<sub>2</sub> = Boiling time, C. fibre = Crude fibre; CHO = Carbohydrate

Table 4 shows the Summary of ANOVA and Coefficient Estimate of the proximate composition of the Bambara groundnut flour for the terms that showed significant model, non-significant lack of fit and P-value < 0.05. The ideal regression equation showing the response variables (proximate parameters) as a function of the independent (process variables) can be represented in Eq. 1

$$Y = \beta_0 + \beta_1 A + \beta_2 B + \beta_{12} AB + \beta_{11} A^2 + \beta_{22} B^2 + \varepsilon \quad (1)$$

Where: Y= response variable,  $\beta_0$ =intercept, A= soaking time, B = boiling time,  $\beta_1, \beta_2, \beta_{12}, \beta_{11}, \beta_{22}$  = coefficients of the linear, interaction and square of A and B respectively,  $\varepsilon$  = estimated error.  $\varepsilon$  is often neglected and not factored into the equation


**Table 4: Summary of ANOVA and Coefficient Estimate of the proximate composition of the Bambara groundnut flour for the terms that showed significant model, insignificant lack of fit and p-value < 0.05**

Term	Coefficient	Moisture	Ash	Fat	Crude Fibre	Carbohydrate
n Intercept	$\beta_0$	6.60	2.20	5.84	0.7266	81.12
(A)	$\beta_1$	-0.6137	-0.4688	2.42	0.1929	1.13
(B)	$\beta_2$	0.3082	-	-0.0895	-0.3225	-
(AB)	$\beta_{12}$	0.4197	-	-0.9471	0.1329	-
(A <sup>2</sup> )	$\beta_{11}^2$	-	0.4063	0.9213	0.0137	-4.25
(B <sup>2</sup> )	$\beta_{22}^2$	-	-	-0.0421	0.2055	-
R <sup>2</sup> adj.		0.5728	0.6648	0.7857	0.6382	0.4710
% CV		7.08	12.31	16.14	23.72	2.84

#### Moisture content

The regression model for the moisture content of the flour samples is presented in Eq.2 and figure 1 as the contour plot. It was observed from equation 1 and figure 1 that reducing the soaking time, while increasing the boiling time and their interaction, increased the moisture content of the flour. This is because soaking time exhibited positive coefficient, while boiling time and interaction of soaking time and boiling time had positive coefficients. The positive coefficients signify direct relationship between the soaking and boiling times and their interactions with the moisture content of the flour, while the negative coefficients indicate an inverse relationship between the process variables and the moisture content of the flour samples. R<sup>2</sup>adj was 57.28% which is high, showing that the 57.28% change in the moisture content was caused by soaking and boiling times and their interactions.

$$\text{Moisture} = 6.60 - 0.6137A + 0.3082B + 0.4197AB \quad (2)$$




**Fig.1: Effect of soaking time and boiling time on the moisture content of the Bambara flour samples**

#### Ash content

The regression model for the ash content of the flour samples is presented in Eq.3, while figure 2 shows the effect of soaking time and boiling time on the ash content of flour samples. Increasing the soaking time reduces the ash contentment of the flour since soaking time has negative coefficient. The square of soaking time however had a slight increase in the ash contentment of the flour shown in the 3D surface plot (figure 2) and as also shown in equation 3.  $R^2_{adj}$  was 66.48% which is high, and the coefficient of variation (% CV) wasn't high (12.31). This shows that the model explains 66.48% of the effect of the soaking and boiling times on the ash content of the flour samples, while the difference in the percentage (33.52%) could come from extraneous variables not factored into the design

$$\text{Ash} = 2.20 - 0.4688A + 0.4063A^2 \quad (2)$$




**Figure 2: Effect of soaking time and boiling time on the ash content of soaked and boiled BGN flour**

#### Fat content

The regression model of the fat content of the samples is presented in Eq. 4 and figure 3.

$$\text{Fat} = 5.84 + 2.42A - 0.0895B - 0.9471AB + 0.9213A^2 - 0.0421B^2 \quad (4)$$

The model was significant with a satisfactory adjusted coefficient of determination,  $R^2_{adj}$  of 0.7857 (78.57%) as shown in Table 2. The high coefficient of determination showed excellent correlation between the independent variables (soaking time and boiling time) and the fat content. This is an indication that the response (fat) model is adequate, and can explain 78.57% of the total variability in the fat content of the Bambara ground nut flour samples.




**Fig.3. Contour plot for the effect of soaking time and boiling time on the fat content of BGN flour**

**Crude Fibre Content**

The regression model of the crude fibre content of the flour samples is shown Eq.5 and fig.4

$$\text{Crude fibre} = 0.7266 + 0.1929A + 0.3225B - 0.1329AB - 0.0137A^2 + 0.2055B^2 \quad (5)$$

Increasing both soaking time and boiling time increases the crude fibre content of the bambara ground nut flour samples. However, the interaction of the two independent variables and the square of soaking time reduced the crude fibre content as evidenced in equation 5 and fig.4.




**Fig. 4: Contour plot for the effect of soaking time (A) and boiling time (B) on the crude fibre content of soaked and boiled Bambara ground nut flour samples**

**Carbohydrate Content**

The regression model for the carbohydrate content is shown in Eq. 6 and figure 5 respectively

$$\text{Carbohydrate} = 81.12 + 1.13A - 4.25A^2 \quad (6)$$

The carbohydrate exhibited significant model. However, the adjusted coefficient of determination ( $R^2_{adj.}$ ) was low (0.4710) suggesting that only 47 % of the changes in carbohydrate were caused by the process variables and their interaction in the experiment. From Eq. 5, increasing the soaking time increased the carbohydrate content, while the square term reduced the carbohydrate also evidenced in figure 5 which shows the 3D surface plot of the carbohydrate content of soaked-boiled Bambara ground nut flour samples



**Figure 5: 3D surface plot of the carbohydrate content of soaked-boiled Bambara ground nut flour**

**Numeric optimization for the proximate composition**

Table 5 shows the generated Numerical Optimization solution of the proximate composition of soaked-boiled Bambara ground nut flour samples. The main criteria for constraints optimization of process parameters for the proximate composition were minimum possible soaking and minimum boiling time which generated the solution in Table 5 with desirability of 61.6%. The percentage desirability is high and acceptable. However, desirability of 100% is the most ideal if it could be obtained. It shows that if the selected critical values of 12 h of soaking and 46.26 min boiling are employed in the production of Bambara ground nut flour, that the flour would exhibit proximate composition of 7.15, 3.07, 4.79, 0.83 and 0.62% for moisture, ash, fat and carbohydrate respectively.

**Table 5: Numeric optimization Solutions of the proximate composition of the BGN flour**

Soaking time (h)	Boiling time (min)	Moisture	Ash	Fat	Crude fibre	Carbohydrate	Desirability	
12.000	46.261	7.150	3.072	4.794	0.827	75.738	0.616	Selected

The mean of the triplicate result of the functional properties of the processed (soaked and boiled) BGN flours shown in table 6 reveals that there was significant ( $P = .05$ ) differences in the values obtained. The water absorption capacity (WAC) steadily increased (in a general term) from 0.665ml/g, the least value, to 2.70ml/g due to 30 h soaking and, 60 min boiling of sample H, and the 48 h soaking and, the 30min boiling of sample J. 12 h soaking yielded 38.46 % increase in WAC and 508.88 % relatively high increase due to 30 h soaking, while 48 h soaking yielded 130.77 % increase. A combination of soaking and boiling generally enhanced the WAC.

**Table 6: Functional Properties of Processed BGN (Soaked and Boiled) flour samples**

Run	Factors		RESPONSES					Swelling C.	
	X <sub>1</sub>	X <sub>2</sub>	WAC ml/g	OAC ml/g	Foam C	Foam Stab	Gel. Temp <sup>o</sup> C	ml/g	Solubility
1	12(-1.00)	30(-1.00)	0.90±0.01 <sup>g</sup>	0.79±0.09 <sup>abc</sup>	24.31±1.18 <sup>a</sup>	97.20±1.04 <sup>a</sup>	61.00±1.00 <sup>f</sup>	60.82±0.06 <sup>n</sup>	30.10±0.91 <sup>a</sup>
2	12(-1.00)	45 (0.00)	1.75±0.05 <sup>e</sup>	0.88±0.00 <sup>a</sup>	2.00±0.04 <sup>d</sup>	74.75±0.25 <sup>bc</sup>	64.00±0.00 <sup>e</sup>	62.87±1.84 <sup>n</sup>	19.12±2.33 <sup>b</sup>

3	12(-1.00)	45 (0.00)	2.05±0.05 <sup>c</sup>	0.75±0.04 <sup>def</sup>	2.00±0.00 <sup>d</sup>	50.00±0.00 <sup>de</sup>	68.50±0.50 <sup>d</sup>	63.36±0.70 <sup>n</sup>	14.85±0.98 <sup>a</sup>
4	12(-1.00)	60(1.00)	2.00±0.00 <sup>d</sup>	0.62±0.00 <sup>g</sup>	2.00±0.00 <sup>d</sup>	86.33±1.25 <sup>ab</sup>	64.00±1.00 <sup>e</sup>	71.47±0.20 <sup>g</sup>	19.23±0.49 <sup>b</sup>
5	30 (0.00)	30(-1.00)	2.15±0.15 <sup>cd</sup>	0.70±0.00 <sup>efg</sup>	0.49±0.00 <sup>e</sup>	50.00±0.00 <sup>d</sup>	67.50±0.50 <sup>d</sup>	73.76±0.87 <sup>ef</sup>	14.44±0.60 <sup>d</sup>
6	30 (0.00)	45 (0.00)	2.35±0.05 <sup>b</sup>	0.79±0.09 <sup>abc</sup>	0.98±0.00 <sup>e</sup>	50.00±0.00 <sup>d</sup>	81.00±1.00 <sup>a</sup>	72.39±0.63 <sup>fg</sup>	15.87±0.47 <sup>c</sup>
7	30 (0.00)	45 (0.00)	2.30±0.00 <sup>b</sup>	0.66±0.04 <sup>g</sup>	0.86±0.12 <sup>e</sup>	68.33±21.67 <sup>c</sup>	78.00±0.00 <sup>c</sup>	74.73±1.79 <sup>e</sup>	14.64±0.34 <sup>d</sup>
8	30 (0.00)	60 (1.00)	2.70±0.10 <sup>a</sup>	0.88±0.00 <sup>a</sup>	0.73±0.24 <sup>e</sup>	65.00±5.00 <sup>cd</sup>	77.00±1.00 <sup>c</sup>	93.26±0.74 <sup>a</sup>	15.98±1.74 <sup>c</sup>
9	48 (1.00)	30(-1.00)	1.50±0.00 <sup>f</sup>	0.63±0.09 <sup>g</sup>	5.39±0.49 <sup>c</sup>	38.33±21.6 <sup>efg</sup>	57.00±1.00 <sup>g</sup>	86.11±0.67 <sup>b</sup>	20.52±1.22 <sup>b</sup>
10	48 (1.00)	30(-1.00)	2.70±0.10 <sup>a</sup>	0.70±0.00 <sup>efg</sup>	0.42±0.07 <sup>e</sup>	47.33±2.51 <sup>e</sup>	79.67±0.76 <sup>b</sup>	73.28±0.05 <sup>efg</sup>	15.70±0.93 <sup>c</sup>
11	48 (1.00)	45 (0.00)	2.70±0.10 <sup>a</sup>	0.84±0.04 <sup>ab</sup>	0.98±0.00 <sup>b</sup>	30.84±4.16 <sup>g</sup>	77.50±1.50 <sup>e</sup>	78.79±0.10 <sup>de</sup>	11.57±0.29 <sup>d</sup>
12	48 (1.00)	60 (1.00)	1.50±0.0 <sup>f</sup>	0.79±0.00 <sup>abc</sup>	13.66±0.97 <sup>b</sup>	96.41±0.25 <sup>a</sup>	62.50±0.50 <sup>e</sup>	75.10±2.11 <sup>de</sup>	13.86±1.85 <sup>d</sup>

**Values are mean ± standard deviation of replicated determinations (n=21). Mean values in the same column followed by different letters are significantly (P =.05) different. X<sub>1</sub> = Soaking time; X<sub>2</sub> = Boiling time,**

Table 7 shows the Summary of ANOVA and coefficient estimate of the functional properties of the Bambara groundnut flour for the terms that showed significant model, insignificant lack of fit and p-value < 0.05. The ideal regression equation showing the response variables (functional properties) as a function of the independent (process variables) can be represented in Eq.7

$$Y = \beta_0 + \beta_1A + \beta_2B + \beta_{12}AB + \beta_{11}A^2 + \beta_{22}B^2 + \varepsilon \quad (7)$$

Where: Y= response variable,  $\beta_0$ =intercept, A= soaking time, B = boiling time,  $\beta_1, \beta_2, \beta_{12}, \beta_{11}, \beta_{22}$  = coefficients of the linear, interaction and square of A and B respectively,  $\varepsilon$  = estimated error.  $\varepsilon$  is often neglected and not factored into the equation

**Table 7: Summary of ANOVA and coefficient estimate of the functional properties of the Bambara groundnut flour for the terms that showed significant model, insignificant lack of fit and p-value < 0.05**

Term	Coefficient	Foam capacity	Gelatinization Temp.	Solubility
n intercept	$\beta_0$	-2.74	81.30	17.56
A	$\beta_1$	-2.13	2.26	-2.71
B	$\beta_2$	-2.35	3.12	-3.23
AB	$\beta_{12}$	7.15	1.05	-
A <sup>2</sup>	$\beta_{11}$	5.58	-9.10	-
B <sup>2</sup>	$\beta_{22}$	6.76	-10.34	-
R <sup>2</sup> .adj		0.6806	0.6463	0.3909
% CV		91.56	7.13	21.83

**Foam capacity**

The regression model for foam capacity is shown in Eq. 8 and the 3D surface and contour plots shown in figures 6a and 6b respectively

$$\text{Foam capacity (\%)} = -2.74 - 2.13A - 2.35B + 7.15AB + 5.58A^2 + 6.76B^2 \quad (8)$$




Fig. 6a: 3D Surface plot for the foam capacity of soaked and boiled BGN flour,





Fig 6b: Contour plot for the effect of soaking time (h) and boiling time (min) on the foam capacity of soaked and boiled BGN flour


#### Gelatinization temperature

The regression model for gelation temperature is shown in Eq. 9, while the Surface and Contour plots are shown in figures 7a and 7b respectively

$$\text{Gelatinization temperature} = +81.30 + 2.26A + 3.12B + 1.05AB - 9.10A^2 - 10.34B^2 \quad (9)$$



**Figure 7a: 3D Surface plot for the gelatinization temperature of soaked and boiled BGN flour**




**Figure 7b: Contour plot for the effect of soaking time and boiling time on the gelatinization temperature of soaked and boiled BGN flour**


**Solubility**

The regression model for solubility of the samples is shown in Eq. 10, while the 3D surface and contour plots are shown in figures 8a and 8b respectively

$$\text{Solubility (\%)} = 17.56 - 2.71A - 3.23B \quad (10)$$



**Figure 8a: 3D Surface plot for the solubility of soaked and boiled BGN flour**



**Figure 8b: Contour plot for the effect of soaking time (A) and boiling time (B) on the solubility of soaked and boiled BGN flour**

**Numeric optimization for the functional properties**

The numeric optimization Solutions for the functional properties of the flour is shown in Table 8. The main criteria for constraints optimization of process parameters for the functional properties were minimum possible soaking and minimum boiling time which generated the solution in Table 8 with desirability of 60.30 %. This value is high and acceptable. However, desirability of 100 % is the most ideal if it could be obtained (Ishiwuet *al.*, 2014). It shows that if the selected critical values of 12 h of soaking and 9.97 min boiling are employed in the production of Bambara ground nut flour, that the flour would exhibit functional properties of 14.33, 63.95°C and 22.42% for foam capacity, gelation temperature and solubility respectively.

**Table 9: Numeric optimization Solutions for the functional properties of the flour**

Soaking	Boiling time	Foam	Gelatinization	Solubility	Desirability
---------	--------------	------	----------------	------------	--------------

time (h)	(min)	Capacity	Temp.			
12.000	9.966	14.327	63.945	22.423	0.603	Selected

### Graphical optimization for the functional properties

The solution displayed in the numerical optimization could be graphically shown as figure 10




Fig. 10: Optimization plot of the functional properties of Bambara ground nut flour

## 4. CONCLUSION

From this study, it may be concluded that the combination of soaking and boiling/water blanching had significant effect on nutrient losses however; longer soaking times yielded the most fat. Minimum soaking time yielded better functional properties. Models for the prediction of optimal nutrient responses and optimal functional properties were developed.

## REFERENCES

- Abdulsalami, M.S. and Sheriff H.B. Effect of processing on the proximate composition and mineral content of bambara groundnut (*Voandzeia subterranean*). *Bayero Journal of Pure and Applied Sciences*, 2010 3(1): 188 - 190.
- Annan, N.T.; Plahar, W.A.; Nti, C.A. Dissemination of improved bambara groundnut processing technologies through a new coalition arrangement to enhance rural livelihoods in Northern Ghana, in Training of Trainees workshop. Tamale Ghana. In Mubaiwaet *al.*, 2003. Doi:10.1080/87559129.2016.1149864,
- AOAC Association of official Analytical Chemists. Official Methods of Analysis of the AOAC. Washington D. C. USA. 2005.
- Arise, A. K., Aliyu, B. N., Ajidagba, S. D. Effect of thermal processing and fermentation on the chemical composition, protein digestibility and functional properties of bambara protein isolates. *Carpathian Journal Food Science Technology*, 2020; **12**, 148-156
- Bamshaiye, O. M., Adegbola, J. A., Bamishaiye, E. I. Bambara groundnut: An under-utilized nut in Africa. *Advances in Agricultural Biotechnology* 2011; **1**, 60–72.
- Barimalaa, I. S., Brown, T., Oboh, C.A. and Jaja, E. T. Effects of soaking conditions on certain functional properties of bambara groundnut flour important in Okpa preparation in Niger Delta *Bioglobia*: 2000<sup>b</sup>; **3** (1), 53-59.
- Erbay, Z. and Icier, F. A Review of Thin Layer Drying of Foods: Theory, Modeling and Experimental Results. *Critical Reviews in Food Science and Nutrition*, 2010; **50**, 441-464. <https://doi.org/10.1080/10408390802437063>
- Ishiwu, C. N, Nkwo, V. O, lwouno J. O, Obiegbuna, J. E and Uchegbu, N. N. Optimization oftasteand

textureofbiscuit produced from blend of plantain,sweetpotato and malted sorghum flour,*AfricanJournal of Food Science*,2014; Vol.8 (5), 233-238.Doi:10. 5897/AJFS2013.1112 <http://www.academicjournals.org/AJFS>

Karuppaiya, V., Sasikumar, M. E., and Vijayagopa, V. I. Optimization of process variables using Response Surface Methodology (RSM) for ethanol production from cashew apple juice by *Saccharomyces cerevisiae**Asian Journal of Food and Agro-Industry* 2010, 3 (4), 462-473

Mazahib, A.M.; Nuha, M.O.; S., S.I.; Babiker, E.E. Some nutritional attributes of bambara groundnut as influenced by domestic processing. *International Food Research Journey*, 2013; 20 (3), 1165-1171.

Mkandawire, C.H. Review of Bambara Groundnut (*Vigna subterranea* (L.)Verdc.). Prod. In*Sub-Sahara African Agricultural Journal* 2007; 2(4), 464–470.

Murevanhema, Y.Y and Jideani, V.A. Potential of Bambara Groundnut (*Vigna subterranean* (L.)Verdc) Milk as a Probiotic Beverage.A Review. *Critical Rev. Food Science Nutrition*, 2013; 53, 954–967

Myers, R. H, Montgomery, D. C and Anderson-Cook, C. M. Response Surface Methodology (RSM), Process and Product Optimization using Designed Experiments 3<sup>rd</sup> Edition.John Wiley & Sons. 2009; Pp. 33-190

Nomathemba, G. M., Abe S. G., and Hussein S. Bambara Groundnut (*Vigna subterranea* [L.] Verdc.)Production, Utilization and Genetic Improvement in *Sub-Saharan Africa. Review Agronomy*, 2021; 11, 1345. <https://doi.org/10.3390/agronomy11071345>

Nti, C.A. Effects of bambara groundnut (*Vigna subterranea*) variety and processing on the quality and consumer appeal for its products. *International Journal Food Science Technology*, 2009; 44, 2234–2242.

Onwuka, G. I. Food analysis and instrumentation. Theory and practices, Naphtali Prints, Lagos, Nigeria. 2005

Yao D.N, Kouassi, K.N, Erba D, Scazzina F, Pellegrini N, Casiraghi M.C. Nutritive evaluation of the Bambara groundnut Ci12 landrace [*Vigna subterranean* (L.) Verdc (*Fabaceae*)] produced in Cote d'Ivoire. *International Journal of Molecular Sciences*. 2015; 16:21428-21441; doi:10.3390/ijms160921428

## **DEFINITIONS, ACRONYMS, ABBREVIATIONS**

Here is the Definitions section. This is an optional section.

**Term:** Definition for the term

## **APPENDIX**