

Original Research Article

An Analysis of farmers extent of Adoption and Knowledge of Bt. Cotton in the Jangaon District of **Telangana**

Comment [CC1]: Please cheep the name "Bt. cotton" like in manuscriptum text. Please, change an analysis because is not only one analysis. Please change the title of paper!

Abstract –The study was conducted in Jangaon district of Telangana during the year 2022. The main purpose of this study is to measure the Extent of Adoption and knowledge of Bt cotton in Jangaon District of Telangana. A total 120 respondents were selected purposively from six villages under Lingala ghanpur block because most of the respondents were cotton growers. Descriptive research design was adopted for the study and data was collected by personal interview method by using pre structured interview schedule. This study revealed that the respondents (49.17%) had medium level of adoption followed by low (31.66%) and high (19.16%) levels of adoption among the respondents and also the most of the respondents (38.33%) had high level of knowledge on Bt. Cotton Adoption followed by medium (35%) and low (26.66%) levels of knowledge.

Keywords – Adoption, Knowledge, Bt. Cotton

INTRODUCTION

Cotton is one of the most important fiber and cash crop of India and plays a dominant role in the industrial and agricultural economy of the country. Cotton is the most important fiber crop not only of India but of the entire world. It provides the basic raw material (cotton fiber) to cotton textile industry, Cotton in India provides direct livelihood to 6 million farmers and about 40-50 million people are employed in cotton trade and its processing. India has the largest area under cotton cultivation in the world though it is the world's third largest producer of cotton after China and the USA. Currently it is grown over 6 per cent of the net sown area.

In Telangana, cotton scenario before introduction of Bt. cotton was not a bright picture. The crisis in cotton revolves around the main issues like raising cost of cultivation, hiked use of pesticides, consequent endemic pest and diseases and

market fluctuations which have hit the producers so severely that most of them end up deep in debt. Epidemics of whitefly and boll worms, besides frequent and severe droughts had forced several farmers to commit suicide. Under these circumstances, Bt. cotton has emerged as an attractive option before the cotton farmers (Singh and Kaushik, 2007).

Cotton is one of the vital commercial fiber crops, play a prominent role in the national and international economy due to its high commercial value, it is also popularly known as 'White Gold'. In India, it is important cash and commercial crop valued for its fiber and vegetable oil. It is a source for earning the valuable foreign exchange by providing employment to millions of people and hence plays a significant role in the national economy. The diverse products obtained from cotton include textile raw material, cotton seed is a major source of vegetable oil and cotton cake as a rich source of high quality protein for livestock feed. Cotton is primarily grown as a fiber crop. It is harvested as 'seed cotton, which is then 'ginned to separate the seed and lint. The long lint fibers are further processed by spinning to produce yarn that is knitted or woven into fabrics. (Kalidasan 2020).

Comment [CC2]: Some notions from the first paragraph are repeated. Please keep the notions uniform.

Research methodology –

Descriptive research design was adopted for the study as it describes the characteristics or phenomena that are being studied. The present study was conducted in Jangaon district Telangana. Out of 6 blocks in Jangaon district, Lingala ghanpur block is selected purposively based on maximum area covered under cotton cultivation. From the selected block, six villages were selected purposively based on the maximum area covered under Bt cotton cultivation.

Comment [CC3]: How many persons are respondents? What were the criteria for choosing them? What methods were used in this research?

RESULTS AND DISSCUSSION

1. Adoption of Bt. Cotton by the farmers –

Table no. 1 distribution of respondents according to their adoption level on Bt. Cotton cultivation.

S.No	Adoption of Bt. Cotton cultivation practices	FA (FULLY ADOPTED)		PA (PARTIALLY ADOPTED)		NA (NOT ADOPTION)	
		<i>F</i>	%	<i>f</i>	P%	<i>f</i>	%
1.	Summer ploughing	25	20.83	71	59.16	24	20.83
2.	FYM	-	-	81	67.5	39	32.5
3.	Recommended spacing	-	-	64	53.33	56	46.66
4.	Recommended seedrate	83	69.16	37	30.83	-	-
5.	Recommended dose of N.P.K	12	10	91	75.83	17	14.16
6.	Inter cultivation	14	11.66	92	76.66	14	11.66
7.	Pre emergence herbicides	30	25	71	59.16	19	15.83
8.	Inter cropping	22	18.33	82	68.33	16	13.33
9.	Hand weeding	-	-	101	84.16	19	15.83
10.	Post emergence herbicide	-	-	-	-	120	100
11.	Mechanical weeding	36	30	65	54.16	19	15.83
12.	N&K should applied in three equal Splits	73	60.83	42	35	5	4.16
13.	Earthing up	-	-	56	48.33	62	51.66
14.	Plant protection measures	31	25.83	65	54.16	24	20

15.	IPM	40	33.33	66	55	14	11.66
16	Trap crops	32	26	54	45	34	28.33
17	Stem applications	37	30.83	63	52.5	20	16.66
18	Monitoring pests by using sticky traps, pheromones and light traps	-	-	62	23.33	58	76.66

From the above table 1 it can be observed that among the (69.16%) of the respondents are completely Adopted the recommended seed rate, (60.83%) of the respondents are Adopted the fertilizers N & K should applied in three equal splits, (33.33%) of the respondents are Adopted IPM practices, (30.83%) of the respondents are Adopted stem applications, (26%) of the respondents are Adopted trap crops, (84.16%) of the respondents are partially Adopted hand weeding, (76.66%) of the respondents are partially Adopted inter cultivation method, (75.83%) of the respondents are partially Adopted recommended dose of N.P.K, (68.33%) of the respondents are partially Adopted inter cropping, (67.5%) of the respondents are partially Adopted FYM, (76.66%) of the respondents are not Adopted Monitoring pests by using sticky traps, pheromones and light traps, (51.66%) of the respondents are not Adopted earthing up, (46.66%) of the respondents are not Adopted recommended spacing , (32.5%) of the respondents are not Adopted FYM, (28.33%) of the respondents are not Adopted trap crops.

Table:2 Distribution of respondents according to their overall Adoption of Btcotton (n=120)

S.No	Adoption	Frequency	Percentage
1.	Low(1-7)	38	31.66
2.	Medium(7-14)	59	49.17
3.	High(above14)	23	19.17
	Total	120	100.00

Comment [CC4]: I believe that 120 respondents are too few to draw the necessary conclusions for the research.

An examination of Table 2 and Fig 1 indicate that the most of the respondents (49.17%) had medium level of adoption followed by low(31.66%) and high (19.16%) levels of adoption among the respondents.**Joshi (2014)**

Fig 1 distribution of respondents according to their overall adoption of Bt. Cotton Cultivation.

2. Knowledge of farmers on Bt. Cotton cultivation .

Table 3 distribution of respondents according to their knowledge level on Bt. Cotton cultivation

S.No	Particulars	A AGREED		UD (UNDECI DED)		DA DISAGREED	
		<i>f</i>	%	<i>f</i>	%	<i>f</i>	%
1.	Ploughing	18	15	42	35	60	50
2.	Time of sowing	76	63.33	32	26.66	12	10
3.	Weed control methods	34	28.33	31	25.83	55	45.83
4.	Pre-emergence herbicides	35	23.33	61	50.83	24	20
5.	Pests and diseases of Bt cotton	26	21.66	59	49.16	35	29.16
6.	Harvesting stage	89	74.16	13	10.83	18	15
7.	Yield	42	35	53	44.16	25	20.83
8.	Marketing	48	40	45	37.5	27	22.5

From the above table 3 it can be observed that among the(74.16%) of the respondents are completely agreed that harvesting stage on time, (63.33%) of the respondents are agreed that on time sowing of crop,(40%) of the


respondents are agreed that on time marketing ,(35%) of the respondents are agreed that getting expected yield,(28.33%) of the respondents are agreed that they are effective controlling weeds,(50.83%) of the respondents are undecided that they are unaware to spray pre emergence herbicides, ,(49.16 %) of the respondents are undecided that they are unable to identity pest and diseases,(44.16%) of the respondents are undecided that they are ,(37.5%) of the respondents are undecided that they having best marketing facilities, ,(35%) of the respondents are undecided regarding ploughing, ,(45.83%) of the respondents are disagree to weed control,(29.16%) of the respondents are disagree regarding pest and disease of Bt cotton, ,(20.83%) of the respondents are disagree regarding cotton yield.

TABLE4 Distribution of respondents based on overall knowledge of Bt.cotton (n=120).

S.No	Knowledge	Frequency	Percentage
1.	Low(6-11)	32	26.66
2.	Medium(12-16)	42	35.00
3.	High(above16)	46	38.33
	Total	120	100.00

An examination of Table 4 and Fig 2 indicate that the most of the respondents (38.33%) had high level of knowledge on Bt. Cotton Adoption followed by medium (35%) and low (26.66%) levels of knowledge. **Jeewanet al.(2020)**

FIG 2 Distribution of respondents based on overall knowledge of Bt.cottoncultivation



CONCLUSION

Comment [CC5]: The authors have more conclusions not only one conclusion.

It is concluded that most of the respondents had medium level of knowledge and adoption towards improved cultivation practices of cotton. And also concluded that medium level of knowledge and adoption of farmers towards improved cultivation practices of cotton. Most of the respondents are having medium level knowledge on time of sowing of crop, harvesting stage and adopting new cultivation practices, recommended seed rate and also fertilizer application rate and partially adopted FYM and not adopting the Integrated Pest Management practices. The respondents expressed their problems such as high cost and non-availability of required hybrid seeds at right time, high cost and non-availability of chemical fertilizers in time, More sucking pest attack, etc., and their suggestions are to provide more practical exposure. Hence the government should provide good provision of improved seed in time with minimum cost, training on identification of pests and diseases and their control

References :

- Balakrishnan, T. and Vasanthkumar, J. (2017).** Knowledge level of System of Rice Intensification (SRI) technology among farmers in Cuddalore district of Tamil Nadu. *International J. of Current Research*. **9**: 065-068.
- Bondarwad, S.P.; Wangikar, S. D. and Desmukh, N. D. (2018).** Correlation of adoption of Bt. cotton production technology. *Agriculture Update*. **300-302**.
- Dharmendra Yadav.; Patel, M. M. and Choudhary, R. K. (2016).** IPM in cotton adoption and constraints. *J. of cotton research and development*. **21**: 230-234.
- Gangadhar, J. (2019).** Marketing behavior of cotton farmers in Warangal district of Andhra Pradesh. *M.Sc. (Ag.) Thesis*. Acharya N G Ranga Agricultural University, Hyderabad.
- Jeewan, R.S.; Sangram, S.; Hanuman, L. and Choudhary, L.R. (2020).** Knowledge level of farmers about improved tomato production technology. *Rajasthan J. of Ext. Edu.* **19**: 139-143
- Joshi, P. L. (2014).** Extent of knowledge and adoption of cotton growers about modern practices of cotton in Bhalarea. *M.Sc. (Ag)*. GAU, Anand Campus, Anand.
- Maraddi, G. N. V. and Sathish, H. S. (2021).** Analysis of farmer's knowledge and adoption of improved cotton cultivation practices. *Agriculture Update*; **9**(1): 1-6.
- Prasanth. (2021).** A study on adoption of organic farming in cotton in Karimnagar district of Andhra Pradesh. *M. Sc. (Ag) thesis*

submitted to Acharya N G Ranga Agricultural University,
Hyderabad.

Rai, D.P. and Bhupendra, S. (2020). Extent of knowledge and constraints in cotton production technology in Madhya Pradesh *.Indian Research J.ofExt.Edu.***10(2):** 78-80.

Singh and Kaushik,(2007) Knowledge of Cotton Growers in the Recommended Package of Practices of Cotton Cultivation.*Indian J. Ext.Edu .45:*7-10s

Trivedi (2020).Success stories of cotton farmers to study the adoption behaviour on Integrated Pest Management of cotton in Nalgonda District. *Agricultural Ext. Review.***17(5):** 22-25