

PATTERN OF CD4 COUNT AND HEMATOLOGICAL INDICES IN HIV
SERODISCORDANT PARTNERS IN JOS, PLATEAU STATE, NIGERIA.

ABSTRACT

This was a cross sectional study designed to evaluate the CD4 count and hematological indices in HIV serodiscordant partners in Jos, Nigeria. A total of 20 discordant HIV couples (40 patients) and 20 controls (40 non HIV couples) aged between 18 and 49 years were included in the study. Each participant provided a 5ml venous blood sample that was collected into EDTA containers for the analysis of the CD4 count and hematological indices. The following blood parameters were measured using a three pack full blood count autoanalyzer: white blood cell (WBC), red blood cell (RBC), hematocrit/packed cell volume (HCT/PCV), hemoglobin (HB), platelet count (PLT), lymphocyte, neutrophil, and mixed cell (NUE/NAC) count. The CD4+-T cell was measured using flow cytometry. The results showed that the mean RBC count, platelet count, mixed cell count (Nue/Nac), HCT/PCV and HB levels, as well as the CD4-T cell count, were all significantly lower while the mean age was higher in the HIV discordant test group compared to control group ($p < 0.05$) respectively. The mean WBC, lymphocyte, and neutrophil counts in the HIV discordant test group did not differ significantly from the control group ($p > 0.05$). The female HIV discordant test group had significantly lower HB, HCT/PCV, RBC, lymphocyte and CD4 count than in the female control group ($p = 0.008$; 0.002 ; 0.000 , 0.008 , 0.000) respectively. The male HIV discordant test group had significantly lower mean neutrophil and CD4 counts than in the male control ($p = 0.000$; 0.012) respectively. The female HIV seropositives had a significantly lower mean RBC count ($p = 0.000$), Hb ($p = 0.037$), PCV/HCT ($p = 0.005$) and CD4 count ($p = 0.000$) than in female control while the female HIV exposed seronegatives had significantly lower mean CD4 count ($p = 0.000$) and Hb ($p = 0.037$) levels than in female control while the male HIV seropositives had significantly lower CD4 count compared to male control ($p = 0.000$). This study has revealed significant changes in CD4-T cell count and hematological indices in HIV discordant couples, which calls for an urgent interventional strategy to prevent the potential anemia, leucocytopenia, and weakened immunity that may result in both HIV seropositives and seronegative exposed couples.

Key words: HIV, Discordant couple/partner, CD4-T cell, Hematological indices, anemia, leucocytopenia, age.

INTRODUCTION

Human immunodeficiency virus (HIV), a member of the lentivirus family, causes a range of symptoms anchored on the reduced immune function of its host by causing devastating effects on the host's innate immune capabilities, allowing HIV replication in the host cell, leading to HIV infecting new immune cells, and resulting in the host's susceptibility to infections and if unchecked, this leads to Acquired Immunodeficiency Syndrome (AIDS) (Aaron *et al.*, 2017; UNAIDS, 2019). HIV is a public health issue affecting the world's population; 79.3 million people have been infected with the virus since the epidemic began, 36.3 million have died from AIDS, and recently 37.7 million people were infected with HIV in 2020 (UNAIDS, 2022). It is the main cause of morbidity and mortality in Sub-Saharan Africa (SSA), accounting for 71% of the global population of people living with HIV (James *et al.*, 2018; Roth *et al.*, 2018). In Nigeria, the first HIV/AIDS patient was identified and reported in Lagos in 1985 (Awofala and Ogundele, 2017). Nigeria, the most populous country in Africa (UNFPA, 2020), has 1.9 million people living with HIV (with prevalence of 1.4%) between 15–49 years, making it the third highest in HIV load (UNAIDS, 2019).

HIV discordance is a situation that occurs when one partner is HIV positive and the other is HIV negative (Mehra *et al.*, 2015). HIV discordant couples are a high-risk group for HIV transmission (Ravikumar and Balakrishna, 2013). In Sub-Saharan Africa, 50% of PLWH are in sero-discordant partnerships (Chemaitelly *et al.*, 2012; Curran *et al.*, 2012). Sero-discordant married or cohabiting couples account for a large proportion of new HIV infections in Sub-Saharan Africa, and transmission within this crucial population is a preventable driver of the epidemic (Irunguet *et al.*, 2016; Patel *et al.*, 2018). Thus, HIV preventive and treatment efforts focus on sero-discordant couples.

HIV causes AIDS, a systemic disease marked by impaired cellular and humoral immune responses (Balasubramaniam *et al.*, 2019; van Woudenberg *et al.*, 2020) which has been linked with hematological abnormalities. Hematological abnormalities are linked to disease progression and death in HIV patients (Damtie *et al.*, 2021). Anemia, leucopenia, neutropenia, lymphopenia, and thrombocytopenia have been observed in HIV-infected people before or after ART initiation (Bhardwaj *et al.*, 2020; Damtie *et al.*, 2021), indicating that hematological abnormalities in HIV individuals are induced by either the virus or ART. More so, anemia, lymphocyte count, and thrombocytopenia are associated with CD4 levels (Bhardwaj *et al.*, 2020). In addition, hematopoietic progenitor cells express CD4 receptors, type 4 C-X-C chemokine receptors and type 5 chemokine receptors, making them susceptible to being infected by HIV (Tsukamoto, 2020). In light of the above, it is important to evaluate the CD4 Count and hematological indices in HIV discordant partners in Jos, Nigeria. Hematological indices offer physiological insights on the reticuloendothelial system and the blood picture (Okoroiwue *et al.*, 2017), and this has not been investigated in this group before.

MATERIALS AND METHODS

Study Area and Location

The study area for this work was Jos North Local Government Area of Plateau State and location includes APIN (Aids Preventive Initiative of Nigeria) section of Our Lady of Apostles (OLA) Hospital, Faith Alive Foundation Hospital and Plateau State Specialist Hospital where HIV screenings are carried out.

Study Design and Subject Selection

For this investigation, a cross sectional study approach was used. The participants were partners who were known to be HIV positive and exposed seronegatives and are between the ages of 18 and 49 years old. Additionally, HIV-negative couples within the aforementioned age range who appeared to be in good health were employed as controls. The HIV-positive people were already taking medication, but their negative counterparts in a discordant relationship were not on ART.

Study Population

The study population included male and female subjects in discordance relationship within the age of 18 to 49 years attending the APIN section of Our Lady of Apostles Hospital, Faith Alive Foundation and Plateau State Specialist Hospital. A total of 20 discordant HIV couples (40 patients) and 20 controls (40 non HIV couples) were included in the study.

Sample Collection

Each participant gave venous blood sample of 5 ml, which was then drawn into EDTA containers after the collection site had been cleaned with 70% alcohol. This was utilized for the determination of full blood count as well as CD4 count.

Inclusion criteria

Participants who tested HIV-negative and had been in a stable, discordant relationship for at least three months were included. The study comprised participants who were registered patients at the hospital under investigation, between the ages of 18 and 49, and who had the necessary status documentation. Additionally, a control group of people who were within the age bracket and appeared healthy was included in the study (non HIV subjects).

Exclusion Criteria

Participants who were already bedridden due to AIDS, those who were not registered patients or had improper documentation with the institution under research, and those who refused to give informed consent were all excluded from the study.

Ethical Approval

Ethical approval was obtained from the Ethics Committees of the hospitals: Plateau State Specialist Hospital (PSSH/ADM/ETH.CO/2019/005); Faith Alive Foundation Hospital (FAFEC/08/34/25) and Our Lady of Apostles Hospital (dated 13th June, 2018) where the study was carried out.

Laboratory methods

Full blood count was determined using a three pack automated full blood count analyzer while the CD4 count was assayed using Flow cytometry.

Statistical Analysis

The data obtained were analyzed using independent t-test and one-way analysis of variance (ANOVA) with the aid of SPSS statistics tool version 23.0 software. Significant level was assumed at $p < 0.05$.

RESULTS

When compared to the control group, the HIV discordant test group's mean age was significantly higher ($p=0.000$). However, when compared to the control group, the mean RBC count, platelet

count, mixed cell count (Nue/Nac), HCT/PCV and HB levels, as well as the CD4-T cell count, were all significantly lower in the HIV discordant test group ($p < 0.05$). The mean WBC, lymphocyte, and neutrophil counts, however, did not differ significantly from the control group ($p > 0.05$ in each case). See table 1.

The male control group's mean age was significantly higher than that of the female control group ($p = 0.000$). The female test group's mean age was significantly higher than that of the female control group ($p = 0.000$). Additionally, the mean age of the male test group was significantly higher than that of the female control group and the female test group who tested positive for HIV ($p = 0.000$, respectively). See table 2.

In comparison to the control groups, as well as between the male and female test groups, there was no statistically significant difference in the mean level of total WBC ($p = 0.524$). See table 2.

In comparison to the values seen in the female and male control groups, respectively, the mean level of RBC seen in the female test group was significantly lower ($p = 0.000$). Additionally, the mean RBC level was significantly lower in the male test group than in the female control group ($p = 0.000$). See table 2.

In comparison to the female (12.95 ± 0.98) and male (12.81 ± 1.22) control groups, the mean HB level was significantly lower in the female test group (11.48 ± 1.43) ($p = 0.008$). See table 2.

There was significantly lower HCT/PCV level observed in the female test group (34.90 ± 3.82) when compared to the female (41.00 ± 2.90) and male (39.80 ± 3.30) control groups ($p = 0.002$).

Also, the mean HCT/PCV level was significantly lower in the male test group than in female control group (35.98 ± 9.40 Vs 41.00 ± 2.90 ; $p = 0.002$).

There was significantly lower platelet count observed in the male test group when compared to the female control groups (220.70 ± 70.99 Vs 280.20 ± 60.41 ; $p=0.022$). See table 2.

When the test groups of males and females as well as the controls were examined, the mean NUC/NAC levels did not differ significantly ($p>0.05$). See table 2.

The mean lymphocyte counts were significantly higher in the male test group than in the female test group (52.85 ± 8.41 Vs 44.85 ± 11.16 ; $p=0.008$), but significantly lower in the female test group compared to the female (52.00 ± 1.86) and male (52.30 ± 4.23) control groups. See table 2.

In comparison to the female control group, the mean neutrophil counts in the male test group were significantly lower ($34.508.66$ Vs $40.906.84$; $p=0.012$), but it did not differ significantly from those found in the female test group ($p>0.05$). See table 2.

Compared to the values seen in the male and female control groups, the mean CD4 count was significantly lower in the female and male test groups ($p=0.000$) respectively. Also, the mean CD4 count was significantly in the male test group than in the female test group ($p=0.000$). See table 2.

When the mean ages of the study subjects were compared between the groups, there were significant differences ($f\text{-value}=15.054$; $p=0.000$). The mean ages of the individuals who were male controls, female HIV positives, and male HIV positives were significantly different from those of the participants who were female controls ($p=0.000$, respectively). Additionally, as compared to female controls and HIV positive people, the mean age of the male HIV exposed seronegative participants was significantly higher ($p=0.000$, respectively). See table 3.

When compared across the groups under study, the mean total white blood cell count (WBC), platelet count, and NUE/NAC levels did not differ significantly ($p>0.05$). See table 3.

The mean red blood cell count was significantly lower in the female HIV seropositives than in the male and female control groups ($p=0.000$, respectively) and significantly different between the groups ($F=5.709$; $p=0.000$). See table 3.

When the mean hemoglobin (HB) levels of the groups were examined, there was a significant difference ($F=2.523$; $p=0.000$). See table 3.

When participants' mean levels of haematocrit and packed cell volume were compared across the study groups, there were significant differences ($F=3.726$; $p=0.005$) as well as between the female HIV positive and male HIV exposed seronegative groups than in the female control group. See table 3.

The mean lymphocyte level varied significantly amongst the groups under investigation ($F=2.456$; $p=0.041$). See table 3.

Comparing the levels of neutrophils in the various study groups revealed additional significant differences ($F=2.423$; $p=0.043$). In addition, male HIV exposed seronegatives had significantly lower mean neutrophil levels than female HIV positives ($p=0.043$). See table 3.

Comparing the mean CD4 counts of the study groups revealed significant differences ($F=16.6672$; $p=0.000$). Compared to the values seen in the male and female control groups, the mean CD4 count was significantly lower in the female HIV seropositives, female exposed HIV seronegatives, and male HIV seropositives ($p=0.000$, respectively). Male HIV exposed seronegatives had a significantly lower mean CD4 count than female HIV positives ($p=0.000$).

See table 3.

Table 1: CD4 count and levels of Hematological indices in the HIV discordant and control groups studied (Mean±SD).

*Statistically significant at $p < 0.05$.

Table 2: Levels of CD4 count and hematological indices in the male and female HIV discordant test groups studied (mean±SD).

Parameter	Control group (n=40)	HIV Discordant Test group (n=40)	t-value	p-value
Age	35.6±6.9	40.5±6.1	3.350	0.000*
WBC	5610.00±1401.43	5220.00±1502.85	1.200	0.234
RBC	4.64±0.43	4.10±0.54	4.879	0.000*
HB	12.88±1.10	11.92±1.76	2.923	0.005*
HCT/PCV	40.40±3.13	35.41±7.10	4.041	0.000*
Platelet	277.95±56.29	241.20±75.48	2.468	0.016*
NUE/NAC(Mixed cells)	7.40±1.68	6.65±1.17	2.322	0.023*
Lymphocyte	52.15±6.17	48.85±10.56	1.706	0.092
Neutrophil	40.45±5.46	38.20±9.71	1.277	0.205
CD4	1035.20±141.11	616.48±348.28	7.047	0.000*

Comment [U1]: Standard Deviation for WBC is very high, data should be cross checked

Parameter	Female control (n=20)	Male control (n=20)	Female discordant test group (n=20)	Male discordant test group (n=20)	f-value	p-value
Age	30.5±4.5	40.7±4.6 ^a	37.5±5.6 ^a	43.5±5.2 ^{a,b}	25.128	0.000
WBC	5820.0±1821.97	5400.00±786.73	5790.00±1622.57	5250.00±1414.77	0.753	0.47
RBC	4.71±0.51	4.57±0.33	3.98±0.38 ^{a,b}	4.23±0.65 ^a	9.246	0.002
HB	12.95±0.98	12.81±1.22	11.48±1.43 ^{a,b}	12.37±1.97	4.192	0.008
HCT/PCV	41.00±2.90	39.80±3.30	34.90±3.82 ^{a,b}	35.98±9.40 ^a	5.648	0.002
Platelet	280.20±60.41	275.70±53.34	261.70±75.97	220.70±70.99 ^a	3.392	0.022
NUE/NAC (mixed cells)	7.10±1.86	7.70±1.45	6.80±1.28	6.50±1.05	2.522	0.064
Lymphocyte	52.00±1.86	52.30±4.23	44.85±11.16 ^{a,b}	52.85±8.41 ^c	4.189	0.008
Neutrophil	40.90±6.84	40.00±3.73	41.90±9.49	34.50±8.66 ^a	3.881	0.012
CD4	1042.95±109.86	1027.45±169.33	469.20±270.79 ^{a,b}	763.75±360.55 ^{a,b,c}	23.909	0.000

Comment [U2]: Standard Deviation for WBC is very high, data should be cross checked

Comment [U3]: Check Standard deviation values

Key:

a=compared with female control

b= compared with male control

c= compared with female test group

Table 3: Levels of CD4 and hematological indices in the participants studied (mean±SD).

Parameter	Female control (n=20)	male control (n=20)	Female positives (n=16)	Female negatives (n=4)	male positives (n=16)	male negatives (n=4)	f-value	p-value
Age	30.5±4.5	40.7±4.6 ^a	37.1±5.6 ^a	38.8±6.4	45.3±7.5 ^a	43.1±4.6 ^{a,c}	15.054	0.000
WBC	5820.0±1821.97	5400.00±786.73	5168±1709.47	5275.00±1431.49	5087.50±1340.09	5415.00±1457.07	0.644	0.667
RBC	4.71±0.51	4.57±0.33	3.99±0.24 ^{a,b}	3.95±0.79	4.29±0.65	4.37±0.56	5.709	0.000
HB	12.95±0.98	12.81±1.22	11.54±1.05	11.23±2.72	12.43±1.98	12.40±1.53	2.523	0.037
HCT/PCV	41.00±2.90	39.80±3.30	34.88±3.28 ^a	35.00±6.22	35.16±9.85	37.92±6.00 ^a	3.726	0.005
Platelet	280.20±60.41	275.70±53.34	263.88±78.03	253.00±77.36	225.63±67.35	259.5u8±68.70	2.101	0.075
NUE/NAC (Mixed cells)	7.10±1.86	7.70±1.45	6.94±1.29	6.25±1.26	6.50±1.03	7.03±1.48	1.631	0.163
Lymphocyte	52.00±1.86	52.30±4.23	44.69±12.31	45.50±5.45	52.94±7.61	50.50±8.75	2.456	0.041
Neutrophil	40.90±6.84	40.00±3.73	42.31±10.40	40.25±4.92	33.94±8.05	39.33±7.91 ^c	2.423	0.043
CD4	1042.95±109.86	1027.45±169.33	442.00±264.68 ^{a,b}	578.00±309.25 ^{a,b}	828.88±357.65 ^{a,b}	825.84±337.79 ^c	16.6672	0.000

Comment [U4]: Standard Deviation for WBC is very high, data should be cross checked

Comment [U5]:

Comment [U6]: Standard Deviation for CD4 is very high, data should be cross checked

Key:

a=compared with female control

b= compared with male control

c= compared with female HIV positive

DISCUSSION

Individuals with HIV-1 infection, those not receiving antiretroviral medication, and those with the disease in its advanced stages typically report having hematological abnormalities of the major blood cell lines (Duguma *et al.*, 2021). Due to its destructive effects on the immune system and various organs in the human body, which result in considerable unfavorable alterations in several hematological indices, HIV continues to be a topic of enormous health relevance and debate around the world. The present study evaluated the CD4 count and hematological indicators in HIV discordant partners in JOS, Nigeria.

In the current investigation, the mean age was significantly higher in the HIV discordant test group (40.5 ± 6.1) than in the control group (35.6 ± 6.9). Given that the mean age of HIV-positive people indicates that they are among the country's working population, proper and prompt therapeutic management of these people will further increase their productivity throughout the economic cycle.

The mean RBC count, platelet count, mixed cell count (Nue/Nac), HCT/PCV and HB levels, as well as the CD4-T cell count, were all significantly lower in the HIV discordant test group when compared to the control group. The lower levels of RBC count, HCT/PCV and HB recorded in this study shows that the HIV serodiscordant couples are at risk of anemia. Previous studies have also reported similar findings to the present study (Ositadinma *et al.*, 2016; Aaron *et al.*, 2021). Furthermore, lower level of platelet count documented in the HIV serodiscordant couple in this study is an indication that these persons are at greater risk of thrombocytopenia. This is in

Comment [U7]: does

Comment [U8]: receive

keeping with findings in previous studies involving HIV subjects (Enawgawet *et al.*, 2014; Aaron *et al.*, 2021). Anemia, leucopenia, neutropenia, lymphopenia, and thrombocytopenia have been observed in HIV-infected people before or after ART initiation (Bhardwaj *et al.*, 2020; Damtie *et al.*, 2021), indicating that hematological abnormalities in HIV individuals are induced by either the virus or ART. More so, anemia, lymphocyte count, and thrombocytopenia are associated with CD4 levels (Bhardwaj *et al.*, 2020).

Expectedly, this study found that the CD4+T-cell count was significantly lower in HIV serodiscordant participants compared to controls. This finding is in line with earlier studies (Nsonwu-Anyanwuet *et al.*, 2017; Ezeugwunneet *et al.*, 2021; Ezeugwunneet *et al.*, 2021). A progressive decrease in CD4+ T-cell populations, along with a steady decline in cellular immunity characterizes HIV infection.

However, there was no significant difference in the mean WBC, lymphocyte, or neutrophil counts observed in the discordant group compared to the control group. This finding is consistent with that of Asemota and colleagues, who found no significant differences between control and HIV-naive patients and HIV-infected participants in terms of the mean WBC counts (Asemota *et al.*, 2018). The limited sample size used in this study as well as the stage of HIV infection may have had an impact on the findings.

The present study showed that the mean level of total WBC did not differ statistically significantly from the control groups or between the male and female HIV discordant test groups. Also, the mean NUE/NAC (mixed cells) levels did not differ significantly between the test groups of males and females HIV discordant couples as well as the controls. This finding is consistent with that of Asemota and colleagues, who found no significant differences between control and HIV-naive patients and HIV-infected participants in terms of the mean WBC and

mixed cells counts (Asemota *et al.*, 2018) but differs with some other reports (Aaron *et al.*, 2021). This result may imply that in the participants in the study, gender differences do not significantly affect either the WBC or the mixed cells count (which includes monocyte, eosinophil, and basophil count).

There was significantly lower platelet count observed in the male HIV discordant test group when compared to the female control group. This demonstrates that male HIV-positive discordant couples may be more susceptible to thrombocytopenia than the general or healthy population. Thrombocytopenia has been reported in the literature among the HIV positive individuals previously (Abdulrahman *et al.*, 2014) which agrees with the current results.

Thrombocytopenia is the second most frequent complication of human immunodeficiency virus (HIV) infection with varied prevalence documented across the globe (Akinbami *et al.*, 2010; Gunda *et al.*, 2017; Butale, 2019; Gebreweld *et al.*, 2020; Patil and Patil, 2020).

In this study, the mean level of RBC in the female HIV discordant test group was significantly lower than the levels observed in the female and male control groups, respectively. Additionally, compared to the female control group, the mean RBC level was significantly lower in the male HIV discordant test group. The mean HB level in the female HIV discordant test group was significantly lower than it was in the male and female control groups. When compared to the female and male control groups, the HCT/PCV level in the female HIV discordant test group was significantly lower. Additionally, the mean HCT/PCV level in the male HIV discordant test group was significantly lower than in the female control group. This research demonstrates that while men are similarly susceptible to anemia brought on by HIV infection and exposure, it affects women more severely. Impaired haematopoiesis, immune-mediated processes, opportunistic infections, lymphoma, and the myelotoxic effects of antiretroviral medications are

some of the aetiopathogenesis factors for anemia in HIV infection (Hoffbrand *et al.*, 2006).

Anemia in HIV-positive people has been shown in several investigations as evidenced by significantly lower mean RBC, Hb, and PCV levels (Ositadinma *et al.*, 2016; Aaron *et al.*, 2021). Furthermore, the mean lymphocyte counts were significantly higher in the male HIV discordant test group than in the female HIV discordant test group, but significantly lower in the female HIV discordant test group compared to the female and male control groups. In comparison to the female control group, the mean neutrophil counts in the male HIV discordant test group were significantly lower, but it did not differ significantly from those found in the female test group. This suggests that while the male discordant test group may be more susceptible to neutropenia than the female discordant group, the female discordant group may be more susceptible to lymphopenia. Numerous studies have recorded varying degrees of leucopenia characterized by lymphopenia and/or neutropenia (Balogun *et al.*, 2020) which corroborates with the present findings. Leucopenia and neutropenia have been linked to HIV infection in previous research (Munyaza *et al.*, 2012; Parinitha and Kulkarni, 2012) While neutropenia makes HIV patients more vulnerable to bacterial infections, leucopenia is believed to increase the frequency of opportunistic infections (Tagoe and Asantewaa, 2011). According to reports, lymphopenia and a low lymphocyte total count are frequent symptoms of HIV infection (Dangana *et al.*, 2010; Parinitha and Kulkarni, 2012).

Interestingly, the mean CD4 count was significantly lower in the male and female discordant test groups, respectively, than the values observed in the male and female control groups, while the mean CD4 count was significantly higher in the male discordant test group than in the female discordant test group. This result is consistent with past research (Nsonwu-Anyanwu *et al.*, 2017; Ezeugwunne *et al.*, 2021; Ezeugwunne *et al.*, 2021).

The mean red blood cell count was significantly different between the groups and significantly lower in the female HIV seropositives than in the male and female control groups respectively. There was a significant difference between the groups' mean hemoglobin (HB) levels as well. In comparison to controls, female HIV seropositives and exposed HIV seronegatives had significantly lower levels of hemoglobin (Hb). Furthermore, compared to female controls, PCV/HCT were significantly lower in female HIV-seropositives as well as exposed seronegative males. These results suggest that anemia caused by HIV infection and/or exposure affects both female HIV positives and HIV exposed seronegatives in a similar way. The fact that the current study identified partner HIV exposure as a risk factor that can result in anemia in such individuals is quite significant. Several studies have documented anemia in HIV infected individuals indicated by significantly lower mean levels of RBC, Hb and PCV (Ositadinmaet *et al.*, 2016; Aaron *et al.*, 2021). The most prevalent hematologic abnormality among HIV-positive individuals is anemia, which is also linked to the course of the disease and a higher mortality risk for the patients (Cao *et al.*, 2022). Patients with HIV experience anemia for a variety of reasons. Hematopoietic stem/progenitor cells (HSPCs), which are found in the bone marrow, may be negatively affected by HIV both directly and indirectly (Durandt *et al.*, 2019; Marchionatti and Parisi, 2021). Additionally, the proliferation and differentiation of HSPCs during hematopoiesis may be impacted by ART medications, inflammatory mediators generated during HIV infection, coinfections, or opportunistic infections (Durandt *et al.*, 2019; Marchionatti and Parisi, 2021). So, either the gradual depletion of HSPCs or the inhibition of their action may be the cause of anemia. In addition, a number of publications have noted that some ART combinations or monotherapy can cause anemia (Zerihunet *et al.*, 2019; Kayodeet *et al.*, 2020), which is consistent with the results of the current study. An important indicator that the anemia experienced by the

HIV exposed seronegative participants in this study may be caused by their exposure to HIV infection in their partners is the fact that their HB, RBC, and PCV/HCT were significantly lower than in control subjects. As a result, it is important to monitor these indices in the exposed participants.

As anticipated, this study found that the CD4+T-cell count was significantly lower in the female HIV seropositives, female exposed HIV seronegatives, and male HIV seropositives as compared to the values seen in the male and female control groups. In comparison to female HIV positives, male HIV exposed seronegatives had a considerably lower mean CD4 count. As anticipated, this study found that the CD4+T-cell count was significantly lower in HIV-positive participants compared to controls. This finding is in line with earlier studies (Nsonwu-Anyanwuet *al.*, 2017; Ezeugwunneet *al.*, 2021; Ezeugwunneet *al.*, 2021). A progressive decrease in CD4+ T-cell populations, along with a steady decline in cellular immunity and an increase in vulnerability to opportunistic infections, is the defining feature of HIV infection and, subsequently, AIDS pathogenesis (Okoye and Picker, 2013). Furthermore, previous studies has shown no significant alterations in mean CD4 counts in HIV infected male and female participants (Ogbodoet *al.*, 2021). This finding suggests that seronegative exposed partners also experience diminishing CD4+ T-cell counts as a result of HIV exposure, and as a result, continual depletion of these immune cells makes these people more susceptible to infections that can worsen immunological function. Therefore, it may be appropriate to suggest that these people start taking ART in order to further protect them from the rising viraemia caused by untreated HIV infection.

CONCLUSION

This study has revealed significant changes in CD4-T cell count and hematological indices, most notably significant declines in PCV/HCT, HB, RBC, WBC count as well as CD4-T cell count in

HIV discordant couples, which calls for an urgent interventional strategy to prevent the potential anemia, leucocytopenia, and weakened immunity that may result in both HIV seropositives and seronegative exposed couples.

REFERENCES

Dangana, A., Nuhu, A., & Thomas, K. (2010). Evaluation of Haematological Variations among HIV Infected Patients Attending Antiretroviral Clinic at BarauDikko Specialist Hospital Kaduna State Northwest Nigeria. *International Journal of Biomedical and Healthcare Science*; **6**:215-217.

Tagoe, D.N.A., & Asantewaa, E. (2011). Profiling Haematological Changes in HIV Patients Attending Fevers Clinic at the Central Regional Hospital in Cape Coast, Ghana: A Case-Control Study. *Archives of Applied Science Research*; **3**:326-331.

Munyazesa, E., Emile, I., Mutimura, E., Hoover, D. R., Shi, Q., McGinn, A. P., Musiime, S., Muhairwe, F., Rutagengwa, A., Dusingize, J. C., & Anastos, K. (2012). Assessment of haematological parameters in HIV-infected and uninfected Rwandan women: a cross-sectional study. *BMJ open*; **2**(6):e001600.

Parinitha, S.S., & Kulkarni, M.H. (2012). Haematological Changes in HIV Infection with Correlation to CD4 Cell Count. *Australasian Medical Journal*; **5**:157-162.

Hoffbrand, A.V., Moss, P.A.H., & Pettit, J.E. (2006) *Essential Haematology*. 5th Edition, Blackwell Publishing, Oxford, 328-329.

Abdulrahman, Y., Yeldu, M.H., Zama, I.I., & Amina, A. (2014). Some Haematological Profile of HIV/AIDS Patients on Highly Active Antiretroviral Therapy (HAART) in UsmanuDanfodiyo University Teaching Hospital Sokoto, North Western Nigeria. *American Journal of Science and Technology*; **2**(1):27-32.

Balogun, T.M., Alao, A.O., Olaosebikan, O.F., Aremu, A.J., & Adegbite, O.O. (2020). The Pattern of Haematological Changes in the Baseline Blood Cell Counts and the CD4+ T Lymphocyte Levels among Antiretroviral Therapy Naïve Adult HIV Positive Patients in a Nigerian Hospital. *Open Journal of Blood Diseases*; **10**:77-88.

Aaron, U. U., Okonko, I. O., & Frank-Peterside, N. (2021). Haematological Abnormalities Among HIV Positive Patients on Antiretroviral Treatment in a Nigerian State, South of the Niger Delta. *Biomedicine and Nursing*; **7**(3):1-7.

Enawgaw, B., Alem, M., Addis, Z., & Melku, M. (2014). Determination of hematological and immunological parameters among HIV positive patients taking highly active antiretroviral treatment and treatment naïve in the antiretroviral therapy clinic of Gondar University Hospital, Gondar, Northwest Ethiopia: a comparative cross-sectional study. *BMC hematology*; **14**(1):8.

Asemota, E.A., Okafora, I.M., Okoroiwua, H.U., Ekonga, E.R., Anyanwua, S.O., Efiomb, E.E., & Udomah, F. (2018). Zinc, copper, CD4 T-cell count and some hematological parameters of HIV-infected subjects in Southern Nigeria. *Integrative Medicine Research*; **7**:53–60.

Nsonwu-Anyanwu, A.C., Egbe, E.R., Agu, C.E., Ofors, S.J., Usoro, C.A., & Essien, I.A. (2017). Nutritional indices and cardiovascular risk factors in HIV infection in Southern Nigeria. *Journal of Immunology and Microbiology*; **2**:34–42.

Okoye, A. A., & Picker, L. J. (2013). CD4(+) T-cell depletion in HIV infection: mechanisms of immunological failure. *Immunological Reviews*; **254**(1):54–64.

Okoroiwu, H.U., Okafor, I.M., Uko, E.K., &Atangwho, I.J. (2017). Somehematological parameters of Wistar rats treated with*Chromolaenaodorata*leave extracts. *Journal of Biological Research*; **90**:51–55.

Ositadinma, I.M., Odozi, E.B., Meludu, S.C., &Okeke, C.O. (2016). Effects ofHIV infection on some haematological parameters andimmunoglobulin levels in HIV patients in Benin City, Southern Nigeria. *Journal of HIV and Retrovirus*; **2**:2.

Cao, G., Wang, Y., Wu, Y., Jing, W., Liu, J., &Liu, M. (2022). Prevalence of anemia among people living with HIV: A systematic review and meta-analysis. *eClinical Medicine*; **44**:101283.

Durandt, C., Potgieter, J. C., Mellet, J., Herd, C., Khoosal, R., Nel, J. G., Rossouw, T., & Pepper, M. S. (2019). HIV and haematopoiesis. *South African Medical Journal*; **109**(8b):40–45.

Marchionatti, A., &Parisi, M. M. (2021). Anemia and thrombocytopenia in people living with HIV/AIDS: a narrative literature review. *International Health*; **13**(2):98–109.

Ezeugwunne, I.P., Ogbodo, E.C., Ezeuduji, O.O., Iwuji, J.C., Okwara, N.A., Obi-Ezeani, C.N., Amah, A.K., Odumodu, I.O., &Izuchukwu, E.C.O. (2021). Assessment of Alpha-Fetoprotein, Albumin, Cd4+ and Some Liver Enzymes in HIV Infected Adult on Art in NauthNnewi, South Eastern Nigeria. *Advances in Bioresearch*; **12**(4):199-205

Ezeugwunne, I.P., Ogbodo, E.C., Analike, R.A., Okwara, N.A., Nnamdi, J.C., Iwuji, J.C., Obi-Ezeani, C.N., Amah, A.K., Odumodu, I.O., &Onyenekwe, C.C. (2021). The pattern of alpha-

fetoprotein, CD4+ count, albumin, AST, ALT and ALP in HIV subjects on long term antiretroviral therapy in NauthNnewi, Anambra State, Nigeria. *Indian Journal of Forensic and Community Medicine*; **8**(1):45-51.

Ogbodo, E.C., Ezeugwunne, I.P., Ezeuduji, O.O., Analike, R.A., Okezie, O.A., Onuora, I.J., Okwara, E.C., Nnamdi, J.C., Obi-Ezeani, C.N., & Onyenekwe, C.C. (2021). Gender Comparison Of Alpha-Fetoprotein, CD4, Albumin and Some Liver Enzymes in Symptomatic HIV Subjects on Antiretroviral Therapy. *Journal of Progressive Research in Biology*; **5**(1):17-25.

Duguma, N., TesfayeKiya, G., AdissuMaleko, W., & Bimerew, L. G. (2021). Hematological parameters abnormalities and associated factors in HIV-positive adults before and after highly active antiretroviral treatment in Goba Referral Hospital, southeast Ethiopia: A cross-sectional study. *SAGE open medicine*; **9**: 20503121211020175.

Kayode, E. M., Usiegbodi, D. O., Ajiboye, M. E., Omony, I. S., Febut, M. N., & Buru, A. S. (2020). Assessment of the effect of anti-retroviral therapy on haematological parameters in HIV positive individuals in Zaria. *Journal of AIDS and HIV Research*; **12**(2):17-23.

Zerihun, K.W., Bikis, G.A., & Muhammad, E.A. (2019). Prevalence and associated factors of anemia among adult human immune deficiency virus positive patients on anti-retroviral therapy at Debre Tabor Hospital, Northwest Ethiopia. *BMC Research Notes*; **12**(1):168.

Gebreweld, A., Fiseha, T., Girma, N., Haileslasie, H., & Gebretsadik, D. (2020). Prevalence of cytopenia and its associated factors among HIV infected adults on highly active antiretroviral therapy at MehalMeda Hospital, North Shewa Zone, Ethiopia. *PloS one*; **15**(9):e0239215.

Gunda, D.W., Godfrey, K.G., Kilonzo, S.B., & Mpondo, B.C. (2017). Cytopenias among ART-naive patients with advanced HIV disease on enrolment to care and treatment services at a tertiary hospital in Tanzania: a cross-sectional study. *Malawi Medical Journal*; **29**(1):43–52.

Butale, P. (2019). Study of hematological profile in hiv infected patients with correlation to CD4 cell count. *International Journal of Medical Science and Diagnosis Research*; **3**(2): 171-176.

Akinbami, A., Oshinaike, O., Adeyemo, T., Adediran, A., Dosunmu, O., Dada, M., Durojaiye, I., Adebola, A., & Vincent, O. (2010). Hematologic abnormalities in treatment-naive HIV patients. *Infectious Diseases: Research and Treatment*; **3**:45–49.

Patil, S.S., & Patil, V.S. (2020). Correlation of Blood Profile and CD4 Count in AIDS Patients Before and After HAART, Study in Western Maharashtra. *Biomedical and Pharmacology Journal*; **13** (1):101–105.

UNAIDS. (2022). Global HIV and AIDS statistics — Fact sheet. Retrieved from:

<https://www.unaids.org/en/resources/fact-sheet> (28 June, 2022).

Roth, G. A., Abate, D., Abate, K. H., Abay, S. M., Abbafati, C., Abbasi, N., Abbastabar, H., Abd-Allah, F., Abdela, J., Abdelalim, A., Abdollahpour, I., Abdulkader, R. S., Abebe, H. T., Abebe, M., Abebe, Z., Abejie, A. N., Abera, S. F., Abil, O. Z., Abraha, H. N., Abrham, A. R. & 30 others. (2018). Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*; **392**(10159):1736–1788.

James, S.L., Abate, D., Abate, K. H., Abay, S. M., Abbafati, C., Abbasi, N., Abbastabar, H., Abd-Allah, F., Abdela, J., & Abdelalim, A. (2018). Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*; **392**(10159):1789–1858.

Awofala, A.A., & Ogundele, O.E. (2018). HIV epidemiology in Nigeria. *Saudi Journal of Biological Science*; **25**(4):697-703.

UNFPA. (2020). World population dashboard Nigeria, 2020.

https://scholar.google.com/scholar_lookup?title=World%20population%20dashboard%20Nigeria&publication_year=2020 (Retrieved June 29, 2022).

UNAIDS (2019). Press Release

[https://www.unaids.org/en/resources/presscentre/pressreleaseandstatementarchive/2019/march/20190314_nigeria]. Retrieved June 29, 2022.

vanWoudenberg, E., Irvine, E.B., Davies, L., de Kock, M., Hanekom, W.A., Day, C.L., Fortune, S., & Alter, G. (2020). HIV Is Associated with Modified Humoral Immune Responses in the Setting of HIV/TB Coinfection. *mSphere*; **5**(3):e00104-20.

Balasubramaniam, M., Pandhare, J., & Dash, C. (2019). Immune Control of HIV. *Journal of Life Science* (Westlake Village); **1**(1):4-37.

Damtie, S., Workineh, L., Kiros, T., Eyayu, T., & Tiruneh, T. (2021). Hematological Abnormalities of Adult HIV-Infected Patients Before and After Initiation of Highly Active Antiretroviral Treatment at Debre Tabor Comprehensive Specialized Hospital, Northcentral Ethiopia: A Cross-Sectional Study. *HIV AIDS* (Auckl); **13**:477-484.

Bhardwaj, S., Almaeen, A., Ahmed, Wani, F., & Thirunavukkarasu, A. (2020). Hematologic derangements in HIV/AIDS patients and their relationship with the CD4 counts: a cross-sectional study. *International Journal of Clinical and Experimental Pathology*; **13**(4):756-763.

Tsukamoto, T. (2020). Hematopoietic stem/progenitor cells and the pathogenesis of HIV/AIDS. *Frontiers in Cellular and Infection Microbiology*; **10**:60.

Ravikumar, B., & Balakrishna, P. (2013). Discordant HIV Couple: Analysis of the Possible Contributing Factors. *Indian Journal of Dermatology*; **58**(5):405.

Chemaitelly, H., Cremin, I., Shelton, J., Hallett, T.B., & Abu-Raddad, L.J. (2012). Distinct HIV discordancy patterns by epidemic size in stable sexual partnerships in sub-Saharan Africa. *Sexually Transmitted Infections*; **88**:51.

Curran, K., Baeten, J.M., Coates, T.J., Kurth, A., Mugo, N.R., & Celum, C. (2012). HIV-1 prevention for HIV-1 serodiscordant couples. *Current HIV/AIDS Reports*; **9**:160.

Irungu, E.M., Heffron, R., Mugo, N., Ngure, K., Katabira, E., Bulya, N., Bukusi, E., Odoyo, J., Asimwe, S., Tindimwebwa, E., Celum, C., Baeten, J.M., & Partners Demonstration Project Team (2016). Use of a risk scoring tool to identify higher-risk HIV-1 serodiscordant couples for an antiretroviral-based HIV-1 prevention intervention. *BMC Infectious Disease*; **16**:1.

Patel, R.C., Leddy, A.M., Odoyo, J., Anand, K., Stanford-Moore, G., Wakhungu, I., Bukusi, E.A., Baeten, J.M., & Brown, J.M. (2018). What motivates serodiscordant couples to prevent HIV transmission within their relationships: findings from a PrEP implementation study in Kenya. *Culture, Health & Sexuality*; **20**:625.

UNDER PEER REVIEW