

In vitro Estimation of Phytochemicals & flavonoids in *Citrus sinensis* & study their Antimicrobial & Antioxidant activity

ABSTRACT:

Orange (*C. sinensis*) peel flavonoids have antibacterial, antioxidant, anticancer, anti-inflammatory, and antiviral effects. However, there is a scarcity of information in the literature about orange peels flavonoids. So, we estimated phytochemicals in hydro-methanolic extract of *Citrus sinensis* peel and separated the flavonoids by Column chromatography and confirmed with TLC and HPLC. After that we evaluated the antimicrobial, antioxidant and nitric oxide scavenging activity. The MIC value of flavonoids against *E. coli*, *B. subtilis*, *S. aureus* and *P. aeruginosa* is 19.40, 20.48, 12.45 and 21.44 μ g respectively. The IC₅₀ value of flavonoids for antioxidant and Nitric oxide scavenging activity is 121.4 and 0.88 μ g respectively. In the current study, *Citrus* peel flavonoids were found to have antibacterial, antioxidant, and nitric oxide scavenging action in methanolic extracts. This study suggests that *C. sinensis* flavonoids could be exploited as an antibacterial and antioxidant source in the future.

Keyword: *Citrus sinensis*, Flavonoids, Antimicrobial activity, MIC, Antioxidant and Nitric Oxide scavenging Activity

INTRODUCTION:

Around 80-85 percent of the total population relies on Ayurveda for disease treatment, which could be due to its ease of accessibility, low cost, and lack of unfavourable side effects (Ekor, 2014). Ayurveda is very popular among patients and is widely available in developing countries such as India, Pakistan, Bhutan, Srilanka and China (Rastogi, 2021, Chaterjee et al, 2012). India is the world's leading producer of herbal medicine and the birthplace of Ayurveda (Mills et al, 2019). *Citrus sinensis* (Orange) is a juicy fruit, belonging to the *Rutaceae* family and the most widely cultivated plant species in the world. Orange peel is proven to have many medicinal properties as a tonic for the digestive and respiratory system, immune system, skin, antiviral, antibacterial, antioxidant and anticancer (Favela-Hernández et al, 2016, Pepe et al, 2017). Orange peel extract showed antibacterial activity against *Klebsiella pneumonia*, *Streptococcus mutans*, *Lactobacilli acidophilus*, *Pseudomonas aeruginosa*, *Escherichia coli* and *Bacillus subtilis* (Zubair, 2020, Pepe et al, 2017).

The reactive oxidizing and nitrogenous species participate in the oxidation of many biological processes as protective hosts and generate more free radicals such as nitric oxide, O₂, \cdot OH and damage bio-molecules such as lipids, proteins and nucleic acids (Meo et al, 2020, Trinity et al, 2016). It contributes to immune pathologies including cancer, inflammation, diabetes, hypertension, AIDS and aging, etc (López-Navarro et al, 2020). Antioxidants have the ability to reduce or inhibit the oxidation of bio-molecules such as lipids, proteins, etc (Kumar et al, 2020). Food degradation with lipid peroxidation affects the shelf life of food due to the overproduction of free

radicals. During the manufacturing process, the use of antioxidants can extend the shelf life of foods (Lourenço et al, 2019). In recent times, there is a tendency to use natural phytochemicals as sources of antibacterial and antioxidant substances found in all plants and their parts. Many phytochemicals such as phenolic content are used as a source of antioxidants to reduce oxidative degradation in food and maintain food quality in the long term (Gutiérrez-Del-Río et al, 2021, Shehata et al, 2021).

Citrus sinensis (Orange) is a widely distributed species of plant in the family Rutaceae. It is grown in all regions for its juices of flavor and medicinal importance. Orange peel also has medicinal properties used to strengthen the immune system, digestive system, vitamin deficiencies and skin. Orange peel extracts and flavonoids have shown antibacterial and antioxidant activities. Flavonoids are effective against antibiotic-resistant bacteria, Gram-positive and Gram-negative bacteria, AIDS, Covid-19 and cancer (Sharma et al, 2019, Koolaji et al, 2020, Antonio et al, 2021).

With my research of the literature, I did not find any studies on the flavonoids in orange peels on bacteria, antioxidants and nitric oxide scavenging activity. Therefore, the present work was carried out with the given objective:

1. Evaluation of phytochemical estimates and isolation and purification of flavonoids from orange peels.
2. Evaluation of antibacterial activity and determination of minimum inhibitory concentration (MIC) of flavonoids
3. For the evaluation of antioxidants, the scavenging activity of nitric oxide and the determination of the IC₅₀ value in order to find new drugs against bacteria and free radicals.

MATERIAL AND METHOD:

Orange peels were collected near to Chandigarh, India and identified by Gaurav Srivastav (Botanist) and dried at room temperature in absence of light after washed. Obtained powder (100gm) after milling was subjected to extraction with methanol & water in ratio of 60:40 at Soxhlet for 24hrs. The obtained extract was concentrated in oven at 60⁰C and quantify the extracts recovered mass by following formula and store at 4-8⁰C for further used.

Y (%) = Mass of extract / mass of sample X 100.

Phytochemical Screening:

A minimum amount of extracts were used for phytochemical screening for the compound as total phenol, alkaloids, flavonoids, proteins, amino acid, tannins, saponins, steroids, terpenoids, carbohydrate, reducing sugar and glycosides (Kumar et al, 2020, Harborne, 1998). According to selected method with little modification, 2.0gm of plant extract was dissolved in 10ml double distilled water and filtered with filter and further used for phytochemical screening:

- **Test for alkaloid:** Take 3.0 ml filtrate extract solution with addition of few drops Dragendorff's reagent, Appearance of an orange-red colour precipitate indicates the presence of alkaloids.
- **Test for phenolic compound:** To 1.0 ml of the extract, 3.0 ml of distilled water followed by few drops of 10% aqueous Ferric chloride solution was added. The origin of blue or green colour indicates the presence of phenols.
- **Test for flavonoids:** Take 2.0 ml of test solution and add a small quantity of reduced lead acetate solution (10%). The solution will form yellow colour ppt. In addition to an increasing amount of sodium hydroxide to the residue, Yellow colour of a solution formed which decolourizes after the addition of acids.
- **Test for carbohydrate:** Take 2.0 ml extract and add few drops of α -naphthol solution in alcohol. Shake and add 2.0 ml conc. H_2SO_4 from the side of test tube. A violet/purple ring at the junction of two liquids appeared.
- **Test for reducing sugar:** Add 20 ml of Fehling (A+B) to 5.0 ml of extract in a test tube and boil for 15mins. Then neutralize with 10% sodium hydroxide to pH-7.00. Solution appeared brick red ppt.
- **Test for glycosides:** Take 2.0 ml extract solution and add 1.0 ml glacial acetic acid and 1.0ml 5% $FeCl_3$ and 1.0ml conc. H_2SO_4 . The reddish-brown colour at the junction of two liquid layers appeared and the upper layer appears bluish-green colour.
- **Test for protein:** Take 3.0 ml test solution and added 4% NaOH and few drops of 1% $CaSO_4$ solution. The solution appeared violet in colour.
- **Test for amino acid:** Take 5.0 ml solution, adds few drops of 40 % NaOH and 10 % lead acetate solution and then boil it. Black ppt. of lead sulphate formed.
- **Test for steroid:** To 1.0ml of extract and added 1.0ml of chloroform, 2-3ml of Acetic Anhydride, 1 - 2 drops of conc. H_2SO_4 . Dark green colour showed.
- **Terpenoids:** Take 2.0ml of extract solution, add 2.0ml of chloroform and 2.0ml of conc. H_2SO_4 and shake well. Chloroform layer (upper) appeared red in colour and acid layer (lower) formed greenish-yellow fluorescence.
- **Tannin:** 1.0 ml of water extract was mixed with 10 ml of distilled water and the filtered. Ferric chloride ($FeCl_3$) reagent 2-4 drops were added to filtrate. A blue-black or green precipitate confirmed the presence of tannins.
- **Saponins:** Take 0.5ml of extract and 5.0ml of water and shake well. Foam formed.

Determination of Total Phenolic Component:

The total amount of phenolic contents was determined using the method by Folin Ciocalteu's phenol reagent with slight modification. 1.0ml of the sample was mixed with 2.0ml of Folin Ciocalteu's phenol reagent (10%). After 3.0min, 1.0 ml of

saturated Na_2CO_3 (~35 %) was added to the mixture and made up to 10 ml by adding distilled water. The reaction was kept in the dark for 90min, observed under UV-Vis spectrophotometer at 760nm absorbance. Tannic acid was used as a standard with varied concentrations from 50 ppm to 200 ppm. The results were expressed as mg of Tannic acid equivalents/g of extract and the same procedure was done with extract and calculated as mean value \pm SD (n = 3) (Kumar et al, 2020, Harborne, 1998).

Determination of Flavonoid Content:

Flavonoids content of the isolated crude extract was determined by method with slight modification. Take 0.5ml of the sample (Extract) containing 1.25ml of distilled water. Then added 0.075ml of 5 % sodium nitrite solution and allowed to stand for 5 min. Added 0.15ml of 10 % aluminium chloride, after 6 min 0.5ml of 1.0M sodium hydroxide was added and the mixture was diluted with another 0.275ml of distilled water. The absorbance of the mixture at 510nm was measured immediately. The flavonoid content was expressed as mg of Quercetin equivalents/g sample and the same procedure was done with Extract of peel and calculated as mean value \pm SD (n=3) (Kumar et al, 2020, Harborne, 1998).

Separation & Purification of flavonoids from Crude extract of peels

Combined solvent (methanol: water) was used as mobile phase in silica gel column chromatography (200g, 60-200 mesh). 10gm extracts were loaded in a glass column with silica gel and a solvent system (methanol: water) was used for running the sample. For confirmation of separation, thin layer chromatography was performed by taking sample with mobile phase (Glacial acetic acid: Methanol: water) using 10 μ l of sample and fractions were further screened for flavonoids (Yahaya et al, 2015, Kumar et al, 2020).

For determination of wavelength, spectrophotometry was considered on UV/Visible spectrophotometer with spectra (200-500nm) using Quercetin dihydrate as standard. 1ml of fraction, 1ml pure water, 5ml acetate buffer (pH-3.8 \pm 0.5), and 3ml AlCl_3 (0.1M) solution added and after shaking kept for 30minuts (Kumar et al, 2019). Absorbance was measured and considered wavelength and then, reversed-phase high-performance chromatography (HPLC) was performed for the determination of flavonoids. The 20 μ l of column chromatography purified liquid extracts was injected into HPLC (reversed-phase, C18, 250mm x 416mm, Shimadzu) at 350nm for 16min at 1.0ml/min with HPLC grade Methanol: Water (60:40) as mobile phase (Yahaya et al, 2015, Kumar et al, 2020, Kumar et al, 2019).

Antibacterial Activity and MIC determination:

Four pathogenic bacteria namely *Escherichia coli* (MTCC 25922), *Bacillus subtilis* (MTCC 3256), *Pseudomonas aeruginosa* (MTCC 1688) and *Staphylococcus aureus* (MTCC 6810) were collected from MTCC, Chandigarh. After subculture, Agar well diffusion was used for antimicrobial activity with slight manipulation. 0.5ml (<1.0 x 10⁵ CFU/ml) of 48 hrs old cultures of test organisms were inoculated into different sterile Petri-plates and 15-20ml sterile Muller Hilton agar media was poured into

each Petri-plate (90 x 90mm). The Petri-plates were gently shaken for proper mixing and allowed to solidify. Thereafter, four wells were punched of 5mm diameter with a sterilized cork borer. For each well 50µl (25 - 100µg/ml) of flavonoids were added. The plates were incubated at 37⁰C for 24hrs and then the zone of inhibition was measured in mm (Kumar et al, 2019). The experiment was carried out in triplicates and calculated as mean value ± SD (n = 3) and manually calculated minimum inhibitory concentration (MIC) (Kumar et al, 2019, Musa et al, 2019).

Antioxidant Activity:

The antioxidant activity of the flavonoids and the standard was assessed on the basis of the radical scavenging effect of the stable 1, 1-diphenyl-2-picrylhydrazyl (DPPH-0.002% in methanol) method (Kumar et al, 2019). The diluted working solution of the standard (Ascorbic acid) and flavonoids (10mg/ml) was prepared using the respective solvents. In 3ml of the total reaction solution, 2ml of flavonoids/standard solution (200-1000µg/ml) and 1.0ml of DPPH were mixed and allowed to react at 37⁰C for 30 min. afterward; absorbance values were measured at 520nm and converted into percent antioxidant activity. The percentage antioxidant activity was calculated by the following formula and calculated as mean value ± SD (n = 3) (Lu et al, 2019, Liew et al, 2018).

Percent (%) inhibition of DPPH activity = $\frac{(A - B)}{A} * 100$

Where A = Absorbance of the blank
 B = Absorbance of the sample

Nitric Oxide Radical Scavenging Activity:

Scavengers of nitric oxide are complete with oxygen, leading to reduced production of nitrite ions. The standard (Ascorbic acid) and *Citrus* peel flavonoid were dissolved in distilled water. 2ml Sodium Nitroprusside (5mM) in phosphate buffer saline (0.025mM, pH-7.4) was incubated with 0.5ml of different concentrations (200-1000µg/ml) of standard and flavonoids were incubated at 30⁰C for 3 hrs. After 3 hrs incubation, samples were diluted with 1ml of Griess reagents (prepared by mixing equal volume of 1% sulphanilamide in 2% phosphoric acid and 0.1% Naphthylethylene diamine dihydrochloride in water) and incubated for 30min (Kumar et al, 2020). The absorbance of the colour developed during diazotization of nitrite with sulphanilamide and its subsequent coupling with Naphthylethylene diamine hydro-chloride was observed at 550nm on a spectrophotometer. Calculated the % inhibition by formula and plot graph in compared to standard and calculated as mean value ± SD (n = 3) (Chang et al, 2016, Kim et al, 2019).

Nitric Oxide Scavenging (%) = $\frac{A_{control} - A_{test}}{A_{control}} * 100$

Where, A_{control} = Absorbance of control reaction
 A_{test} = Absorbance in the presence of the samples of extract

RESULT & DISCUSSION:

Herbs contain a variety of ingredients that can be separated using chromatography techniques. There are numerous references available for describing the various components and their functions. Flavonoids, a phenolic compound, have been shown in recent studies to be antimicrobial agents, antioxidants, and nitric oxide scavengers. Flavonoids, which are secondary metabolites of plants, have antibacterial, antioxidant, cleansing, carcinogenic, and other disease-fighting properties (Barreca et al, 2017, Suntar et al, 2018).

The presence of total phenol, alkaloids, flavonoids, proteins, amino acids, tannins, saponins, steroids, terpenoids, carbohydrate, reducing sugar, and glycosides were observed in a hydro-methanolic extract of *Citrus sinensis*. During the screening, we discovered that alkaloids, phenolic compounds, steroids, terpenoids, flavonoids, tannins, saponins, and reducing sugar seem to have been present, but glycosides, carbohydrates, proteins, and amino acids have not been (Table-2). The peels' phenolic content, flavonoids, terpenoids, and alkaloids may contribute to antimicrobial, antioxidant, anti-inflammatory, and anti-cancer activity. We estimated the content of phenolic and flavonoids using a spectrophotometer after acknowledging their existence in the peels. 26.5mg hydro-methanolic extract was found in 300gm of orange peels. The total phenolic content and flavonoids in hydro-methanolic extract powder were 32.4µg Tannic Acid equivalent/gm and 62.59µg Quercetin equivalent/gm, respectively, using the standard curve (tannic acid- $y=0.015x + 1.004$, $R^2 = 0.994$ and Quercetin- $y=0.001x + 0.019$, $R^2 = 0.998$) (Table-1). The solvent system (Methanol + Water) was used for column chromatography, spectrum and HPLC and total of five fractions were collected. After qualitative confirmation of flavonoids, the entire fraction were mixed and further screened for re-confirmation. The yield is 62.59µg and had percentage of 0.21%. By spectrum scanning of wavelength of flavonoids sample, the corresponding wavelength of most absorption peak had been identified in Orange – 287nm and 332nm although HPLC was performed at 350nm due to common similarity with standard spectrum (Figure - 1).

Table:1- Crude Extracts, Total Phenolic & Flavonoids content of *Citrus sinensis* peels

Plant Name	Plant Part	Quantity (gm)	Extract (mg)	Total Phenolic Content (µg of TAE/serving)	Flavonoids (µg of QUE /serving)
<i>Citrus sinensis</i>	Peel	300	26.5	32.43	62.59

Table: 2 - Qualitative phytochemical analysis

Parameter	Alkaloids	Glycosides	Reducing sugar	Phenolic compound	Steroids	Terpenoids
Citrus peel	+	-	+	+	+	+
Parameter	Carbohydrate	Protein	Amino	Flavonoid	Tannin	Saponins

r	s		acids	s	s	
Citrus peel	-	-	-	+	+	+




Figure:1- HPLC graph of Orange peels Flavonoids

Figure: 02- Antimicrobial Activity of *Citrus sinensis* peels Flavonoids

Flavonoids were tested for antimicrobial activity against four microorganisms (*Escherichia coli*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*) using the agar plate diffusion method. Ciprofloxacin was used as the standard, and the zone of inhibition was measured in millimetre. For each pathogen, the flavonoids concentrations are 25, 50, 75, and 100µg/ml. Flavonoids demonstrated concentration dependent antimicrobial activity against pathogens, with the highest zone of inhibition at 100µg/ml for *E. coli* and *B. subtilis*. As shown in Figure-02, the zone of inhibition for standard (Ciprofloxacin-30µg/ml) is 28, 30, 27,

32mm for selected pathogens. In comparison to *S. aureus* and *P. aeruginosa*, *E. coli* and *B. subtilis* are more susceptible to flavonoids. Despite the reality that pathogen antimicrobial susceptibility is closer to standard than sample. However, when we calculated the minimum inhibitory concentration (MIC), we discovered that *P. aeruginosa* ($21.45 \pm 2.21 \mu\text{g/ml}$) outperformed *E. coli* ($19.41 \pm 2.75 \mu\text{g/ml}$), *B. subtilis* ($20.49 \pm 4.32 \mu\text{g/ml}$), and *S. aureus* ($12.46 \pm 1.41 \mu\text{g/ml}$).

The antioxidant properties of plants or phytochemicals are tested using various methods such as DPPH, hydrogen peroxide, and nitric oxide. The antioxidant activity of flavonoids found in orange peel was studied in comparison to Ascorbic acid (standard). The DPPH scavenging activity was determined using different concentrations of ascorbic acid and flavonoids, which were 100, 400, 600, 800, and 1000 $\mu\text{g/ml}$, respectively (Figure-03). The interpretation of the data revealed that as the concentration of flavonoids increased, so did the scavenging activity. Similarly, the IC_{50} was calculated using regression analysis of the graph and was found to be $10.06 \pm 2.19 \mu\text{g}$ for ascorbic acid and $121.41 \pm 0.21 \mu\text{g}$. By scanning the wavelength of five flavonoids sample, the corresponding wavelength of most absorption peak had been identified in Orange – 287nm, 332nm g for orange flavonoids. Free radicals start chain reactions, which can cause cell damage or death. Flavonoids act as antioxidants by reducing oxidation reactions or removing free radicals through neutralization with peroxide, lipid peroxy, and/or hydroperoxide.




Figure: 03 -Antioxidant Activity of *Citrus sinensis* peels Flavonoids

Flavonoids from a hydro-methanolic extract of orange peels were evaluated for nitric oxide scavenging activity. The nitric oxide scavenging activity of flavonoids was tested at concentrations ranging from 200 to 1000 $\mu\text{g/ml}$ (Figure - 4). The IC_{50} value of flavonoids was $0.88 \pm 0.21 \mu\text{g}$, calculated using the standard curve (Ascorbic acid = $y=0.070x$ and $r^2=-42.6$). The flavonoids demonstrated free radicals scavenging by participating in scavenge of nitrite radicals generated by sodium nitroprusside at neutral pH. The antioxidant activity increased with increasing flavonoid concentration, which is concentration dependent (Shen et al, 2020, Miyake et al, 1999).

Free radical scavenging activity of flavonoids against DPPH and nitric oxide is assessing that higher scavenging activity; more anti-oxidizing agent might be possessing anticancer properties.

Figure: 04 - Nitrite scavenging Activity of *Citrus sinensis* peels Flavonoids

CONCLUSION:

The presence of active compounds such as phenolic content, alkaloids, carbohydrates, terpenoids, and flavonoids was observed in the phytochemical analysis of orange peels. It contains a high concentration of phytochemicals that have biological activity. Flavonoids were also found to have antimicrobial and antioxidant properties. As a result of these findings, plant extracts and their flavonoids with significant biological activity should be chosen. Further research into pure flavonoids may open the way for the discovery of a core constituent of *Citrus sinensis* peels as a biological activity of a pure or mixed compound.

NOTE:

The study highlights the efficacy of "Ayurved" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

REFERENCE:

1. Ekor M. (2014). The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Frontiers in pharmacology*, 4, 177.
2. Chatterjee, B., Biswas, P. C., & Pancholi, J. (2012). Health awareness and popularity of alternative medicines among people of Jamnagar town: A cross-sectional study. *Ayu*, 33(1), 33–37.
3. Rastogi S. (2021). Emanating the specialty clinical practices in Ayurveda: Preliminary observations from an arthritis clinic and its implications. *Journal of Ayurveda and integrative medicine*, 12(1), 52–57.
4. Mills, P. J., Peterson, C. T., Wilson, K. L., Pung, M. A., Patel, S., Weiss, L., Kshirsagar, S. G., Tanzi, R. E., & Chopra, D. (2019). Relationships among

classifications of ayurvedic medicine diagnostics for imbalances and western measures of psychological states: An exploratory study. *Journal of Ayurveda and integrative medicine*, 10(3), 198–202.

5. Favela-Hernández, J. M., González-Santiago, O., Ramírez-Cabrera, M. A., Esquivel-Ferriño, P. C., & Camacho-Corona, M. (2016). Chemistry and Pharmacology of Citrus sinensis. *Molecules (Basel, Switzerland)*, 21(2), 247.
6. Pepe, G., Pagano, F., Adesso, S., Sommella, E., Ostacolo, C., Manfra, M., Chieppa, M., Sala, M., Russo, M., Marzocco, S., & Campiglia, P. (2017). Bioavailable Citrus sinensis Extract: Polyphenolic Composition and Biological Activity. *Molecules (Basel, Switzerland)*, 22(4), 623.
7. Zubair M. (2020). Antimicrobial and Anti-Biofilm Activities of Citrus sinensis and Moringa oleifera Against the Pathogenic Pseudomonas aeruginosa and Staphylococcus aureus. *Cureus*, 12(12), e12337.
8. Shehata, M. G., Awad, T. S., Asker, D., El Sohaimy, S. A., Abd El-Aziz, N. M., & Youssef, M. M. (2021). Antioxidant and antimicrobial activities and UPLC-ESI-MS/MS polyphenolic profile of sweet orange peel extracts. *Current research in food science*, 4, 326–335.
9. Gutiérrez-Del-Río, I., López-Ibáñez, S., Magadán-Corpas, P., Fernández-Calleja, L., Pérez-Valero, Á., Tuñón-Granda, M., Miguélez, E. M., Villar, C. J., & Lombó, F. (2021). Terpenoids and Polyphenols as Natural Antioxidant Agents in Food Preservation. *Antioxidants (Basel, Switzerland)*, 10(8), 1264.
10. Kumar A., Mahajan A.2, Begum Z.. Phytochemical Screening and In vitro Study of Free Radical Scavenging Activity of Flavonoids of Aloe vera. *Research J. Pharm. and Tech* 2020; 13(2):593-598.
11. Di Meo, S., & Venditti, P. (2020). Evolution of the Knowledge of Free Radicals and Other Oxidants. *Oxidative medicine and cellular longevity*, 2020, 9829176.
12. Trinity, J. D., Broxterman, R. M., & Richardson, R. S. (2016). Regulation of exercise blood flow: Role of free radicals. *Free radical biology & medicine*, 98, 90–102.
13. López-Navarro, M. E., Jarquín-Martínez, M., Sánchez-Labastida, L. A., Ramírez-Rosales, D., Godínez-Victoria, M., Quintas-Granados, L. I., & Trujillo-Ferrara, J. G. (2020). Decoding Aging: Understanding the Complex Relationship among Aging, Free Radicals, and GSH. *Oxidative medicine and cellular longevity*, 2020, 3970860.
14. Lourenço, S. C., Moldão-Martins, M., & Alves, V. D. (2019). Antioxidants of Natural Plant Origins: From Sources to Food Industry Applications. *Molecules (Basel, Switzerland)*, 24(22), 4132.
15. Sharma, Kavita, Mahato, Neelima and Lee, Yong Rok. "Extraction, characterization and biological activity of citrus flavonoids" *Reviews in Chemical Engineering*, vol. 35, no. 2, 2019, pp. 265-284.
16. Koolaji, N., Shammugasamy, B., Schindeler, A., Dong, Q., Dehghani, F., & Valtchev, P. (2020). Citrus Peel Flavonoids as Potential Cancer Prevention Agents. *Current developments in nutrition*, 4(5), nzaa025.

17. Antonio, A. S., Wiedemann, L., Galante, E., Guimarães, A. C., Matharu, A. S., & Veiga-Junior, V. F. (2021). Efficacy and sustainability of natural products in COVID-19 treatment development: opportunities and challenges in using agro-industrial waste from *Citrus* and apple. *Heliyon*, 7(8), e07816.
18. Garrat DC: The quantitative analysis of drugs. 3rd edition. Japan. Chapman and Hall. 1964.
19. Harborne, J.B. (1998) Textbook of Phytochemical Methods. A Guide to Modern Techniques of Plant Analysis. 5th Edition, Chapman and Hall Ltd, London, 01-36.
20. Harborne, J.B. (1998) Textbook of Phytochemical Methods. A Guide to Modern Techniques of Plant Analysis. 5th Edition, Chapman and Hall Ltd, London, 37-99.
21. Yahaya Mobmi Musa. Isolation and Purification of Flavonoids from the leaves of locally Produced *Carica papaya*. International Journal Of Scientific and Technology Research, 2015, 4:282-284.
22. Kumar, A., Sharma, P., Mahajan, A., and Begum, Z. In Vitro Comparative Study of Antioxidant and Antibacterial Activity of Selected Dietary Plants. Nepal Journal of Biotechnology, 2019, 6(1):39-45.
23. Musa, D.D.; Sangodele, F.; Hafiz, S.S. Phytochemical analysis and antibacterial activity of orange (*Citrus sinensis*) peel. FUDMA J. Sci. 2019,3, 375–380
24. Chang, Y. H., Seo, J., Song, E., Choi, H. J., Shim, E., Lee, O., & Hwang, J. (2016). Bioconverted Jeju Hallabong tangor (*Citrus kiyomi* × *ponkan*) peel extracts by cytolase enhance antioxidant and anti-inflammatory capacity in RAW 264.7 cells. *Nutrition research and practice*, 10(2), 131–138.
25. Kim, C., Ji, J., Ho Baek, S., Lee, J. H., Ha, I. J., Lim, S. S., Yoon, H. J., Je Nam, Y., & Ahn, K. S. (2019). Fermented dried *Citrus unshiu* peel extracts exert anti-inflammatory activities in LPS-induced RAW264.7 macrophages and improve skin moisturizing efficacy in immortalized human HaCaT keratinocytes. *Pharmaceutical biology*, 57(1), 392–402.
26. Lu, Q., Huang, N., Peng, Y., Zhu, C., & Pan, S. (2019). Peel oils from three *Citrus* species: volatile constituents, antioxidant activities and related contributions of individual components. *Journal of food science and technology*, 56(10), 4492–4502.
27. Liew, S. S., Ho, W. Y., Yeap, S. K., & Sharifudin, S. (2018). Phytochemical composition and in vitro antioxidant activities of *Citrus sinensis* peel extracts. *PeerJ*, 6, e5331.
28. Shen, C. Y., , Lin, J. J., , Jiang, J. G., , Wang, T. X., , & Zhu, W., (2020). Potential roles of dietary flavonoids from *Citrus aurantium* L. var. *amara* Engl. in atherosclerosis development. *Food & function*, 11(1), 561–571.
29. Miyake, Y., Murakami, A., Sugiyama, Y., Isobe, M., Koshimizu, K., & Ohigashi, H. (1999). Identification of coumarins from lemon fruit (*Citrus limon*) as inhibitors of in vitro tumor promotion and superoxide and nitric oxide generation. *Journal of agricultural and food chemistry*, 47(8), 3151–3157.

30. Barreca, D., Gattuso, G., Bellocco, E., Calderaro, A., Trombetta, D., Smeriglio, A., Laganà, G., Daglia, M., Meneghini, S., & Nabavi, S. M. (2017). Flavanones: Citrus phytochemical with health-promoting properties. *BioFactors (Oxford, England)*, *43*(4), 495–506.
31. Suntar, I., Khan, H., Patel, S., Celano, R., & Rastrelli, L. (2018). An Overview on *Citrus aurantium* L.: Its Functions as Food Ingredient and Therapeutic Agent. *Oxidative medicine and cellular longevity*, *2018*, 7864269.