

Review Article

Internal Transcribed Spacer (ITS) Region as Molecular Marker for Parasite Identification in Aquaculture –A Review

ABSTRACT

Molecular techniques are potentially faster and more sensitive than the culture, serology, and histology methods traditionally used to identify fish parasites. Parasite can be detected from asymptomatic fish by molecular diagnostic techniques so that disease outbreaks can be prevented. This review ~~journal~~ discusses the use of Internal Transcribed Spacer (ITS) markers to detect parasite in aquaculture and their potential applications are discussed. It was found that ~~it~~ is a more reliable marker and species identification in combination with other DNA markers. ITS region are highly polymorphic non-coding region with sufficient taxonomic units. Thus, sequences can be separated at the species level. ~~It~~ Even though ribosomal ITS as a universal barcode marker for parasite identification is still hampered by few limitations, the ITS will remain as the key choice for parasite identification. It remains the most important option for parasite identification. Many studies have been carried out using ITS markers for parasites identification such as cestode, nematodes, *Pseudoterranova* spp., *Anisakis* type1, *Diplostomum*, *Gyrodactylids* *Hysterothylacium* sp, *Glochidia* larvae. This identification of parasites are taken from the species Rainbow trout (*Oncorhynchus mykiss*), Argentine sea bass, (*Acanthistius patachonicus*), and other samples from various places.

Formatted: Highlight

Keywords: ITS (Internal Transcribed Spacer), Molecular Marker, Parasite, Aquaculture

1. Introduction

Aquaculture plays an important role in providing animal protein for the global community, and Indonesia is no exception (1). Therefore, fisheries are currently in great demand by various groups and the demand for its products continues to increase. However, in the production of fisheries, there are many obstacles, including parasitic attacks. This parasite can cause significant production losses by interfering with fish health. The high pathological effect of this parasite can cause death in the host (2).

This is partly because the risk factors associated with aquaculture production in infectious aquatic environments are analyzed and the reasons for the absence of zoonotic parasites in trout aquaculture are pinpointed. Based on the analysis recommendations for management practices securing parasite freedom in cultured fish are given (3). One of

them based on research revealed that the life cycle of *P. cattani* mainly involves demersal and benthic organisms, with preferences marked by the surrounding environment (4).

Fish disease management is a major concern for aquaculture activities. The ability to quickly identify the presence of pathogenic organisms in fish would have significant economic benefits. An additional benefit will also be obtained if concentrations of infectious organisms can be determined in fish or the environment so that changes in the abundance of these pathogenic organisms can be monitored. The development of a system that can evaluate fish parasites and disease carriers in an introduced area will also assist in the development of aquaculture management programs (5).

In controlling parasites, it is still hampered due to difficulties in identifying them. Most parasitic eggs have very similar morphological characteristics, so identification down to the

species level is not always possible with a microscope(6). The fact that certain species of parasite infect more successfully than other species of the same genus is also true in the case of treatment where some strains show sensitivity to certain drugs while others are resistant to the same drug. Therefore, studying and identifying organisms molecularly is important to know their taxonomic status, whether the species are distant or close relatives (2).

For example parasites species *Hysterothylacium amoyense* was the most common species (prevalence 47.2%, mean intensity 14.9 ± 17.1). *Hysterothylacium thalassini* and *R. lophii* were reported in Japanese freshwater fish for the first time. Two distinct genotypes of *A. typica* (3.4% overall prevalence; mean intensity 1.7 ± 0.9) were discovered in the South China Sea for the first time (7).Therefore, one solution is to carry out molecular identification which makes it possible to detect the unique species-specific nucleotide sequences from the internal transcribed spacer (ITS) for a particular species. (6).

The development of molecular techniques for taxonomic analysis of parasites has led to great improvements in their precise identification and factuality. These molecular techniques, in particular the use of molecular markers, have been used by researchers to identify and validate parasites. Although, improvements in marker detection systems in particular rDNA elements such as 18S, ITS, and 28S used in parasites, among others, monogeneans have allowed great progress to be made in recent years in various countries (8).

The ITS (Internal Transcribed Spacer) region is widely used as a molecular marker for taxonomy and phylogenetic analysis(9). The advantages of ITS over other non-coding regions are: (1) Multiple copies of rRNA genes; (2) Can be used for PCR amplification with multiple universal primers for different organisms; (3) The size of the fragment is not long to facilitate sequencing; and (4) high variation at the generic and species level due to the presence of nucleotide polymorphisms or insertions/deletions in sequences (10,11).So this journal was compiled to review the application of the ITS region as a molecular marker for the identification of parasites in aquaculture.

2. What is ITS?

One of the phylogenetic studies with a molecular approach is to use DNA characters from target organisms. DNA characters are known to be more consistent in establishing phylogenetic relationships than morphological characters (12). One of the molecular characteristics that can be used is the nuclear genome. The nuclear genome or nucleus that is often used to conclude a phylogenetic is ribosomal DNA called rDNA. rDNA is a region of the genome that codes for ribosomal RNA(13).

The highly conserved coding DNA region (18S, 28S rDNA) is a major evolutionary region that is often used for comparison at the species and related genera levels. Each rDNA unit in a set of chromosomes has a coding region of 18S, 5.8S, and 28S flanking ITS1 and ITS2 (14).The 18S rDNA gene, along with the two ITS regions and the 5.8S rDNA gene, has a total length of 2600 bp, apart from the 28S rDNA gene which has a length of 3300 bp. Molecular markers that can be used for taxonomic and phylogenetic studies at the species level for Zingiber are molecular markers ITS (15).

ITS in the 18S-28S nuclear rDNA region is the main focus for use in phylogenetic reconstruction. This is because the ITS region has a high level of variation compared to other regions in the small subunit and large subunit rDNA(14). Baldwin(16)added, thatthe ITS region is small, which is approximately 700 base pairs, and has many copies in the core genome. This character causes the ITS region to be easily isolated, amplified, and analyzed. Provides extensive phylogenetic information in prokaryotes and eukaryotes, collectively known as ribosomal DNA (rDNA), consisting of coding regions (18S, 5.8S, and 28S) and non-coding regions 2 internal transcription spacers (ITS-1, ITS-2) and one non-transcriptional spacer (NTS)(17).




Figure 1. Ribosomal subunits and internal transcribed spacer region(18)

The rDNA units can be seen in Figure 1. In the section between the 18S region and the 5.8S region, there are several hundred base pairs of DNA called ITS1 and between the 5.8S and 28S regions, there are ITS2 regions. The ITS region has advantages compared to other target molecular regions, including having a high level of sensitivity because it has about 100 replicates in the genome. The ITS region also has a high rate of evolution and is present in all eukaryotic rDNA genes (19).

3. Molecular Identification of Fish Parasites with ITS

Diseases of fish are one of the most common problems in aquaculture and export-import businesses. Diseases do not only attack humans but also attack livestock and plants, as well as fish that live in water. attack diseases caused by parasites, fungi, viruses, and those caused by some kind of chemical substance in the form of pollution so that the fish's organs are damaged and disease arises (20).

Parasitism in fish is an ecological coexistence between two organisms in close contact. Some parasites form a balance of host-parasite relationships to become encysted or dormant until conditions are favorable for reproduction (21). Fish encounter many disease agents, including parasites, in their environment. Parasites are mechanically and functionally harmful to living things and also have an exploitative effect on preying, eating host nutrients, and interfering with respiration. This parasite reduces demand in the market and causes huge economic losses, weakens and

even kills fish by taking advantage of their food(22).

Some of the parasites that are often found attacking fish in aquaculture are Platyhelminthes of the Trematoda class, both from the Monogenea and Digenea subclasses. Platyhelminthes is one of the largest which includes a group of parasites commonly found on the gills, skin and fins of fish, lower aquatic invertebrates and some other amphibian vertebrates (8).

Digenean genera that affect mainly humans and livestock are best studied to date based on immunology, epidemiology and other medical fields (23). Recently, efforts have been made toward understanding the origin, diversification and phylogenetic interrelationships. The other genus *Hysterothylacium* has a cosmopolitan and circumpolar distribution and is mainly found in temperate and cold waters (24).

Another case study is the discovery of the parasite *Anisakis* sp. the prevalence of infection was generally higher in bigger fish than in smaller ones (25). Other parasites are *Diplostomum* spp. sampled in Spain: *D. spathaceum* with only one haplotype shared with adult isolates from central and northern Europe. No specific geographic pattern of the distribution of the novel haplotypes was found. This first molecular exploration of the diversity of *Diplostomum* spp. in southern Europe indicates much lower species richness compared with the northern regions of Europe(26)

In addition, the application contains parasites found in morpho-molecular characterization of *Gyrodactylus* parasites of

No.	Host	Origin	Parasites	ITS Primer	References
				CATT) and NC2(5-TTAGTTTCTTTTCTCCGCT-3),	
5.	Threadfin bream <i>Nemipterus japonicus</i>	China	Ascaridoid nematodes	A (forward: 5'-GTC GAA TTC GTA GGT GAA CCT GCG GAA GGA TCA-3') and B (reverse: 5'-GCC GGA TCC GAA TCC TGG TTA GTT TCT TTT CCT-3')	(32)
6.	<i>Katsuwonus pelamis</i> , <i>Auxis thazard</i>	Southern Makassar Strait, Indonesia	Anisakis type I	NC5 (5'-GTAGGTGAACCTGCGGAAGGAT CATT-3') and NC2 (5'-TTAGTTTCTTTTCTCCGCT-3')	(25)
7.	Pseudochondrostoma willkommii and <i>Salmo trutta</i>	Different regions in Spain	Diplostomum	D1 (forward: 5'-AGG AAT TCC TGG TAA GTG CAA G-3') and D2 (reverse: 5'-CGT TAC TGA GGG AAT CCT GGT-3')	(26)
8.	Tilapia	Mexico and Kenya	Gyrodactylids	ITS1-fm (5'-TAG AGG AAG TAC AAG TCG -3') and ITS2-rm (5'-GCT YGA ATC GAG GTC AGG AC-3')	(33)
				internal primers ITS3A (5'-GAG CCG AGT GAT CCACC-3') and ITS4.5 (5'-CAT CGG TCT CTC GAACG-3')	
9.	<i>Lophius piscatorius</i> , <i>Zeus faber</i> , <i>Engraulis encrasicolus</i> , <i>Merluccius merluccius</i> , <i>Mullus barbatus</i> , <i>Trachurus trachurus</i> , <i>Trachinus draco</i> , <i>Trigla lyra</i> , <i>Conger conger</i> , <i>Phycis phycis</i> , and <i>Pagellus erythrinus</i> .	Southern Mediterranean Sea (Southern Italy)	<i>Hysterothylacium</i> sp.	NC5 primer (forward: 50-GTA GGT GAA CCT GCG GAA GGA TCA TT- 30) and the NC2 primer (reverse: 50-TTA GTT TCT TTT CCT CCG CT-30)	(24)
10.	<i>Salmo trutta</i>	North and Central European freshwater	Glochidia larvae	Primer sequences were ITS-1-F (for-ward): AGACTGGGTTGCGGAGGT; and ITS-1-R (reverse): CGAGTGATCCACCGCTTAGA.	(28)

Formatted: Indonesian (Indonesia)

Formatted: Indonesian (Indonesia)

Formatted: French (France)

Formatted: Indonesian (Indonesia)

4. Limitations of ITS in aquaculture

However, it is generally accepted that ITS copies within the genome can be highly homogeneous due to coordinated evolution (34,35). Importantly, there are observed a low level of ambiguity in sequence data. Nevertheless, the limitations mentioned here also apply to mitochondrial markers(36).

Inappropriate homogenization of rDNA repeats can be compared with heteroplasmy of different mitochondrial haplotypes in a single organism (37,38), and rDNA pseudogenes are the nuclear copies of mitochondrial genes(39). Therefore, one should be aware of this complication with either ribosomal or mitochondrial markers, although they are not common enough to seriously affect their broad applicability (38).

5. Conclusion and Prospect

ITS markers have many advantages including having multiple copies of rRNA genes; being suitable for PCR amplification with multiple universal primers for different organisms; average sequence length for sorting; and a high degree of variation at the generic and species levels due to frequent nucleotide polymorphisms or insertions/deletions in sequences. So it is effective to be applied to detect parasites in fish. Many studies have been carried out using ITS markers, including Rainbow trout (*Oncorhynchus mykiss*), Argentine sea bass, (*Acanthistius patachonicus*), and other samples from various places.

The ITS region is a highly polymorphic noncoding region with enough taxonomic units to segregate sequences at the species level. Although ribosomal ITS as a universal barcode marker is still hampered by several limitations, ITS remains a major option for parasite identification in aquaculture in future.

Reference

1. DJPB KKP. Report DJPB 2021. Marine and Fisheries Ministry 2021. 74 p.
2. Zimik P, Roy B. Molecular identification

Moreover, there are several nucleotide sequencedatabases in the world. GenBank is one of the most important and the most influential nucleotide sequencedatabase that can be accessed publicly. As GenBank is constructed by direct submissions from individual laboratories, some of the sequence data may be unreliable due to incorrect identification of the organism from where the submitted sequence originates(40).

However, as dedicated databases are built in various locations, initial morphological identification at the genus level must be completed before any gene can be applied. For this reason, ribosomal ITS as a universal barcode marker for fungi is still hampered by several limitations(41). Moreover, ITS remains the primary choice for identification..

of two cestodes species parasitizing freshwater fishes in India. J Parasit Dis [Internet]. 2019;43(1):59–65. Available from: <https://doi.org/10.1007/s12639-018-1058-z>

Formatted: French (France)

3. Karami AM, Marnis H, Korb R, Zuo S, Jaafar R, Duan Y, et al. Absence of zoonotic parasites in salmonid aquaculture in Denmark: Causes and consequences. Aquaculture [Internet]. 2022;549(November 2021):737793. Available from: <https://doi.org/10.1016/j.aquaculture.2021.737793>

Formatted: French (France)

4. Timi JT, Paoletti M, Cimmaruta R, Lanfranchi AL, Alarcos AJ, Garbin L, et al. Molecular identification, morphological characterization and new insights into the ecology of larval *Pseudoterranova cattan* in fishes from the Argentine coast with its differentiation from the Antarctic species, *P. decipiens* sp. E (Nematoda: Anisakidae). Vet Parasitol [Internet]. 2014;199(1–2):59–72. Available from: <http://dx.doi.org/10.1016/j.vetpar.2013.09.033>

5. Altinok I, Kurt I. Molecular Diagnosis of Fish Diseases: a Review. Turkish J Fish Aquat Sci. 2003;3:131–8.
6. Ahmed M, Singh MN, Bera AK,

- Bandyopadhyay S, Bhattacharya D. Molecular basis for identification of species/isolates of gastrointestinal nematode parasites. *Asian Pac J Trop Med.* 2011;4(8):589–93.
7. Guo N, Chen HX, Zhang LP, Zhang JY, Yang LY, Li L. Infection and molecular identification of ascaridoid nematodes from the important marine food fish Japanese threadfin bream *Nemipterus japonicus* (Bloch) (Perciformes: Nemipteridae) in China. *Infect Genet Evol* [Internet]. 2020;85(September):104562. Available from: <https://doi.org/10.1016/j.meegid.2020.104562>
 8. Chaudhary A, Verma C, Singh HS. Ribosomal DNA as molecular markers and their applications in the identification of fish parasites (Platyhelminthes: Monogenea) from India. *Asian Pacific J Trop Dis.* 2014;4(1):82–4.
 9. Porter C, Collins F. Species-diagnostic differences in a ribosomal DNA internal transcribed spacer from the sibling species *Anopheles freeborni* and *Anopheles hermsi* (Diptera:Culicidae). *Am J Trop Med Hyg.* 1991;45(2):271–9.
 10. Poczai P, Hyvönen J. Nuclear ribosomal spacer regions in plant phylogenetics: Problems and prospects. *Mol Biol Rep.* 2010;37(4):1897–912.
 11. Calonje M, Martín-Bravo S, Dobeš C, Gong W, Jordon-Thaden I, Kiefer C, et al. Non-coding nuclear DNA markers in phylogenetic reconstruction. *Plant Syst Evol.* 2009;282(3–4):257–80.
 12. Noer S. Identifikasi Bakteri secara Molekular Menggunakan 16S rRNA. *EduBiologia Biol Sci Educ J.* 2021;1(1):1.
 13. Srivastava AK, Schlessinger D. Structure and organization of ribosomal DNA. *Biochimie.* 1991;73(6):631–8.
 14. Soltis DE, Soltis PS. Choosing an Approach and an Appropriate Gene for Phylogenetic Analysis. *Mol Syst Plants II.* 1998;1–42.
 15. Kress WJ, Prince LM, Hahn WJ, Zimmer EA. Unraveling the evolutionary radiation of the families of the zingiberales using morphological and molecular evidence. *Syst Biol.* 2001;50(6):926–44.
 16. Baldwin B., Sanderson MJ, Porter JM, Wojciechowski MF, Campbell CS, Donoghue MJ. The ITS Region of Nuclear Ribosomal DNA: A Valuable Source of Evidence on Angiosperm Phylogeny. 1995. p. 247–77.
 17. Wei NWV, Wallace CC, Dai CF, Pillay KRM, Chen CA. Analyses of the ribosomal internal transcribed spacers (ITS) and the 5.8S gene indicate that extremely high rDNA heterogeneity is a unique feature in the scleractinian coral genus *Acropora* (Scleractinia; Acroporidae). *Zool Stud.* 2006;45(3):404–18.
 18. Choudhary K, Verma AK, Swaroop S, Agrawal N. A review on the molecular characterization of digenean parasites using molecular markers with special reference to ITS region. *Helminthol.* 2015;52(3):167–87.
 19. Jorgensen R, Synder C, Jones JDG. T-DNA is organized predominantly in inverted repeat structures in plants transformed with *Agrobacterium tumefaciens* C58 derivatives. *Mol Gen Genet MGG.* 1987;207:471–7.
 20. Arannilewa ST, Salawu SO, Sorungbe AA, Ola-Salawu BB. Effect of frozen period on the chemical, microbiological and sensory quality of frozen tilapia fish (*Sarotherodon galiaenus*). *African J Biotechnol.* 2005;4(8):852–5.
 21. Schäperclaus W. *Fish diseases.* 5th Ed. vol. 1. Rotterdam: A.A. Balkema; 1992. 545 p.
 22. Özcan M, Bozdoğan N. Molecular identification of *Neoechinorhynchus rutili* parasite diagnosed in some fish species caught in Menzelet dam lake in Kahramanmaraş province (Turkey). *Saudi J Biol Sci.* 2020;27(7):1717–21.
 23. Snyder SD. Phylogeny and paraphyly among tetrapod blood flukes (Digenea: Schistosomatidae and Spirorchidae). *Int J Parasitol.* 2004;34(12):1385–92.
 24. Costa A, Graci S, Cammilleri G, Buscemi

- MD, Collura R, Vella A, et al. Molecular Identification of *Hysterothylacium* spp. in Fishes from the Southern Mediterranean Sea (Southern Italy). *J Parasitol.* 2018;104(4):398–406.
25. Anshary H, Sriwulan, Freeman MA, Ogawa K. Occurrence and molecular identification of *Anisakis* Dujardin, 1845 from marine fish in southern Makassar Strait, Indonesia. *Korean J Parasitol.* 2014;52(1):9–19.
 26. Pérez-del-Olmo A, Georgieva S, Pula HJ, Kostadinova A. Molecular and morphological evidence for three species of *Diplostomum* (Digenea: Diplostomidae), parasites of fishes and fish-eating birds in Spain. *Parasit Vectors.* 2014;7(1):502.
 27. García-Vásquez A, Pinacho-Pinacho CD, Guzmán-Valdivieso I, Calixto-Rojas M, Rubio-Godoy M. Morpho-molecular characterization of *Gyrodactylus* parasites of farmed tilapia and their spillover to native fishes in Mexico. *Sci Rep* [Internet]. 2021;11(1):1–17. Available from: <https://doi.org/10.1038/s41598-021-93472-6>
 28. Zieritz A, Gum B, Kuehn R, Geist J. Identifying freshwater mussels (Unionoida) and parasitic glochidia larvae from host fish gills: A molecular key to the North and Central European species. *Ecol Evol.* 2012;2(4):740–50.
 29. Gweon HS, Oliver A, Taylor J, Booth T, Gibbs M, Read DS, et al. PIPITS: An automated pipeline for analyses of fungal internal transcribed spacer sequences from the Illumina sequencing platform. *Methods Ecol Evol.* 2015;6(8):973–80.
 30. Karami AM, Marnis H, Korbut R, Zuo S, Jaafar R, Duan Y, et al. Absence of zoonotic parasites in salmonid aquaculture in Denmark: Causes and consequences. *Aquaculture.* 2022;549(November 2021):737793.
 31. Timi JT, Paoletti M, Cimmaruta R, Lanfranchi AL, Alarcos AJ, Garbin L, et al. Molecular identification, morphological characterization and new insights into the ecology of larval *Pseudoterranova cattani* in fishes from the Argentine coast with its differentiation from the Antarctic species, *P. decipiens* sp. E (Nematoda: Anisakidae). *Vet Parasitol.* 2014;199(1–2):59–72.
 32. Guo N, Chen HX, Zhang LP, Zhang JY, Yang LY, Li L. Infection and molecular identification of ascaridoid nematodes from the important marine food fish Japanese threadfin bream *Nemipterus japonicus* (Bloch) (Perciformes: Nemipteridae) in China. *Infect Genet Evol.* 2020;85(September):104562.
 33. García-Vásquez A, Pinacho-Pinacho CD, Guzmán-Valdivieso I, Calixto-Rojas M, Rubio-Godoy M. Morpho-molecular characterization of *Gyrodactylus* parasites of farmed tilapia and their spillover to native fishes in Mexico. *Sci Rep.* 2021;11(1):1–17.
 34. Kovarik A, Pires JC, Leitch AR, Lim KY, Sherwood AM, Matyasek R, et al. Rapid concerted evolution of nuclear ribosomal DNA in two tragopogon allopolyploids of recent and recurrent origin. *Genetics.* 2005;169(2):931–44.
 35. Koch MA, Dobeš C, Mitchell-Olds T. Multiple hybrid formation in natural populations: Concerted evolution of the internal transcribed spacer of nuclear ribosomal DNA (ITS) in North American *Arabis divaricarpa* (brassicaceae). *Mol Biol Evol.* 2003;20(3):338–50.
 36. Hu ZM, Guiry MD, Duan DL. Using the ribosomal internal transcribed spacer (ITS) as a complement marker for species identification of red macroalgae. *Hydrobiologia.* 2009;635(1):279–87.
 37. Kmiec B, Woloszynska M, Janska H. Heteroplasmy as a common state of mitochondrial genetic information in plants and animals. *Curr Genet.* 2006;50(3):149–59.
 38. Sonnenberg R, Nolte A, Tautz D. An evaluation of LSU rDNA D1-D2 sequences for their use in species identification. *Front Zool.* 2007;4:1–12.
 39. Pons J, Vogler AP. Complex pattern of coalescence and fast evolution of a mitochondrial rRNA pseudogene in a

Formatted: French (France)

recent radiation of tiger beetles. *Mol Biol Evol.* 2005;22(4):991–1000.

40. Fajarningsih ND. Internal Transcribed Spacer (ITS) as Dna Barcoding to Identify Fungal Species: a Review. *Squalen Bull Mar Fish Postharvest Biotechnol.* 2016;11(2):37.
41. Schoch CL, Robbertse B, Robert V, Vu D, Cardinali G, Irinyi L, et al. Finding needles in haystacks: Linking scientific names, reference specimens and molecular data for Fungi. *Database.* 2014;2014:1–21.