

## Original Research Article

### **Rice bran Stabilization by Solid-State Fermentation: Effect on Chemical Composition, Functional Properties, and Antioxidant Activity**

#### **Abstract**

Rice bran is the main by-product of rice grain processing. It is produced in large quantities worldwide and it contains a high amount of valuable nutrients and bioactive compounds with significant health-related properties. Despite that, its application in the food industry is still scarce because of its sensitivity to oxidation processes, instability, and poor technological suitability. Thus, this study aimed to investigate the chemical composition, functional properties, and antioxidant activity of rice bran fermented by lactic acid bacteria (*Lactobacillus plantarum*), and *Saccharomyces cerevisiae* as active dry yeast in the solid state. Bioactive compounds (free, bound, and total phenolic acids), antioxidant activity, water-extractable arabinoxylans and functional properties were evaluated. Similarly, phytic acid degradation was determined in fermented and non-fermented rice bran. After fermentation processes, protein, fiber, and ash contents in modified bran were raised compared to non-modified bran. Total, free phenolic content, and antioxidant activity increased after solid-state fermentation. Phytic acid content decrement of 40.01% and water-extractable arabinoxylans increment four times were observed in fermented bran. As well as, protein solubility, water holding capacity, water-solubility, oil holding capacity of fermented rice bran were enhanced. Thus, the solid-state fermentation is an efficient technique and can be applied to improve the nutritional and functional properties of rice bran, as well as utilized as ingredients in food applications.

**Comment [A1]:** Kindly do the Font size 12 point.

**Comment [A2]:** What is the use of this line?

**Keywords:** Rice bran, Lactic acid bacteria, *Saccharomyces cerevisiae*, functional properties, water-extractable arabinoxylans, phytic acid, phenolics, and antioxidant activity

**Comment [A3]:** Use keywords 5-6.

#### **1. Introduction**

World rice production in 2020 was approximately 500 million metric tons [1]. Before consumption, almost all rice is milled and as a result, rice bran is obtained

**Comment [A4]:** Cite the reference as per guidelines with proper font colour.

in large amounts around 63 million tons annually as a by-product of rice milling [2]. The bran consists of the pericarp, aleurone layer, embryo, and some endosperm, constituting about 10% of the unprocessed rice [3]. Rice bran is rich in protein, essential nutrients like minerals, vitamins, essential fatty acids, dietary fiber, and antioxidants, such as tocopherols and oryzanols [4]. It has been reported that rice bran is mostly comprised of protein (13.2%–18.6%), ash (9.2%–13.9%), lipid (9.5%–22.9%), starch (16.1%–26.7%) and dietary fiber (20.5%–33.3%) [5, 6]. The antioxidants, including anthocyanin and various phenolic acids, were found in rice bran, which has a strong ability to inhibit the formation of or to scavenge the reactive cell-damaging free radicals [7, 8]. Further, the antioxidant capacity of phenolic compounds is associated with health-promoting functions, such as obesity prevention, and anti-cardiovascular disease, as well as anti-inflammatory and anti-cancer effects [9]. In rice grains, phenolic compounds and phenolic acids exist in three forms: soluble free, soluble conjugated, and insoluble bound states [10]. Therapeutic effects were closely related to the nutrient components such as phenolic, dietary fiber, and antioxidants of rice bran [11]. Furthermore, each year 90% of the rice bran produced in the world is utilized cheaply as a feed stock for cattle and poultry and the remainder is used for the extraction of rice bran oil [12, 13], rice bran has very short shelf life owing to high-fat content and potent enzyme lipase which degrades the oil, making bran rancid and inedible.

Phytic acid is an anti-nutritional factor. It is decreasing the dietary bioavailability of nutrients minerals such as  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$ ,  $\text{Fe}^{++}$ , and  $\text{Zn}^{++}$  [14]. In rice kernel, phytic acid is mainly distributed in the outer layers, pericarp and germ [15]. Arabinoxylans are the main components of cell walls in cereal grains. Also, the content of arabinoxylan in rice is highest among all polysaccharides except starch [16], and they exist in both insoluble water and soluble states. The total

arabinoxylan content of rice bran is 4.84%–5.11%, while water-soluble arabinoxylan is 0.35%–0.77% [17]. Numerous investigations have suggested that the insoluble form has negative results on the rheological characteristics of dough [18].

To improve utilization of rice bran, development of feasible stabilization methods and the value-added processes of the nutrients and active compositions are necessary. Many investigations have concentrated on modifying features of rice bran. Numerous investigations have highlighted that fermentation of rice bran with lactic acid bacteria (LAB) or yeast, is a useful pre-treatment process to enhance technological, nutritional, and sensory properties of bran-containing products [11, 19], and to reduce anti-nutritive factors like phytic acid as well as enhancing mineral bioavailability [20]. Similarly, bran fermentation improves bioavailability and the contents of various functional compounds such as total free phenols, and soluble fiber as well as improving functional properties [8, 21]. Therefore, the aim of the current study was to modify autoclaved rice bran by solid-state fermentation with yeast and LAB. Also, study these effects on gross chemical composition, functional properties, phenolics, phytic acid contents, and antioxidant activity of rice bran.

## **2. Material and Methods**

### **2.1 Raw materials**

Commercial rice bran was obtained from Kafr El-Sheikh Mills company (Kafr El-Sheikh, Egypt), *Saccharomyces cerevisiae* as active dry yeast (Commercial baker's yeast with high sugar tolerance) was obtained from local market, and *Lactobacillus plantarum* ATCC 14917 was purchased from Microbiological Resources center (Cairo MIRCEN) Ain Shams University, Cairo, Egypt.

## 2.2 Inoculant preparation

*Lactobacillus* bacteria strains (*Lactobacillus plantarum*) were activated by inoculating in sterile MRS broth (9 ml) and incubation at 37 °C for 24 h. The cells were separated from the broth by centrifuging and re-suspended in sterile saline solution (9 ml) with final concentration 10<sup>8</sup> CFU/mL [22]. Active dry yeast was used directly without incubation.

## 2.3. Solid-state fermentation

The fermentation processes of autoclaved Rice bran (121°C at 15 min), was carried out as follows:

**Group □ treated by *L. plantarum*:** The autoclaved water (2:1) was added to rice bran, the activated culture of probiotic bacteria (10<sup>8</sup> CFU/mL) was added to autoclaved bran by a concentration of 1% and incubated at 37 °C for 24 and 48h.

Comment [A5]: What this symbol indicates?

**Group □ treated by yeast:** The autoclaved water (2:1) was added to rice bran and 1.25% active dry yeast was mixed well and fermented for 24 and 48h.

Comment [A6]: Refere proper bullets and no. for highlighted points.

**Group □ treated by LPY:** The autoclaved water (2:1) was added to rice bran, the activated culture of probiotic bacteria (10<sup>8</sup> CFU/mL) was added to autoclaved bran by a concentration of 1% and 1.25% active dry yeast was mixed well and fermented for 24 h and 48h. Rice bran autoclaved and non-autoclaved samples without starters were incubated at 37 °C and used as control. After fermentation all samples were air-dried, then milled by a laboratory mill (Braun, Germany) to pass through a 60-mesh sieve and stored at -20°C until analyses.

## 2.4. Gross chemical composition of Rice bran:

Comment [A7]: Article must be single spaced only. Do this correction in remain all.

The chemical composition of fermented and non-fermented rice brans including protein, fat, fiber, and ash contents were determined according to official methods as described in [23] Carbohydrate was calculated by difference.

## 2.5 Phytic acid:

The phytic acid was determined in terms of its phosphorous content, using the method described by [24].

## 2.6 phenolic compounds:

Total phenolic content was extracted from rice bran samples using methods described by the method of [25], while free phenolic compounds were extracted according to [26]. Total and free phenolic content determined by the Folin–Ciocalteu’s method according to [27]. The results were expressed as milligrams of gallic acid equivalents (GAE) per100 gram of bran sample on dry weight basis. Bound phenolic compounds were calculated by subtract free phenolics from total phenolics.

## 2.7 Determination of antioxidant activity

Samples were extracted using methods described by [28]. The 2,2-Diphenyl-1-picrylhydrazyl (DPPH) assay was carried out according to the method described by [29] with some modifications. The stock reagent solution ( $10^{-3}$  Mol) was prepared by dissolving 22 mg of (DPPH) in 50 ml of methanol and stored at 20°C until use. The working solution ( $6 \times 10^{-5}$  Mol) was prepared by mixing 6 mL of stock solution with 100 mL of methanol to obtain an absorbance value of  $0.8 \pm 0.02$  at 515 nm, as measured using a spectrophotometer (6505 UV/Vis, Jenway LTD., Felsted, Dunmow, UK). Extract solution of tested samples (0.1 ml) were vortexed for 30 s with 3.9 ml of DPPH solution and left to react for 30 min, after which the absorbance was measured at 515 nm and recorded. A control with no added extract was also analyzed. Scavenging activity was calculated as follows:

$$\text{DPPH radical scavenging activity (\%)} = \left[ \frac{(\text{Ab control} - \text{Ab sample})}{\text{Ab control}} \right] \times 100$$

Where Ab is the absorbance at 515 nm.

**Comment [A8]:** Cite the reference in uniform way only.

**Comment [A9]:** Space between two words must be only one. Do this correction in remain all.

## 2.8 Water extractable arabinoxylans (WEAX)

The WEAX was determined following previous study [30] by extracting 0.4 g of bran samples with 20 mL of distilled water at room temperature. The extracts were centrifuged at 5000 rpm for 10 min. 100 µL of supernatant, 100 µL of distilled water and 2 mL of freshly prepared reaction solution (1 g phloroglucinol in 5 mL anhydrous ethanol, 2 mL chlorohydric acid, 110 mL acetic acid, 1 mL 17.5 g/L glucose solution) were pipetted into stoppered glass tube. The tubes were vigorously boiling in water bath for 25 min and then cooled in flowing water immediately. The absorbance was measured at 552 nm and 510 nm successively. D- (+)-Xylose was used as standard. Calculate the content of WEAX by subtraction of the absorbance at 510 nm from 552 nm and comparison of the results with a standard curve and conversion of D-(+)-xylose to pentosan with a scaling factor (0.88).

**Comment [A10]:** Write proper abbreviations for specific units. Do this for remain all.

## 2.9 pH value:

The pH of the samples was determined according to approved methods 02-52 and 02-31 [31]. one gram of fermented and non-fermented Rice brans was first diluted and homogenize in 10 ml of distilled water then the pH of suspension was measured by a pH meter (Hanna Instruments, Padova, Italy).

## 2.10. Functional properties

Water-holding capacity and water solubility was achieved according to the method of [32]. Oil Holding Capacity was determined according to the method described by [33]. The protein solubility was achieved according to the method of [34].

**Comment [A11]:** Refer only one uniform proper style of referencin as per guidelines.

## 2.11 Statistical analysis

**Table 1. Gross chemical composition of fermented and non-fermented rice brans (D.W bases)**

---

Data were analyzed statistically by one-way analysis of variance (ANOVA) using SPSS software (IBM SPSS Statistics, version 22). When the difference between the samples was statistically significant ( $p < 0.05$ ), the Duncan test was used to determine the differences among the mean.

### **3. Results and Discussion**

#### **3.1 Chemical composition**

The results of gross chemical composition of fermented and non-fermented rice bran were illustrated in Table 1. It could be observed that the rice bran raw (RBR) contained 17.81% protein, 8.74% crude fiber, 6.99% ash, 17.27% fat, and 49.19% carbohydrate. The gross chemical composition of RBR in this work are agreement with that reported by [5, 6]. The protein, crude fiber and fat of rice bran was increased after treatment by fermentation compared to autoclaved sample RBA, the high protein content recorded for sample RBYP48h (21.13 %) followed by sample RBP48h (19.56 %) and RBP24h (19.44%). It's been reported that, yeasts and lactic acid bacteria have some proteinases and peptidases [35], while crude fiber increased from 7.02% to 8.64% in RBA and RBY48h samples, respectively, similar results were reported by [36]. In the other hand, there are slight differences in ash content of rice bran after fermentation process compared to non-treated sample (RBR). Small decreased differences were observed in carbohydrate contents of modified rice bran comparison with unmodified sample.

Samples	protein	fiber	Ash	fat	Carbohydrate*
<b>RBR</b>	17.81±1.20c	8.74±0.93a	6.99±0.19bc	17.27±0.73c	49.19±5.69b
<b>RBA</b>	16.98±1.09d	7.02±0.19c	6.86±0.86 <sup>bcd</sup>	17.16±0.84c	51.98±3.44a
<b>RBYP24h</b>	18.40±0.60c	8.43±0.42ab	7.13±0.03b	17.80±0.13 <sup>bc</sup>	48.24±0.92bc
<b>RBYP48h</b>	18.48±0.52c	8.64±0.32a	7.68±0.35a	18.10±0.10 <sup>abc</sup>	47.10±0.65c
<b>RBP24h</b>	19.44±0.46b	7.47±0.21c	6.54±0.05d	18.13±0.37 <sup>abc</sup>	48.43±5.09bc
<b>RBP48h</b>	19.56±0.44b	8.29±0.12ab	6.69±0.14cd	18.47±0.53ab	46.99±5.52c
<b>RBYP24h</b>	19.35±0.35b	7.74±0.23bc	6.67±0.27cd	18.87±0.27a	47.37±4.78c
<b>RBYP48h</b>	21.13±0.42a	8.45±0.17ab	7.14±0.03b	18.97±0.77a	44.31±5.74d

\*Carbohydrate by deference

Values are the mean of triplicate determinations with standard division.

The different letters at the column means significant differences at ( $p \leq 0.05$ ) and the same letters means no significant differences.

**RBR** raw bran; **RBA** autoclaved rice bran; **RBYP24h** fermented autoclaved bran with yeast for 24h; **RBYP48h** fermented autoclaved bran with yeast for 48h; **RBP24h** fermented autoclaved bran with lactic acid bacteria for 24h; **RBP48h** fermented autoclaved bran with lactic acid bacteria for 48h; **RBYP24h** fermented autoclaved bran with lactic acid bacteria and yeast for 24h; **RBYP48h** fermented autoclaved bran with lactic acid bacteria and yeast for 48h

**Comment [A12]:** Do the page set up properly. As table name indicates in one page and tale reflects in one page.

### 3.2. Phenolic compounds and antioxidant activity of fermented and non-fermented rice bran.

The effect of fermentation on phenolic compounds and antioxidant activity of rice bran are presented in Table 2. The results indicated that the free phenolic, total phenolic content and antioxidant activity of rice bran under different fermentation conditions by yeast, LAB, and Yeast combined with LAB were increased after fermentation compared to untreated sample. The highest free content of modified rice bran was observed in RBP48h (721.03 mg GAE /100g), RBYP48h (672.39 mg GAE/100g), and RBYP48h (665.41 mg GAE /100g), while the lowest 586.34 mg GAE /100g was for RBA. While the high of total phenolic content was recorded as follows: 852.33 mg GAE/100g > 806.64 mg GAE/100g > 781.83 mg GAE/100g > for RBYP48h, RBP48h, RBYP24h, respectively. The increase in the total and free phenolic content can be explained by the ability of yeast and LAB to degrade

organic materials due to their highly efficient enzymatic system. Similar results have been previously reported by [37, 38].

From the same table, it was observed that arabinoxylans after fermentation, the content of water-extractable arabinoxylans in rice bran improved significantly ( $P<0.05$ ) Table 2, increasing from 2.75 (WBR) to 4.72 mg/g for (WBY48h) sample. This enhancement means that fermentation, could be caused the transition from water unextractable arabinoxylans to water-extractable arabinoxylans partly. Water extractable arabinoxylans in bran increased after fermentation, and the highest content was observed in RBPY48h (4.72 mg/g), RBP48h (3.16 mg/g), RBY48h (3.07 mg/g), and RBPY24h (2.91 mg/g), while the lowest 2.12 mg/g was for RBP24h, Similar results have been previously reported by [39, 40].

**Table 2. Phenolic compounds, antioxidant activity and water extractable arabinoxylans of fermented and non-fermented rice bran**

Samples	Total phenolic content mg/100g			Antioxidant activity (%)	Water extractable arabinoxylans mg/g
	Total	Bound	Free		
<b>RBR</b>	771.22±1.20c	115.22±0.63b	656.00±1.00d	17.77±0.17g	1.13±0.03g
<b>RBA</b>	691.58±2.83d	105.24±0.92b	586.34±0.66g	19.73±0.70cd	1.18±0.01g
<b>RBY24h</b>	762.31±6.69c	158.21±6.60a	604.10±5.14e	20.56±0.56b	2.75±0.02e
<b>RBY48h</b>	753.38±4.62c	80.99±8.00b	672.39±3.39b	21.75±0.05a	3.07±0.02c
<b>RBP24h</b>	770.13±9.87bc	172.80±7.90a	597.33±3.17f	18.89±0.09ef	2.12±0.03f
<b>RBP48h</b>	806.64±5.36b	85.61±1.38b	721.03±4.03a	20.26±0.07bc	3.16±0.09b
<b>RBYP24h</b>	781.83±7.04bc	188.96±3.40a	592.87±2.76fg	18.43±0.06f	2.91±0.05d
<b>RBYP48h</b>	852.33±3.67a	186.92±2.10a	665.41±3.59c	19.41±0.41de	4.72±0.06a

Values are the mean of triplicate determinations with standard division.

The different letters at the column means significant differences at ( $p\leq 0.05$ ) and the same letters means no significant differences.

**RBR** raw bran; **RBA** autoclaved rice bran; **RBY24h** fermented autoclaved bran with yeast for 24h; **RBY48h** fermented autoclaved bran with yeast for 48h; **RBP24h** fermented autoclaved bran with lactic acid bacteria for 24h; **RBP48h** fermented autoclaved bran with lactic acid bacteria for 48h; **RBYP24h** fermented autoclaved bran with lactic acid bacteria and yeast for 24h; **RBYP48h** fermented autoclaved bran with lactic acid bacteria and yeast for 48h

### 3.3. Phytate phosphorus, phytic acid degradation, phytic acid and pH values of fermented and non-fermented rice bran

Table 3 shows the effect of fermentation on phytate phosphorus, phytic acid degradation, phytate content, and pH values of rice bran. It was observed that the highest values of phytate phosphorus (0.85%) and phytate content (3.02%) was recorded in raw rice bran. In contrast, the phytate phosphorus of fermented rice bran decreased significantly ( $P < 0.05$ ) from 0.75 to 0.51%, while phytate content decreased from 2.66 to 1.81%, and the highest phytate phosphorus 0.75% and phytic acid 2.66% was documented in RBY24h, followed by RBP24h (0.73 and 2.59%), RBY48h (0.66 and 2.34%), and RBYP24h (0.62 and 2.20%), the lowest 0.51% and 1.81% were recorded in RBP48h, these result in same line reported **by** [20]. Fermentation has been shown to reduce the phytic acid content of grains because of the action of phytases produced by yeast or lactic acid bacteria [41]. In the same trend, the high value of phytic acid degradation were recorded as follows: 40.01% > 37.75% > 27.15 > 22.52% for RBP48h, RBYP48h, RBYP24h, and RBY48h respectively, whereas phytic acid degradation was zero in RBC sample (Table 3). Phytic acid degradation may be due to phytase, and phosphatase enzymes present in microorganisms which can hydrolyze the phytic acid expressed in *Lactobacillus rhamnosus*1473, as previously reported in strains of the same species [42, 43]. The pH reduction caused by the production of carbon dioxide and organic acids by yeast increases phytase activity thereby, increasing phytic acid solubility. Results regarding the pH values, was observed that a considerable decrease in pH values was (from  $6.51 \pm 0.01$  to  $5.15 \pm 0.06$ ).

**Comment [A13]:** P is always italicized and capitalized.

**Comment [A14]:** Who is the inventor? It seems the sentence is not completed. Also make the uniformity while writing sentences some words are bold some are not.

Do this correction in remain all.

**Table 3. Phytate phosphorus, phytic acid degradation, phytic acid and pH values of fermented and non-fermented rice bran**

Samples	Phytate phosphorus %	Phytic acid degradation %	Phytic acid %	PH
<b>RBR</b>	0.85±0.12 <sup>a</sup>	00	3.02±0.44 <sup>a</sup>	6.51±0.01 <sup>a</sup>
<b>RBA</b>	0.81±0.25 <sup>ab</sup>	4.97	2.87±0.09 <sup>ab</sup>	6.50±0.08 <sup>a</sup>
<b>RBYP24h</b>	0.75±0.01 <sup>b</sup>	11.92	2.66±0.21 <sup>b</sup>	6.20±0.04 <sup>b</sup>
<b>RBYP48h</b>	0.66±0.06 <sup>cd</sup>	22.52	2.34±0.16 <sup>cd</sup>	5.78±0.05 <sup>e</sup>
<b>RBP24h</b>	0.73±0.04 <sup>bc</sup>	14.24	2.59±0.31 <sup>bc</sup>	6.11±0.03 <sup>c</sup>
<b>RBP48h</b>	0.51±0.01 <sup>e</sup>	40.01	1.81±0.37 <sup>e</sup>	5.15±0.06 <sup>g</sup>
<b>RBYP24h</b>	0.62±0.08 <sup>d</sup>	27.15	2.20±0.19 <sup>d</sup>	5.98±0.07 <sup>d</sup>
<b>RBYP48h</b>	0.53±0.09 <sup>e</sup>	37.75	1.88±0.33 <sup>e</sup>	5.54±0.10 <sup>f</sup>

Values are the mean of triplicate determinations with standard division.

The different letters at the column means significant differences at ( $p \leq 0.05$ ) and the same letters means no significant differences

**RBR** raw bran; **RBA** autoclaved rice bran; **RBYP24h** fermented autoclaved bran with yeast for 24h; **RBYP48h** fermented autoclaved bran with yeast for 48h; **RBP24h** fermented autoclaved bran with lactic acid bacteria for 24h; **RBP48h** fermented autoclaved bran with lactic acid bacteria for 48h; **RBYP24h** fermented autoclaved bran with lactic acid bacteria and yeast for 24h; **RBYP48h** fermented autoclaved bran with lactic acid bacteria and yeast for 48h

### 3.4. Functional properties of fermented and non-fermented rice bran.

The results in Table 4 indicate the functional properties of fermented and non-fermented rice bran. It was observed that the functional properties of rice bran including the water holding capacity, water-solubility, oil holding capacity, and protein solubility of fermented rice bran were improved and showed higher values than that observed for non-fermented sample and showed significant variations at ( $p \leq 0.05$ ). The highest WHC was recorded in RBYP48h (223.26%), followed by RBP48h (219.12%), RBYP48h (218.95%) and RBYP24h (218.61%). while the lowest was in RBR (213.33%), Similar results have been previously reported by [44]. These results demonstrate that all fermentation can be improving the WHC of rice bran. This increase in WHC may be due to an increment of soluble dietary fiber because it can absorb more water than insoluble dietary fiber. It has been reported that fermentation by lactic acid bacteria or yeast can enhance the content of soluble

dietary fiber of bran [45, 46]. In addition, the water solubility (WS) of raw rice bran was 12.14%, and it was significantly increased up to 17.50% after fermentation ( $p < 0.05$ ). This increase in WS of modified rice bran could be due to the increase of soluble dietary fiber.

After-fermentation the OHC of rice bran was enhanced obviously. The mean values showed higher oil holding capacity for RBYP48h (187.76%), followed by RBP24h (184.10%), RBPY24h (183.59%) and RBY48h (183.45%), while the lowest 176.50% was found in RBA. Similar results of wheat bran have been previously reported by [46]. Regarding protein solubility results (Table 4), the protein solubility of raw rice bran (RBR) was 10.85%, while autoclaving and fermentation processes increased the soluble protein of rice bran significantly ( $p < 0.05$ ). The highest values were recorded as follows: 16.04% > 14.93% > 14.13% > 13.83% for RBYP48h, RBP48h, RBY48h, and RBPY24h, respectively. These results are in line with those reported by [47]. These increasing in soluble protein, maybe due to degradation of proteins in rice brans after fermentation process of rice bran treatment by yeast or lactic acid bacteria. It was reported that Lactic fermentation enhanced indirectly protein hydrolysis via endogenous proteases [48].

**Table 4. Functional properties of and non-fermented rice brans (D.W bases)**

Samples	Water holding capacity (%)	water solubility	Oil holding capacity (%)	soluble protein as % of total sample protein
---------	----------------------------	------------------	--------------------------	--

<b>RBR</b>	213.33±1.67 <sup>c</sup>	12.14±0.36 <sup>d</sup>	176.88±1.88 <sup>d</sup>	10.85±0.50 <sup>e</sup>
<b>RBA</b>	215.18±1.82 <sup>d</sup>	11.89±0.53 <sup>d</sup>	176.50±1.55 <sup>d</sup>	12.54±0.40 <sup>d</sup>
<b>RBYP24h</b>	217.83±0.45 <sup>bc</sup>	12.66±0.33 <sup>d</sup>	182.33±1.67 <sup>c</sup>	13.74±0.87 <sup>c</sup>
<b>RBYP48h</b>	218.95±1.63 <sup>b</sup>	13.98±0.02 <sup>c</sup>	183.45±0.96 <sup>b</sup>	14.13±0.26 <sup>c</sup>
<b>RBP24h</b>	216.87±2.04 <sup>c</sup>	14.23±0.77 <sup>bc</sup>	181.37±1.41 <sup>c</sup>	12.85±0.87 <sup>d</sup>
<b>RBP48h</b>	219.12±0.97 <sup>b</sup>	15.00±0.50 <sup>b</sup>	184.10±1.80 <sup>b</sup>	14.93±0.07 <sup>b</sup>
<b>RBYP24h</b>	218.61±1.47 <sup>b</sup>	14.11±0.65 <sup>c</sup>	183.59±1.24 <sup>b</sup>	13.83±0.17 <sup>c</sup>
<b>RBYP48h</b>	223.26±0.75 <sup>a</sup>	17.50±0.47 <sup>a</sup>	187.76±2.10 <sup>a</sup>	16.04±0.46 <sup>a</sup>

Values are the mean of triplicate determinations with standard deviation.

The different letters at the column means significant differences at ( $p \leq 0.05$ ) and the same letters means no significant differences.

**RBR** raw bran; **RBA** autoclaved rice bran; **RBYP24h** fermented autoclaved bran with yeast for 24h; **RBYP48h** fermented autoclaved bran with yeast for 48h; **RBP24h** fermented autoclaved bran with lactic acid bacteria for 24h; **RBP48h** fermented autoclaved bran with lactic acid bacteria for 48h; **RBYP24h** fermented autoclaved bran with lactic acid bacteria and yeast for 24h; **RBYP48h** fermented autoclaved bran with lactic acid bacteria and yeast for 48h

#### 4. Conclusion

Our data demonstrated that solid-state fermentation is an efficient technique for improving rice bran's nutritional and functional characteristics. After fermentation, protein, fiber, ash contents as well as phenolic compounds, and antioxidant activity increased, phytic acid content reduced partly, while water-extractable arabinoxylans and protein solubility increased significantly ( $p < 0.05$ ). In addition, the functional properties of fermented rice bran including water holding capacity, water-solubility, and oil holding capacity were improved. Solid-state fermentation enhances the bioavailability of rice bran as well as increases its use as a functional ingredient in the food industry.

#### 5 -References

1. USDA (2021). World agricultural supply and demand estimates. In US Department of Agriculture
2. Webber, A., Hettiarachchy, N.S., Webber, D.M., Sivarrooban, T., Horax, R. Heat-stabilized defatted rice bran (HDRB) as an alternative growth medium for *Saccharomyces cerevisiae*. *Journal of Food and Nutrition*. 2014;1: 1–6.
3. Gul, K., et al., Rice bran: Nutritional values and its emerging potential for development of functional food—A review *Bioactive Carbohydrates and Dietary Fibre*. 2015; **6** (1): 24-30.

**Comment [A15]:** Do the proper alignment for all main headings.

4. Moongngarm, A., N. Daomukda, and S. Khumpika. Chemical compositions, phytochemicals, and antioxidant capacity of rice bran, rice bran layer, and rice germ. *Apcbee Procedia*. 2012; **2**: 73-79.
5. Casas, G.A., et al., Arabinoxylan is the main polysaccharide in fiber from rice coproducts, and increased concentration of fiber decreases *in vitro* digestibility of dry matter. *Animal Feed Science and Technology*. 2019; 247: 255-261.
6. Dang, T.T. and T. Vasanthan. Modification of rice bran dietary fiber concentrates using enzyme and extrusion cooking. *Food Hydrocolloids*. 2019; **89**: 773-782.
7. Nam, S.H., et al., Antioxidative activities of bran extracts from twenty one pigmented rice cultivars. *Food Chemistry*. 2006; **94** (4): 613-620.
8. Shin, H.-Y., et al. Solid-state fermentation of black rice bran with *Aspergillus awamori* and *Aspergillus oryzae*: Effects on phenolic acid composition and antioxidant activity of bran extracts. *Food chemistry*. 2019; 272: 235-241.
9. Min, B., A.M. McClung, and M.H. Chen. Phytochemicals and antioxidant capacities in rice brans of different color. *Journal of Food Science*. 2011; **76**(1): 117-126.
10. Shao, Y., et al., Phenolic acids, anthocyanins, and antioxidant capacity in rice (*Oryza sativa* L.) grains at four stages of development after flowering. *Food Chemistry*. 2014; **143**: 90-96.
11. Kuda, T., New Functional Properties of Fermented Rice Bran in Food Processing and Inflammatory Bowel Disease Model Mice, in *Dietary Interventions in Gastrointestinal Diseases*. 2019, Elsevier. . 197-206.
12. Schramm, R., et al. Fractionation of the rice bran layer and quantification of vitamin E, oryzanol, protein, and rice bran saccharide. *Journal of Biological Engineering*. 2007; **1**(1): 1-9.
13. Zullaikah, S., E. Melwita, and Y. Ju. Isolation of oryzanol from crude rice bran oil. *Food Chemistry*. 2009; 100(1): 299-302.
14. Bloot, A.P.M., et al. A review of phytic acid sources, obtention, and applications. *Food Reviews International*. 2021; 1-20.
15. Zhang, H.W. and X.L. Bai. Optimization of extraction conditions for phytic acid from rice bran using response surface methodology and its antioxidant effects. *Journal of food science and technology*. 2014; 51(2): 371-376.
16. Zhuang, X., Yin, T., Han, W., & Zhang, X. Nutritional ingredients and active compositions of defatted rice bran, in *Rice Bran and Rice Bran Oil*. AOCS Press .2019; 247-270.
17. Zhang, Z., C. Smith, and W. Li. Extraction and modification technology of arabinoxylans from cereal by-products: A critical review. *Food Research International*. 2014; **65**: 423-436.
18. Courtin, C. and J.A. Delcour. Arabinoxylans and endoxylanases in wheat flour bread-making. *Journal of cereal science*. 2002; **35**(3): 225-243.
19. Spaggiari, M., Dall'Asta, C., Galaverna, G. and M. D. del Castillo Bilbao. Rice bran by-product: From valorization strategies to nutritional perspectives. *Foods*. 2021; **10**(1): 85.
20. Ranjan, A., Sahu, N. P., Deo, A. D., and S. Kumar. Solid state fermentation of de-oiled rice bran: Effect on *in vitro* protein digestibility, fatty acid profile and anti-nutritional factors. *Food Research International*. 2019; **119**: 1-5.
21. Jia, M., et al. Structural characteristics and functional properties of soluble dietary fiber from defatted rice bran obtained through *Trichoderma viride* fermentation. *Food Hydrocolloids*. 2019; 94: 468-474.
22. Ujirohene, O.J., et al. Potent  $\alpha$ -amylase inhibitory activity of sprouted quinoa-based yoghurt beverages fermented with selected anti-diabetic strains of lactic acid bacteria. *RSC Advances*. 2019; 9(17): 9486-9493.

23. Association of Official Agriculture Chemists, W.D., Official methods of analysis Official Methods of Analysis 19th Ed.: Analytical Chemists International Arlington, Virginia, U.S.A; 2012.
24. Kent-Jones, D. and A.J. Amos. *Modern Cereal Chemistry*, Northern Pub. Liverpool, England. 1957.
25. Abdel-Gawad, A.S. Isolierung und charakterisierung von pentosanfraktionen aus verschiedenen weizen sorten. Ph.D.Dissertation Natural Science TU. Berlin, D 83 No.132/FB13. (In German). 1982.
26. Adom, K.K., Sorrells, M. E., and R. H. Liu. Phytochemical profiles and antioxidant activity of wheat varieties. *Journal of Agricultural and Food Chemistry*. 2003; **51**(26): 7825-7834.
27. Singleton, V.L., R. Orthofer, and R.M. Lamuela-Raventós, [14] Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent, in *Methods in enzymology*. 1999, Elsevier. p. 152-178.
28. Zieliński, H., Michalska, A., Ceglijska, A. and G. Lamparski. Antioxidant properties and sensory quality of traditional rye bread as affected by the incorporation of flour with different extraction rates in the formulation. *European Food Research and Technology*. 2008; **226**(4): 671-680.
29. Lee, S.-C., et al. Effect of far-infrared radiation on the antioxidant activity of rice hulls. *Journal of Agricultural and Food Chemistry*. 2003; **51**(15): 4400-4403.
30. Douglas, S. A rapid method for the determination of pentosans in wheat flour. *Food Chemistry*. 1981; **7**(2): 139-145.
31. Committee, A.A.o.C.C.A.M., *Approved methods of the American association of cereal chemists*. Vol. 1. 2000: Amer Assn of Cereal Chemists.
32. Singh, J. and N.. Singh. Studies on the morphological and rheological properties of granular cold water soluble corn and potato starches. *Food Hydrocolloids*. 2003; **17**(1): 63-72.
33. Sosulski, F., Humbert, E. S.; Bui, K. and J. O. Jones. Functional properties of rapeseed flours, concentrates and isolate. *Journal of Food Science*. 1976; **41**(6): 1349-1352.
34. Morr, C., et al. A collaborative study to develop a standardized food protein solubility procedure. *Journal of Food Science*. 1985; **50**(6): 1715-1718.
35. Hata, T., R. Hayashi, and E. Doi. Purification of Yeast Proteinases. *Agricultural and Biological Chemistry*, 2014. **31**(2): 150-169.
36. Oliveira, M.d.S., et al. Physico-chemical characterization of fermented rice bran biomass Caracterización físico-química de la biomasa del salvado de arroz fermentado. *CyTA–Journal of Food*. 2010; **8**(3): 229-236.
37. Verni, M., C.G. Rizzello, and R. Coda. Fermentation biotechnology applied to cereal industry by-products: Nutritional and functional insights. *Frontiers in nutrition*. 2019; **6**: 42.
38. Wang, M., et al. Impact of *Lactobacillus plantarum* 423 fermentation on the antioxidant activity and flavor properties of rice bran and wheat bran. *Food chemistry*. 2020; **330**: 127156.
39. Kim, D., and G.D. Han. Technologies, High hydrostatic pressure treatment combined with enzymes increases the extractability and bioactivity of fermented rice bran. *Innovative Food Science and Emerging Technologies*. 2012. **16**: 191-197.
40. Aktaş, K. and N. Akin. Influence of rice bran and corn bran addition on the selected properties of tarhana, a fermented cereal based food product. *LWT*; 2020. **129**: 109574.
41. Mukhopadhyay, N. and A.K.Ray. Improvement of quality of sal (*Shorea robusta*) seed meal protein with supplemental amino acids in feeds for rohu, *Labeo rohita* (Hamiltonian), fingerlings. *Acta Ichthyologica et Piscatoria*. 1999; **29**(1): p. 25-39.
42. Zamudio, M., A. Gonzalez, and J. Medina. *Lactobacillus plantarum* phytase activity is due to non-specific acid phosphatase. *Letters in Applied Microbiology*. 2001; **32**(3): 181-184.

43. Carrizo, S.L., et al., Ancestral Andean grain quinoa as source of lactic acid bacteria capable to degrade phytate and produce B-group vitamins. *Food Research International*. 2016. 89: p. 488-494.
44. Liu, Y., H., Yi, C., Quan, K., and B., Lin. Chemical composition, structure, physicochemical and functional properties of rice bran dietary fiber modified by cellulase treatment. *Food Chemistry*. 2021;342: 128352.
45. Zhao, H.-M., X.-N. Guo, and K.-X.J.F.C. Zhu. Impact of solid state fermentation on nutritional, physical and flavor properties of wheat bran. *Food Chemistry*. 2017; 217: 28-36.
46. Sorour, M. A., M. S., Hussein, M. A. Hassan, and E., Mahmoud. Impact of Solid State Fermentation on Chemical Composition, Functional Properties, and Antioxidant Activity of Wheat Bran. *ournal of Sohag Agriscience (JSAS)*. 2022. 7(1): 38-47.
47. Arte, E., et al., Impact of enzymatic and microbial bioprocessing on protein modification and nutritional properties of wheat bran. *Journal of Agricultural and Food Chemistry*. 2015. 63(39): 8685-8693.
48. Dallagnol, A.M., et al., Fermentation of quinoa and wheat slurries by *Lactobacillus plantarum* CRL 778: proteolytic activity. *Applied Microbiology Biotechnology*. 2013; 97(7): 3129-3140.

**Comment [A16]:** References are must be in uniform way.

**Comment [A17]:** See the font style as per guideline.