

# PREVALENCE OF MALARIA AND ASSOCIATED FACTORS AMONG PREGNANT WOMEN ATTENDING UNIOSUN TEACHING HOSPITAL OSOGBO OSUN STATE, NIGERIA.

## ABSTRACT

Pregnancy-related malaria is still a public health issue, as it contributes to a high infant mortality rate, low birth weight, and maternal mortality. Over 80% of the world's malaria cases were found in sub-Saharan Africa and India. More than 500,000 cases were reported in Nigeria, Madagascar, and the Democratic Republic of Congo. Pregnant women in sub-Saharan Africa, particularly in Nigeria, are most at risk of malaria infection each year, with about 50 million pregnancies at risk. Therefore, the objective of this study was to determine malaria prevalence and associated risk factors among pregnant women in Osun State, Nigeria.

The study was retrospective of five years between January 2017- December 2021. The medical records of 277 pregnant women who tested positive for malaria were retrieved to examine the contributing factors and malaria-related complications. Data were analyzed using the SPSS statistical software package, version 23. The Chi-square test was used to determine statistical significance. Data were presented in their simplest form and P - value < 0.05 was statistical significance.

The prevalence of malaria reported among pregnant women was found to be 98.9% caused by *Plasmodium falciparum*. While most pregnant women who tested positive for malaria parasite had no complications, low birth weight was the most common complication among pregnant women who were infected with malaria, Proportion of malaria in pregnant women during the gestational period was found, according to the findings - 1st Trimester 33.2%, 2nd Trimester 47.3%, and 3rd Trimester 19.5%. There was no significant association between the prevalence of malaria and Genotypes (P=0.498). It was discovered that the prevalence of malaria among pregnant women was high with *P. falciparum* being the most common malaria parasite found in the stud. Prenatal care was found to be a major factor in the early detection of signs and symptoms associated with pregnancy.

**Keywords:** Prevalence, Malaria, Pregnant Women, Associated Factors.

## Introduction

The Plasmodium genus of protists (a type of microorganism) causes malaria, a disease spread by mosquito bites that affect both humans and animals. Female Anopheles mosquitoes transmit malaria

which is also a life-threatening parasitic disease. Around the world, nearly half of the world's population is at risk of being infected with the malaria parasite (WHO, 2020). "Marsh fever" and "ague" were both previously used to describe the illness, which was originally known as "mala-aria" in Medieval Italian because of its connection to swamps and marshes. After being bitten by an infected female mosquito, protists (microorganisms) are released into the circulatory system and eventually develop and proliferate in the liver. As far as the health zone is concerned, malaria is still a major factor in maternal mortality and morbidity around the world and is regarded as contamination linked to poverty (Moya-Alvarez *et al.*, 2015; Ricci, 2016). According to the World Health Organisation (WHO, 2020), there were approximately 241 million new cases of malaria around the world. In addition, 95% of the malaria cases occurred in Africa, followed by Southeast Asia (3.2%), and the Eastern Mediterranean region (2%).

The tropics have long been a hotbed for malaria, which is spread by female Anopheles mosquitoes that are infected with parasites of the genus Plasmodium and reproduce in warm and humid conditions. *P. falciparum*, *P. vivax*, *P. malariae*, *P. knowlesi*, and *P. ovale* are just a few of the five distinct malaria parasite species that can infect humans. The estimated number of death due to malaria is said to be 627,000 in the world and the Africa Region has a high record of 95% malaria cases and 96% death due to malaria. (WHO, 2020). Even though malaria can be prevented, it remains the primary cause of maternal, in-utero, and infant morbidity and mortality. 1.2 billion of the 3.4 billion persons who are at high risk of contracting malaria are pregnant women. About 88% of cases and 90.0% of deaths were linked to malaria in the sub-Saharan Africa region, with more than 35.0% of these cases occurring in Nigeria and the DRC (Democratic Republic of Congo) area (National Multiple Indicator Cluster Survey [NMICS] & Federal Ministry of Health [FMOH], 2015; WHO, 2018, 2019). About 50 million times a year, pregnant women in sub-Saharan Africa, notably in Nigeria, are in danger of contracting malaria. Approximately 70% of pregnant women in Nigeria suffer from malaria, and 11% die as a result of the ailment (NMICS, 2015; Noland *et al.*, 2018). While pregnant women in areas with high rates of malaria transmission are particularly vulnerable to infection, there is no better time than during the first trimester to start building immunity against the disease (Eijk *et al.*, 2015).

*Plasmodium falciparum*, the parasite that causes malaria in pregnant women, is commonly overlooked as a public health issue by clinicians who treat such cases. More than 80% of malaria deaths occur in sub-Saharan Africa, according to the World Health Organisation 2013. Nigeria Malaria Indicator Report (2012) revealed that over 40% of all malaria-related deaths occur in Nigeria and the Democratic Republic of the Congo. Due to this knowledge, the two most significant nations in the world have teamed up to fight the spread of malaria. Malaria during pregnancy is one of the more common chronic infections affecting pregnant women, unborn children, and children under the age of five in Nigeria (National Bureau of Statistics [NBS] & FMOH, 2016). Over 800,000 women in Sub-Saharan Africa had low birth weight due to malaria, according to the World Health Organisation (WHO, 2020). The health of pregnant women and their unborn children has suffered as a result of this contamination. Other issues include pregnancy loss, extreme anemia, intrauterine growth restriction, fetal hypotrophy, and maternal hypoglycemia (Stephanie *et al.*, 2016).

The prevalence and risk factors of malaria among pregnant women in this part of the country need to be critically considered as a major public health issue with daily exposure to the malaria parasite. Therefore, this study's objective is to determine the prevalence of malaria and associated factors among pregnant women in Uniosun Teaching Hospital Osogbo Osun State Nigeria (277)

## **Material and Method**

### **Study design and area**

**Pregnant women were surveyed over five years to determine the prevalence of malaria and its associated factors (January 2017- December 2021). Pregnant women with malaria were counted at the hospital's medical records coding unit from January 2017 to December 2021. Records were checked out from the library to identify what factors were associated with each patient's coding unit number, as well as how malaria affected each patient's symptoms and overall health. This study was conducted at UNIOSUN Teaching Hospital Department of Obstetrics Clinic, Osogbo Osun State. The Hospital was established in 1997 as Ladoke Akintola University of Technology (LAUTECH) Teaching**

Hospital Complex, which is now acknowledged as Uniosun Teaching Hospital. Uniosun is a tertiary health organization located in Osogbo which is the capital of Osun State. Osogbo sits between Latitude 70° and 46° North and Longitude 40° and 36° East. The whole land area is 47km<sup>2</sup>. The town is positioned 88km northeast of Ibadan by road, 100km south of Ilorin via road, and 115km northwest of Akure. 48km from Ife, 32km from Ilesa, Iwo 46km, Ikire 48km and Ila-orangun 46km. The hospital serves the Osogbo Township where it is located and receives cases from other parts of the state. **Being a teaching hospital in the state and a place where most patients attend or are referred to for treatment, this made it suitable for the research setting, and also more well-detailed information will be obtained.**

#### Sample size determination and sampling technique

The **Kish, Leslie, 1965** formula was used to determine the sample size with a related prevalence of 82%, the confidence level was 95% with a margin of error of 5% which gives a total sample size of 277. Following approval from Uniosun Teaching Hospital, Osogbo Osun State Department of Obstetrics and Gynecology. Data on all pregnant women with malaria who visited an antenatal clinic during the study period were available through the coding system. The records of their cases were retrieved from the record library to gather the necessary information.

#### Data collection

The coding section of the hospital's medical records was used to gather the data. The patients who tested positive for malaria infection had their file numbers compiled, and the proforma was used to generate data for analysis.

#### Data analysis

Analysis was done using the Statistical Package for the Social Sciences (SPSS) version 23. The chi-square test was used to test for statistical significance for discrete and continuous data, respectively. The data was summarized and presented as tables and charts. Statistical significance was set at p-value < 0.05%.

## Results

### Frequency Distribution of Patients by Socio-demographic Variable

The highest age range of the patients was 26-32 years 113 (40.8%) and the lowest age range was 47-53 years 4 (1.4%). The highest marital status of the patients was the married group 264 (95.3%) while the lowest was the single group 13 (4.7%). The highest religious practice was Christianity 138 (49.8%) and the lowest religious practice was Traditional religion 2 (0.7%). The majority of the patient's ethnicity was Yoruba 246 (88.8%) while the lowest ethnicity was Hausa 16 (5.8%). The highest level of education of the patients was tertiary education 167 (60.3%) while the lowest level of education was primary 25 (9.0%). The occupational status of the majority of the patients was salaried earner 105 (37.9%) while the lowest occupational status was apprentice 8 (2.9%). The frequency of each socio-demographic variable is shown in (Table 1).

**Table 1: Frequency Distribution of Patients by Socio-demographic Variable**

Variable	Category	Frequency	Percentage
Age Group	19-25 years	62	22.4%
	26-32 years	113	40.8%
	33-39 years	69	24.9%
	40-46 years	29	10.5%
	47-53 years	4	1.4%
	Total		277

<b>Marital Status</b>	Single	13	4.7%
	Married	264	95.3%
	Total	277	100%
<b>Religion</b>	Christianity	138	49.8%
	Islam	137	49.5%
	Traditional	2	0.7%
	Total	277	100%
<b>Ethnicity</b>	Yoruba	246	88.8%
	Igbo	15	5.4%
	Hausa	16	5.8%
	Total	277	100%
<b>Highest level of education</b>	Primary	25	9.0%
	Secondary	84	30.3%
	Tertiary	167	60.3%
	None	1	0.4%
	Total	277	100%
<b>Occupational Status</b>	Salary earners	105	37.9%
	Trader	59	21.3%
	Unemployed	52	18.8%
	Artisan	53	19.1%
	Apprentice	8	2.9%
	Total	277	100%

---

### **Number of Pregnant Women Tested Positive for Malaria per Year**

In terms of the number of pregnant women with malaria over the five years, there was a high prevalence of malaria in the year 2018 70 (25.3%) compared to the year 2017 59 (21.3%), and there was also a decrease in the rate from the year 2019 58 (20.9%) compared to the year 2021 40 (14.4%) (Table 2).

**Table 2: Number of Pregnant Women Tested positive For Malaria per Year**

<b>Years</b>	<b>Number of Pregnant Women</b>	<b>Percentage</b>
2017	59	21.3%
2018	70	25.3%
2019	58	20.9%
2020	50	18.1%
2021	40	14.4%
<b>Total</b>	<b>277</b>	<b>100</b>

### **Frequency Distribution of Patients by Malaria Symptom**

The most prevalent symptoms of malaria among the patients were Fever (31.8%), 23.1% presented with Abdominal Pain, 22% had Spotting bleeding, 11.6% complained of Headache, 7.9% presented with Headache and Fever, 1.4% developed Dizziness, 1.4% had no symptoms, 0.4% presented Preterm Contraction and 0.4% had Vomiting (Table 3).

**Table 3: Frequency Distribution of Patients by Malaria Symptom**

<b>Variable</b>	<b>Category</b>	<b>Frequency</b>	<b>Percentage</b>
Symptoms of Malaria	Vagina bleeding	61	22%
	Fever	88	31.8%
	Abdominal pain	64	23.1%
	Bleeding	22	7.9%
	Headache	32	11.6%
	Preterm Contraction	1	0.4%
	Vomiting	1	0.4%
	Dizziness	4	1.4%
	No symptom	4	1.4%

**Frequency Distribution of Patients by Malaria complication**

The most prevalent complications of malaria among patients were low birth weight (16.2%), followed by premature delivery (15.2%), abortion (12.3%), miscarriage (8.3%), stillbirth (6.9%), and 41.2% had no complication. The frequency of each complication is shown in (Table 4).

**Table 4: Frequency Distribution of Patients by Malaria Complication**

<b>Variable</b>	<b>Category</b>	<b>Frequency</b>	<b>Cumulative Frequency</b>
<b>Complications of Malaria</b>	Stillbirth	19	19
	Premature delivery	42	61
	Low birth weight	45	106
	Miscarriage	23	127
	Abortion	34	163
	No Complication	114	277

### **Frequency Distribution of Patients by Gestational Age & Gravidity**

The majority of the pregnant women were in their 2nd Trimester (47.3%) and tested positive for malaria, 33.2% were in their 1st Trimester and 19.5% were in their 3rd Trimester. The highest Gravidity (35.7%) of pregnant women who tested positive for malaria were Secundigravida i.e. second pregnancy, 30.3% were Primigravida (First pregnancy), and 17.7% were Multigravida with the 4th pregnancy or more while 16.2%

were Multigravida with 3rd pregnancy. The frequency of each category is shown in (Table 5).

**Table 5: Frequency Distribution of Patients by Gestational Age & Gravidity**

<b>Variable</b>	<b>Category</b>	<b>Frequency</b>	<b>Percentage</b>
<b>Gestational age</b>	1st Trimester	92	33.2%
	2nd Trimester	131	47.3%
	3rd Trimester	54	19.5%
	Total	277	100%
<b>Gravidity</b>	Primigravida	84	30.3%
	Secundgravida	99	35.7%
	Multigravida (3rd)	45	16.2%
	Multigravida ( $\geq 4$ th)	49	17.7%
	Total	277	100%

**Frequency Distribution of Patients by ITN Utilisation & Antenatal Care Follow up**

A few of the patients (28.9%) utilized Insecticide-Treated Net while 71.1% did not. The majority (63.9%) of respondents had antenatal care follow-up while only 36.1% did not. The frequency is shown in (Table 6).

**Table 6: Frequency Distribution of Patients by ITN Utilisation & Antenatal Care Follow up**

Variable	Category	Frequency	Percentage
ITN Utilisation	Yes	80	28.9%
	No	197	71.1%
	Total	277	100%
Antenatal Care Follow up	Yes	177	63.9%
	No	100	36.1%
	Total	277	100%

**Chi-Square Test (Association between Prevalence of Malaria and Gestational Age, ITN Utilisation, Gravidity, Genotype, Symptoms, and Complications)**

1. There is a higher risk of *P. Falciparum* and *P. Ovale* among Patients in 2nd Gestational age while *P. Ovale* had a higher risk in 1st Trimester, however, this association was not statistically significant (p value=0.37)

2. There is a higher risk of *P. Falciparum*, *P. Vivax*, and *P. Ovale* among Patients without ITN, however, this association was not statistically significant (p value=0.65)
3. There is a higher risk of *Falciparum* Malaria, *Vivax*, and *Ovale* Malaria among Patients who are Secundigradiva, however, this association was not statistically significant (p value=0.76)
4. There is a higher risk of *P. Falciparum* and *P. Ovale* among Respondents with Genotype AA while *P. Ovale* had a higher risk among respondents with Genotype AS, however, there was no statistically significant association between Prevalence of Malaria and Gestational age (p value=0.49)
5. The risk of symptoms is very high with malaria prevalence (0.99). However, there is no statistically significant association between malaria prevalence and symptoms (p value=0.97)
6. The risk of complications is high with malaria prevalence (0.59). However, there is no statistically significant association between malaria prevalence and complications (p value=0.24)


The result shows that there is a higher risk of symptoms among patients with *P. Falciparum*, *P. vivax*, and *P. ovale* over five years, however, this association was not statistically significant (p value=0.97). There is also a higher risk of complications among patients with *P. falciparum* and *P. ovale*, however, this association was not statistically significant (p value=0.24). Each frequency of each associated factor is shown in (Table 7).

**Table 7: Chi-Square Test (Association between Prevalence of Malaria and Gestational Age, ITN Utilisation, Gravidity, Genotype, Symptoms, and Complications)**

Variable	Option	Falciparum	Vivax	Ovale	Total	X <sup>2</sup> Value	Df	P Value
Gestational Age	1 <sup>st</sup>	91	1	0	92	4.24	2	0.37
	2 <sup>nd</sup>	129	0	2	131			
	3 <sup>rd</sup>	54	0	0	54			
	Total	274	1	2	277			
ITN	Yes	79	0	1	80	0.84	2	0.65
	No	195	1	1	197			
	Total	274	1	2	277			
Gravidity	Primigravida	82	1	1	84	3.37	6	0.76
	Secundigravida	98	0	1	99			
	Multigravida	45	0	0	45			
	Multigravida $\geq 4$	49	0	0	49			
	Total	274	1	2	277			
Genotype	AC	29	0	1	30	5.36	6	0.49
	AA	143	0	1	144			
	AS	99	1	0	100			
	SC	3	0	0	3			
	Total	274	1	2	277			
Symptoms	Yes	270	1	2	273	0.04	2	0.97
	No	4	0	0	4			
	Total	274	1	2	277			
Complication	Yes	161	0	2	163	2.8	2	0.24
	No	113	1	0	114			
	Total	274	1	2	277			

**Figure 1: Frequency Distribution of Patients by Number of Antenatal Care Visits**


The majority (64.6%) of the patients visited Antenatal Clinic four times and more, 21.3% visited Antenatal Clinic three times, 10.1% visited Antenatal Clinic two times and 4% visited Antenatal Clinic only once (Figure 1).



**Figure 1: Frequency Distribution of Patients by Number of Antenatal Care Visits**

**Figure 2: Prevalence of Malaria among Pregnant Women by Plasmodium Species**

The prevalence of *falciparum* Malaria was 98.9%, the prevalence of *ovale* Malaria was 0.7% and the prevalence of *vivax* Malaria was 0.4% (Figure 2).




**Figure 2: Prevalence of Malaria among Pregnant Women by Plasmodium Species**

**Figure 3: Frequency Distribution of Patients by Genotype**

Note AC – Hemoglobin C trait, AA – Homozygous dominant. AS –Sickle-cell gene, SC – Hemoglobin (HB)

More than half (52%) of Patients had an AA Genotype, 36.1% had AS Genotype, 10.8% had AC and only 1.1% had the SC Genotype (Figure 3).



**Figure 3: Frequency Distribution of Patients by Genotype**

## **DISCUSSION**

Due to their weaker immune systems, pregnant women are more susceptible to malaria infection, which makes it a major problem for the country as a whole, particularly in Nigeria. More than half of all pregnant women in the study are married (95.3%), and the largest ethnic group is Yoruba (88.8%).

Most patients had *falciparum* malaria (98.9 %), followed by *ovale* malaria (0.7 %) and *vivax* malaria (0.4%). A higher prevalence of *falciparum* Malaria among pregnant women who visited antenatal care facilities at Uniosun Teaching Hospital, in response to the first question of the study could be observed. *Plasmodium falciparum* is an endemic disease in sub-Saharan Africa, and pregnant women's low immunity makes them vulnerable to *falciparum* malaria (Awosolu *et al.*, 2021).

A total of 52% of patients have the AA genotype, 36% of patients have the AS genotype, 10.8% of patients have the AC genotype, and 1.1% of patients have the SC genotype, according to the research. Pregnant women in the antenatal care clinic at Uniosun Teaching Hospital Osogbo have the AA Genotype, followed by AS Genotype, AC Genotype, and SC Genotype. This report agrees with that of Albiti and Nsiah (2014) in many ways. In comparison to AA, the prevalence of individuals with AS and AC is quite low. Stronger resistance to malaria may exist in people with AS, AC, or SC. Archer *et al.* (2018) findings support this conclusion. The level of susceptibility to malaria has been pronounced to be greater in individuals with AA when compared with those with AS, AC, and SC. Thus, the high frequency of AC and AS in malaria-endemic areas has been attributed to and reduction in malaria morbidity and mortality in malaria-endemic areas (Archer *et al.*, 2018). When compared to those with AS, AC, and SC, those with AA are more susceptible to malaria. A decrease in malaria-related morbidity and mortality has been linked to an increase in the prevalence of AC and AS, which are

common in malaria-endemic regions (Archer *et al.*, 2018). AC and AS take a defensive stance against malaria infection as a result of reduced cytoadhesion of infected red blood cells to microvasculature and impaired rosette formation as a result of the presence of parasite falciparum erythrocyte membrane protein 1 (PfEMP1) antigen on AC and AS (Fairhurst *et al.*, 2012).

Pregnant women at Uniosun Hospital Osogbo Teaching use ITNs at a very low rate in the past (28.9 %). Availability, affordability, and acceptance of ITN may be to blame for this low utilization. According to the National Demographic Health Survey (NPC Nigeria and ICF, 2019), 37.1% of the population was found to be in poor health. The findings are also in line with findings from a study in Edo State (Wagbatsoma *et al.*, 2010). Pregnant women attending an antenatal care clinic at Uniosun Teaching Hospital Osogbo ranged in gestational age from 47.3% in the 2nd trimester to 33.2% in the 1st trimester to 19.5% in the 3rd trimester. Pregnant women in the second trimester are more likely to volunteer for the study because they have had a reasonable amount of time to adjust to their pregnancy, unlike those in the first trimester, who may still be experiencing nausea and vomiting. Pregnant women in their third trimester may also be extremely obese at this time.

More than one-third (35.7%) of the women who took part in this study are classified as Secundigravida, while 30.3% are classified as Primigravida, and 17.7% are classified as Multigravida with four or more pregnancies. Meanwhile, 16.2% are classified as Multigravida with only their third pregnancy. This is expected, as most women prefer to have a small family of two or three, so it is common to see women who are pregnant for

the first or second time. It's odd, though, that there were more Multigravida with their fourth or subsequent pregnancies than there were with their third.

Patients had from one to four prenatal visits, with an average of 3.4 visits. According to the National Health and Nutrition Examination Survey [NDHS] 2018, 56.8% of pregnant women in the United States had at least four antenatal visits. The study also found that the majority of respondents (63.9 %) received Antenatal Care. More than 96% of pregnant women in Osun State were not receiving prenatal care (National Demographic Health Survey (NDHS), 2018).

Although there is a higher risk of *falciparum* malaria among Patients with Genotypes AA (AS), AC (AC), and SC (p- value=0.498), there is not a significant association. There is no significant correlation between pregnant women's genotypes and their likelihood of being infected with malaria in Osogbo, Nigeria, according to the first hypothesis.

ITN use is less common among those with the genotypes AA-SC than among those without, and thus no link can be drawn between the two variables. There appears to be no correlation between genotypes and the use of Insecticide Treated Nets (ITN) among pregnant women attending the antenatal care clinic at Uniosun Teaching Hospital Osogbo.

In the first, second, and third trimesters, women are more likely to contract *falciparum* malaria than *vivax* or *ovale* malaria, but there is no link (p value=0.373) between the prevalence of malaria and gestational age. In other words, this supports the hypothesis that pregnant women attending the antenatal care clinic at Uniosun Teaching Hospital Osogbo are not at increased risk of malaria because of their gestational age.

## **Conclusion**

Within five years, the prevalence of malaria among expectant mothers is reportedly low, making it simpler to further educate them. Due to a lack of use of insecticide-treated nets, which also contributed to greater complications, the prevalence of malaria symptoms among pregnant women is high. The majority of pregnant women go for antenatal care, which will also aid in the early identification of associated factors such as genotypes, gestational age, gravidity, and ITN use.

Education and emphasis on regular use of the insecticide-treated net by pregnant women during the antenatal clinic days. Also, awareness should be made of the importance of malaria in pregnancy prevention among pregnant women in the community through health education and health promotion. This will help in the future prevention of malaria among expected and pregnant women.

## **Ethical Approval**

Uniosun Teaching Hospital Moral Committee Osogbo provided ethical approval for the research. Adeleke University Ethical Committee, Ede, Osun State also provided ethical approval for the research with the reference number (AUERC/FBMS/IND/11). Knowledge obtained from this study has been processed and evaluated without revealing the participants' private details.

## **Acknowledgments**

Our acknowledgment goes to Adeleke University and Lautech ethical committee for giving us ethical approval. Also, my profound gratitude goes to Prof. J.E Atolagbe and Mr. Adeola Oshineye for their support, encouragement, advice, patience, understanding, and effort while supervising this work.

### **Conflict of Interest**

The authors declare no conflicts of interest in this work

### **Author Contributions**

Conceptualization : Odesanmi Oluwanifemi

Writing – original draft: Odesanmi Oluwanifemi

Writing – review & editing: Odesanmi Oluwanifemi, Prof. James Atolagbe, Adeola Oshineye

## References

- Albiti, A.H., Nsiah, K. (2014) Comparative hematological parameters of HbAA and HbAS genotype children infected with Plasmodium falciparum malaria in Yemen. *Hematology*, 19 (3): 169- 174.
- Archer, N.M., Petersen, N., Clark, M.A., Buckee C.O., Childs, L.M., Duraisingh, M.T. (2018). Resistance to Plasmodium falciparum in sickle cell erythrocytes is driven by oxygen-dependent- growth inhibition. *Proceedings of the National Academy of Sciences of the United States of America* 2018,115 (28): 7350-7355.
- Awosolu, O. B., Yahaya, Z. S., Farah Haziqah, M. T., Simon-Oke, I. A., Olanipekun, I. T., & Oniya, M. O. (2021). Epidemiology of falciparum malaria among residents of some rural and periurban communities in Ekiti State, Southwestern Nigeria. *Tropical biomedicine*, 38(1), 14–21. <https://doi.org/10.47665/tb.38.1.003>.

Eijk, A.M., Hill, J., Noor, A.M., Snow, R.W., Kuile, F.O. (2015). Prevalence of malaria infection in pregnant women compared with children for tracking malaria transmission in sub-Saharan Africa: a systematic review and meta-analysis. *Lancet Global Health*.3:e617–28.

Fairhurst, R.M., Bess, C.D., Krause, M.A. (2012). Abnormal PfEMP1/knob display on Plasmodium falciparum-infected erythrocytes containing hemoglobin variants: fresh insights into malaria pathogenesis and protection. *Microbes and Infection*, 14 (10): 851–862.

Federal Ministry of Health, Nepal Multiple Indicator Cluster Survey (2016). National Anti-malaria Treatment Policy.

Federal Ministry of Health. (2015). National guidelines & strategies for malaria Prevention & control during pregnancy. (A publication of the Federal Ministry of Health, Nigeria). Nigeria Maternal Health - malaria in pregnancy.

Kish, Leslie (1965). Survey Sampling. New York: John Wiley and Sons, Inc.

Moya-Alvarez, V., Abellana, R., & Cot, M., (2015). Pregnancy-associated malaria and malaria in Infants: An old problem with present consequences. *Malaria. Journal*. 13, 271.

National Population Commission (NPC) [Nigeria] and ICF. 2019. *Nigeria Demographic and Health Survey 2018*. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF. [natnets.org/attachments/article/65/Monitoring%20&%20Evaluation%20 Plan%202008-2013](http://natnets.org/attachments/article/65/Monitoring%20&%20Evaluation%20Plan%202008-2013).

Noland, G.S., Graves, P.M., Sallau, A., Eigege, A., Emukah, E., Patterson, A.E., Ajiji, J., Okorofor, I., Oji, O.U., & Umar, M. (2018). Malaria prevalence, anemia, and baseline

intervention coverage before mass net distributions in Abia and Plateau States, Nigeria. *BMC Infectious Diseases*.14:168. doi: 10.1186/1471-2334-14-168.

Ricci. F., (2016). Social implications of malaria and their relationships with poverty. *Mediterranean Journal Hematol Infect Dis*; 4(1).

Stephanie, D., Jenny, H., Jane, B., Peter, O., Doris M., Peter, O., & Meghna, D., (2016).Effectiveness of the delivery of interventions to prevent malaria in pregnancy in Kenya15 (221) DOI 10.1186/s12936-016-1261-2

Wagbatsoma, V. A., & Aigbe, E. E. (2010). ITN utilization among pregnant women attending ANC in Etsako West Lga, Edo State, Nigeria. *Nigerian journal of clinical practice*, 13(2), 44–148.

World Health Organization (2013). World Malaria Report. Geneva

WHO Global Malaria Programme (2020). World malaria report 2020. Geneva: World Health Organization

World Health Organization. (2018). World Malaria Report.

World Health Organization. World Malaria Report; WHO: Geneva, Switzerland, 2019; Available online: <https://www.who.int/malaria/publications/world-malaria-report-2019/en/>.

