

## Original Research Article

Assessment of heavy metal concentration in soil and plant and Evaluation of Bioconcentration factor at LOUMBILA market gardening perimeters, Burkina Faso

### Abstract

Agricultural soil quality deterioration resulting from an increase in the level of heavy metals is becoming more and more pronounced, thus raising the question of the safety status of human health and the environment. Determination of heavy metal concentration in soils and plants from Loumbila market gardening, and calculation of bioconcentration factor were undertaken. Heavy metals copper (Cu), nickel (Ni), lead (Pb), chromium (Cr), cadmium (Cd), and zinc (Zn) were measured using atomic absorption spectrophotometer (AAS) model AANALYST 200 from PERKIN ELMER.

In the soil where the pepper was grown, the concentration of chromium was 175 mg/kg, which is higher than the concentration limit which is 150 mg/kg. Also, the concentrations of lead in the soils of onion (118 mg/kg), green bean (118 mg/kg), carrot (178 mg/kg), and pepper (135 mg/kg) were above the limit which is 100 mg/kg. The concentrations of chromium in onion leaves (3.72 mg/kg), onions (4.65 mg/kg), lettuce (4.89 mg/kg), green beans (5.89 mg/kg), and bell peppers (3.56 mg/kg) were concentrations above the limit established by the FAO/WHO, which is 2.3 mg/Kg. The concentrations of lead in carrot, onion leaf, onion, lettuce, green bean, and bell pepper were above the limit proposed by FAO/WHO.

The bioconcentration factors show that the onion leaf (0.731), onion (0.929), lettuce (0.876), green bean (0.987), and bell pepper (0.858) are the plants that accumulate zinc and the carrot (0.524) accumulates nickel. The bioaccumulation of the metals in the vegetable from Loumbila market gardening decreased in the order of Onion > lettuce > green bean > bell pepper > onion leaves > carrot.

**Keywords:** Heavy Metal, Concentration, Bioconcentration factor, soil, plant

### I. Introduction

In countries witnessing rapid industrialization, one of the most pressing environmental issues challenging the global sustainable development agenda is environmental pollution due to toxic metals. Metal pollutants gain entry into the air, soil, and water bodies via natural and anthropogenic routes, including domestic and agricultural use, metals mining and smelting activities, and other industrial productions [1]. Rapid industrialization, inappropriate utilization, and disposal of toxic trace metal-containing wastes, excessive use of fertilizers and pesticides are the main causes of toxic trace metals in agricultural soils, which become hazardous to human and environmental health. The main heavy metals that contribute to environmental pollution, particularly in areas with high anthropogenic activities are lead (Pb), cadmium (Cd), nickel (Ni), cobalt (Co), iron (Fe), zinc (Zn), chromium (Cr). The capacities of heavy metal uptake and accumulation, mechanisms of metal concentration, exclusion, and

compartmentation vary among different plant species and also between various parts of plants [2, 3].

The problem of contaminated soil on the perimeters of market gardening is very worrying today in urban cities in emerging countries. Heavy metals such as lead, cadmium, copper, zinc, and mercury cannot be biodegraded and therefore persist in the environment for long periods. The accumulation of heavy metals in the environment can affect human and animal health [4]. Understanding the origin of heavy metal pollution, the phenomenon of accumulation in soils, and their possible interactions with soil constituents is a priority in many environmental studies. The accumulation of heavy metals in agricultural soils presents an increased risk of food pollution and a potential risk to human health [5].

The use of wastewater can increase crop productivity, but also increases the contamination of heavy metals (Pb, Ni, Cd, Cu, Zn, Mn, Cr...) in the plants. The plants (Amaranthus, Fenugreek, and Spinach) heavy metals concentrations depend on the soil concentration [6] [7]. The vegetable from the contaminated soil can accumulate high concentrations of heavy metals and cause some serious risks to human health [8]. As an example, [9] John et al., show that the vegetable and the soil from Kaduna city are polluted by Pb, Cd, and Cr [9]. The concentration of Pb in cabbage, (10.51), lettuce (10.19), green pepper (9.44), hot pepper (7.61), and ayoyo (9.05) from some parts of Accra were higher than the FAO/WHO maximum recommended limit of 0.30 mg/kg for Pb [10].

The concentration of cadmium (Cd), lead (Pb), and chromium (Cr) in vegetable species (*Telferia occidentalis*, *Talinum triangulare*, *Ocimum gratissimum*, *Celosia argentea*, and *Amaranthus Viridis*) cultivated in Lagos farmlands and floodplains, were above standard limit [11]. Heavy metals in plants can accumulate in the human body and cause diseases such as Colon, stomach, liver, lung, bladder, and kidney cancers [12] [13] [14].

Trace metal elements contained in wastewater are toxic and non-biodegradable substances that accumulate in the soil and, depending on biogeochemical conditions, can lead to soil degradation [15]. Studies show that even treated wastewater also contains heavy metals that can contaminate crops when used for irrigation [16].

Metal elements such as Fe, Zn, Mo, Cu, Co, and Cr are known to be essential elements in the living organism. However, a high concentration can be potentially toxic [17].

Pollution can be due on the one hand to the use of wastewater for irrigation and on the other hand to the proximity of the production sites of major urban traffic axes and industries that emit heavy metals [18]. The accumulation of heavy metals in soil can be due to the irrigation water and thereafter contaminate the vegetable and fruit. Heavy metal concentrations in the vegetable are different from those of the fruits revealing the difference between their capacities for accumulation [19].

The contamination of soils, plants, and water by heavy metals constitutes an important public health issue that requires more information on the various sources of pollution and the rate of contamination. Preliminary studies on heavy metal pollution in the market gardening areas of Ouagadougou and Loubila have revealed a deterioration in soil quality [20]. Concentration values of certain metals (Cr, Mn, Ni, and Hg) in irrigation water were found to be above standards. A high concentration of chromium was observed in tomatoes [21]. The alert resulting from this study on the potential risk of pollution of agricultural soils and transfer of pollutants into the food chain justifies the need to continue studies on the market gardening areas of Ouagadougou and Loubila on other plants.

The bioconcentration factor (BCF) for the heavy metals in the leafy vegetables decreased in the order of Cd (5.85) > Hg (3.83) > Fe (0.31) > Zn (0.11) > Cr (0) = Ni (0). The capacity of leafy vegetables to accumulate Fe, Cd, Zn, Cr, Ni, and Hg decreased in the order of *Amaranthus Spinosa* > *Corchorus solitorius* > *Brassica oleracea* > *Brassica rapa*. In the non-

leafy vegetables, the BCF decreased in the order of Hg (2.46) > Zn (0.12) > Fe (0.1) > Cd (0) = Cr (0) = Ni (0). The capacity of non-leafy vegetables to accumulate Fe, Cd, Zn, Cr, Ni, and Hg decreased in the order of *Daucus carota* > *Raphanus sativus* > *Allium cepa* ~ *capsicum annum*. The low BCF for Cr and Ni in all vegetable species may be due to the toxicity of the crops [22].

Based on the work carried out on heavy metals in plants in West Africa (Ghana, Nigeria, etc.) in general and in Burkina Faso in particular, this study focused on copper (Cu), chromium (Cr), zinc (Zn), lead (Pb), nickel (Ni), and cadmium (Cd) in the most consumed plants [10] [11] [20].

The objective of this study is to estimate the concentration of heavy metals (Cu, Cr, Zn, Pb, Ni, and Cd) in soil and vegetables at the Loumbila market gardening sites and evaluate the bioconcentration factor.

## II. Materials and methods

### II.1. Study Area

In this study, the vegetable samples were collected in different agricultural areas at the Loumbila market garden. Distance of 18 kilometers from Ouagadougou capital city of Burkina, Loumbila market garden is expanding around the dam. The dam is located at a longitude of 01°24'07.4 West and a latitude of 12°29'35.8 North with a water capacity of 42.2 million cubic meters. It is used by market gardeners to irrigate plants [23].  
Columbia's market gardening areas have a much-diversified production of vegetables, namely onion, tomato, okra, zucchini, African eggplant, eggplant, pepper, bell pepper, lettuce, cabbage, carrot, green bean, and potato. Market garden products from Loumbila can be found in most markets in the city of Ouagadougou or exported to neighboring countries. These reasons led to the choice of the Loumbila market gardening areas to carry out this study, by choosing the most consumed vegetables in the city of Ouagadougou.

### II.2. Samples and Sampling Techniques

The samples of soils and vegetables were collected from the market garden of Loumbila. The agricultural soil sample was collected at three or four different points on the diagonal profile of each site. The soils from the different points were mixed and kept in sterile plastic bags.

Sampling was carried out at regular intervals and over the entire plot for each type of sample. The different plants were sampled by separating the different plant tissues (roots, leaves, stems, and fruits) of each plant using a ceramic knife to avoid contamination. To measure an average concentration, at the plot level, the samples were mixed by family and constitute an aliquot. Plant samples for analysis of heavy metals were placed in plastic bags. The vegetables were washed up with tap water thoroughly to remove the attached dust particles, soil, unicellular algae, etc. Then they were washed with distilled water and finally with deionized water. The washed vegetables were dried at room temperature to remove surface water.

Carrot, onion leaves, onion, lettuce, green beam, and bell pepper were collected during the vegetable sampling.

### II.3. Laboratory Analysis

The soils and vegetables samples were analyzed for heavy metals such as chromium (Cr), zinc (Zn), lead (Pb), nickel (Ni), cadmium (Cd), and Mercury (Hg) using Atomic Absorption Spectrometer, model PERKIN ELMER AANALYST 200.

Mechanical preparation consisted of sieving the soil samples through a certified 200 mesh (75 microns) sieve. The test portion used is one gram. The soil samples were weighed using a PA214C balance from OHAUS PIONEER, with a precision of  $10^{-4}$  grams and a capacity of 200g. The mineralization consisted in mineralizing the samples weighed by aqua regia (2.5 mL of  $\text{HNO}_3$  + 7.5 mL of HCl) at a controlled temperature (water bath at  $90 \pm 50^\circ\text{C}$ ) for one hour. The acids used are of the analytical type. The solution obtained is consequently brought to 100 mL by way of demineralized water with a conductivity of less than  $2 \mu\text{S} / \text{cm}$ .

The vegetable sample (1 g) was weighed into a 100 ml volumetric flask and concentrated acids of 10.0 mL of concentrated sulfuric acid were added to each sample. The samples obtained after adding concentrated acid were gently on a hot plate, stirring occasionally until the powder completely dissolved in the solution (about 10 to 15 minutes). Then 10ml of distilled water was added and the whole was heated gently for a few minutes (5 to 10 mins). Finally, the solution was left to stand for it to settle well and the filtrate was taken for analysis.

The analysis results are valid based on the performance of the analysis methods used (limit of detection and quantification, data of duplicates and reference materials inserted in the chemical preparation of the samples, selectivity of the method, the robustness of the method, data for control solutions, data for chemical blanks, etc.).

### II.4. Bioconcentration factor

Bioconcentration factor (BCF) is defined as the ratio of the metal concentration in the plant to the metal concentration in the soil. BCF measures the heavy metal accumulation efficiency in plants. When the BCF values are greater than 1, indicates that the species is a hyperaccumulator of heavy metals [1-3; 22; 24-25].

$$BCF = \frac{\text{heavy metal concentration in plant}}{\text{heavy metal concentration in soil}}$$

## III. Results and discussions

### III.1. Concentration of heavy metals in agricultural soils

Table 1 presents the concentrations of Cu, Ni, Cr, Cd, Pb, and Zn in some market gardening soils at Loumbila. The soils were taken according to the plant produced. The soils for the

production of onions, green beans, lettuce, carrots, and bell peppers are the subject of this study.

Table 1: Concentration of some heavy metals in the market gardening soils of LOUMBILA

	Concentration in ppm (mg/kg)					
	Cu	Ni	Cr	Cd	Pb	Zn
Soil of onion and green beans	34	29	83	ND	118	79
Soil of Lettuce	35	24	69	ND	46	80
Soil of bellpepper	33	22	66	ND	82	82
Soil of carrot	34	21	62	ND	178	90
Soil of pepper	51	19	175	ND	135	107
Average	37.4	23	91	ND	111.8	87.6
Limits values	100	50	150	3	100	300

The copper concentrations in the studied soils vary from 33 mg/kg to 51 mg/kg with an average of 37.4 mg/kg. The copper concentrations obtained in this study were all below the limit value for copper in agricultural soils.

The studied soils present nickel concentrations which were between 19 mg/kg and 29 mg/kg with an average of 23 mg/kg. The limit value for nickel in soils is 50 mg/kg, which is higher than the values obtained in this study.

On the soil where the pepper was grown, the concentration of chromium was 175 mg/kg, which is higher than the concentration limit which is 150 mg/kg. The other soils had chromium concentrations below the limit. This high concentration of chromium in soil can lead to a high concentration of chromium in the pepper.

In all soil samples analyzed in this study, the cadmium concentration was below the detection limit.

The concentration of zinc in soils varies between 79 mg/kg and 107 mg/kg with an average of 87.6 mg/kg. The zinc concentrations in the studied soils were below the limit of 300 mg/kg.

The results of this study show that the concentrations of lead in the soils of onion, green bean, carrot, and pepper were above the limit which is 100 mg/kg. These high lead concentrations in soils can lead to high concentrations of lead in onions, green beans, carrots, and peppers. Lead concentrations in lettuce and pepper soils were below the limit.

The mean concentration of heavy metals in the agricultural soils samples from Loumbila market gardening decreased in the order Pb > Cr > Zn > Cu > Ni > Cd.

### III.2. Concentration of heavy metals in some plants

Table 2 presents the concentrations of Cu, Ni, Cr, Cd, Pb, and Zn in carrot, onion leaf, onion, lettuce, green bean, and bell pepper sampled from some soils of Loumbila.

Table 2: Concentrations of heavy metals in the studied plants

	Concentration in ppm (mg/kg)					
	Cu	Ni	Cr	Cd	Pb	Zn
Carrot	17.6	11	1.3	ND	27.02	42.95
Onion leaves	16.4	11.16	3.72	ND	10.48	57.72
Onion	19.81	10.05	4.65	ND	11.52	73.37
Lettuce	16.9	5.09	4.89	ND	15.44	70.1

<b>Green bean</b>	<b>16.06</b>	<b>8.58</b>	<b>5.89</b>	<b>ND</b>	<b>4.09</b>	<b>78</b>
<b>Bell pepper</b>	<b>19.65</b>	<b>7.01</b>	<b>3.56</b>	<b>ND</b>	<b>1.73</b>	<b>70.39</b>
<b>Average</b>	<b>17.74</b>	<b>8.82</b>	<b>4.00</b>	<b>ND</b>	<b>11.71</b>	<b>65.42</b>
<b>FAO/WHO safelimit</b>	<b>40</b>	<b>-</b>	<b>2.3</b>	<b>0.2</b>	<b>0.3</b>	<b>20</b>

The concentrations of copper in the studied plants vary from 16.06 mg/Kg to 19.81 mg/Kg. All measured concentrations in the studied plants were below the limit established by the FAO/WHO.

The concentrations of chromium in the studied plants vary from 1.3 mg/Kg to 5.89 mg/Kg. Onion leaves, onions, lettuce, green beans, and bell peppers have chromium concentrations above the limit established by the FAO/WHO, which is 2.3 mg/Kg. Carrots have a chromium concentration lower than the limit established by the FAO/WHO. Consumption in large quantities of onion leaf, onion, lettuce, green bean, and pepper from the study area can be the cause of diseases such as stomach cancer, and intestine, ... [14].

Concentrations of lead in carrot, onion leaf, onion, lettuce, green bean, and bell pepper were above the limit proposed by FAO/WHO. The high concentrations may be due to the high concentrations of lead observed in crop soils. The lettuce growing soil had a lead concentration below the limit whereas the lettuce leaves have a lead concentration above the FAO/WHO limit. This high concentration of lead in lettuce leaves may be because lettuce is accumulating lead. Consumption in large quantities of these products from the study area can be the cause of cardiovascular disease (hypertension), kidney damage [12], and liver disease [14].

The concentrations of zinc in carrot, onion leaf, onion, lettuce, green bean, and pepper were higher than the limit proposed by the FAO/WHO although the concentrations in the soils were lower. These high concentrations of zinc in carrots, onion leaves, onions, lettuce, green beans, and bell peppers show that the studied plants were hyper-accumulators of zinc. The mean concentration of heavy metals in the edible portions of the vegetables decreased in the order  $Zn > Cu > Pb > Ni > Cr > Cd$ .

### III.3. Soil-plant concentration

Figure 1 presents a histogram comparing the concentrations of copper, nickel, chromium, cadmium, lead, and zinc in the carrot and its cultivation soil.

Figure 1: Concentration of heavy metals in the carrot and its cultivation soil

The concentration of lead was the highest in carrot cultivation soil, while zinc is the metal with the highest concentration in carrot. Carrots accumulate less chromium than the other heavy metals studied.

Figure 2 presents a histogram comparing the concentrations of copper, nickel, chromium, cadmium, lead, and zinc in lettuce and its growing soil.

Figure 2: Concentration of heavy metals in lettuce and its growing soil

Zinc has a high concentration in lettuce and its growing soil. Chromium has a fairly high concentration in soil, while its concentration in lettuce is the lowest. This reflects a high absorption of zinc and low absorption of chromium by lettuce.

Figure 3 presents a histogram of the concentrations of heavy metals in green beans and their cultivation soil.

Figure 3: Concentration of heavymetals in green beans and theirgrowingsoil

Lead has a high concentration in greenbeansoil but a low concentration in **the green** bean. Zinc has a high concentration in green beans and isvery close to the soil concentration. This reflects a high absorption of zinc and low absorption of lead by the green bean.

Figure 4 presents a histogram of the concentrations of heavymetals in the pepper and its cultivation soil.

Figure 4: Concentration of heavymetals in bellpepper and itsgrowingsoil

Zinc has a high concentration in the pepper and its cultivation soil. Lead has a fairly high concentration in soil, whileits concentration in pepperis the lowest. This reflects a high absorption of zinc and **low** absorption of lead by the bellpepper.

Figure 5 presents a histogramcomparing the concentrations of copper, nickel, chromium, cadmium, lead, and zinc in the onion, theseleaves, and its cultivation soil.

Figure 5: Concentration of heavymetals in onion, onionleaves, and growingsoil

Onion-growingsoil has a high lead concentration and a low nickel concentration. Onionleaves have a high zinc concentration and a lowchromium concentration. Onion has a high zinc concentration and a lowchromium concentration. This reflects a high absorption of zinc and low absorption of chromium by the onion and theseleaves. The onionaccumulates more copper, chromium, lead, and zinc thantheseleaves. As for onionleaves, theyaccumulate more nickel.

The results of soil and plant concentrations show that zinc is the heavymetalthataccumulates the most in onionleaf, onion, lettuce, green bean, and pepper. Lead and chromiumaccumulateless in moststudied plants.

### III.4. Bioconcentration factor

Table 3: heavymetal Bioconcentration factor

	<b>Cu</b>	<b>Ni</b>	<b>Cr</b>	<b>Pb</b>	<b>Zn</b>
<b>Carrot</b>	<b>0.518</b>	<b>0.524</b>	<b>0.021</b>	<b>0.152</b>	<b>0.477</b>
<b>Onion</b>	<b>0.583</b>	<b>0.347</b>	<b>0.056</b>	<b>0.098</b>	<b>0.929</b>
<b>Onionleaves</b>	<b>0.482</b>	<b>0.385</b>	<b>0.045</b>	<b>0.089</b>	<b>0.731</b>
<b>Lettuce</b>	<b>0.483</b>	<b>0.212</b>	<b>0.071</b>	<b>0.336</b>	<b>0.876</b>
<b>Green Bean</b>	<b>0.472</b>	<b>0.296</b>	<b>0.071</b>	<b>0.035</b>	<b>0.987</b>
<b>Bell pepper</b>	<b>0.595</b>	<b>0.319</b>	<b>0.054</b>	<b>0.021</b>	<b>0.858</b>

Table 3presents the bioconcentration factors of copper, nickel, chromium, lead, and zinc in carrot, onionleaf, onion, lettuce, green bean, and bellpepper.

The carrot has a bioconcentrationfactor that varies from 0.021 to 0.524. The most accumulative metal in the carrotwas nickel.

The bioconcentration factor ranges from 0.056 to 0.929 in onions and ranges from 0.045 to 0.731 in onionleaves. The most accumulative metal in the onionand the leavesis zinc, which

has a bioconcentration factor close to one in the fruit. This reflects a strong transfer of zinc from the soil to the plant.

In lettuce, the bioconcentration factor ranges from 0.071 to 0.876. The most accumulating metal in lettuce was zinc.

Green beans have a bioconcentration factor that varies from 0.035 to 0.987. The most accumulative metal in the green bean is zinc, which has a bioconcentration factor roughly equal to one. This reflects a strong transfer of zinc from the soil to the green bean.

The bioconcentration factor varies from 0.021 to 0.858 in bell pepper. The most accumulative metal in bell pepper was zinc.

The bioconcentration factors show that the onion leaf, onion, lettuce, green bean, and pepper are the plants that accumulate zinc and the carrot accumulates nickel.

The bioaccumulation of the metals in the vegetable from Lumbini market gardening decreased in the order of Onion > lettuce > green bean > bell pepper > onion leaves > carrot.

#### IV. Conclusion

This study has revealed that the concentrations of lead in the soils of onion, green bean, carrot, and pepper are above the limit which is 100 mg/kg. These high lead concentrations in soils can lead to high lead concentrations in onions, green beans, carrots, and peppers. Lead concentrations in lettuce and pepper soils are below the limit.

The values of the bioaccumulation show that the onion leaf, onion, lettuce, green bean, and bell pepper were the plants that accumulate zinc and the carrot accumulates nickel.

The mean concentration of heavy metals in the agricultural soils samples from Lumbini market gardening decreased in the order Pb > Cr > Zn > Cu > Ni > Cd. The mean concentration of heavy metals in the edible portions of the vegetables decreased in the order Zn > Cu > Pb > Ni > Cr > Cd. For the individual vegetables, the bioaccumulation of the metals in the edible parts decreased in the order of Onion > lettuce > green bean > bell pepper > onion leaves > carrot.

#### References

- [1] **Kamal Usman, Hareb Al Jabri, Mohammed H. Abu-Dieyeh and Mohammed H. S. A. Alsafran, (2020). Comparative Assessment of Toxic Metals Bioaccumulation and the Mechanisms of Chromium (Cr) Tolerance and Uptake in *Calotropis procera*.** *Frontier Plant Science*, 19 June 2020 | <https://doi.org/10.3389/fpls.2020.00883>
- [2] Hira Amin, Basir Ahmed Arain, Taj Muhammad Jahangir, Muhammad Sadiq Abbasi & Farah Amin (2018) Accumulation and distribution of lead (Pb) in plant tissues of guar (*Cyamopsis tetragonoloba* L.) and sesame (*Sesamum indicum* L.): profitable phytoremediation with biofuel crops, *Geology, Ecology, and Landscapes*, 2:1, 51-60, DOI: 10.1080/24749508.2018.1452464 To link to this article: <https://doi.org/10.1080/24749508.2018.1452464>.
- [3] Anna Chrzan, (2016). Monitoring bioconcentration of potentially toxic trace elements in soils trophic chains. *Environ Earth Sci* (2016) 75:786, DOI 10.1007/s12665-016-5595-4.
- [4] Dung, H.T.M. (2012) Impacts des métaux lourds sur l'interaction plante/ver de terre/microflore tellurique. These, Présentée et soutenue publiquement le 22/12/2012 pour l'obtention du grade de Docteur de L'UNIVERSITE PARIS EST, 169 p.
- [5] Salano, E.M. (2013) Assessment of Heavy Metal Pollution in Soils and Water of Samburu County, Kenya. Thesis, Kenyatta University, Nairobi.
- [6] Puttaih, J.E.T. (2013) Assessment of Heavy Metals Uptake in Leafy Vegetables Grown on Long Term Wastewater Irrigated Soil across Vrishabhavathi River, Bangalore, Karnataka. *IOSR Journal of Environmental Science, Toxicology and Food Technology*, 7, 52-55. <https://www.iosrjournals.org/iosr-jestft/papers/vol7-issue6/10765255.pdf>
- [7] Yadav, A., Yadav, P.K. and Shukla, D.N. (2013) Investigation of Heavy Metal Status in Soil and Vegetables Grown in Urban Area of Allahabad, Uttar Pradesh, India. *International Journal of Scientific and Research Publications*, 3, 1-7.

- [8] Naser, H.M., Sultana, S., Mahmud, N.U., Gomes, R. and Noor, S. (2011) Heavy Metal Levels in Vegetables with Growth Stage and Plant Species Variations. *Bangladesh Journal of Agricultural Research*, 36, 563-574. <https://doi.org/10.3329/bjar.v36i4.11743>
- [9] Jacob, J.O. and Kakulu, S.E. (2012) Assessment of Heavy Metal Bioaccumulation in Spinach, Jute Mallow and Tomato in Farms within Kaduna Metropolis, Nigeria. *American Journal of Chemistry*, 2, 13-16. <https://doi.org/10.5923/j.chemistry.20120201.04>, <http://journal.sapub.org/chemistry>
- [10] Lente, I., Ofosu-Anim, J., Brimah, A.K. and Atiemo, S. (2014) Heavy Metal Pollution of Vegetable Crops Irrigated with Wastewater in Accra, Ghana. *West African Journal of Applied Ecology*, 22, 41-58.
- [11] Adedokun, A.H., Njoku, K.L., Akinola, M.O., Adesuyi, A.A. and Jolaoso, A.O. (2017) Heavy Metal Content and the Potential Health Risk Assessment of Some Leafy Vegetables Cultivated in Some Flood Plains and Farmlands in Lagos, Nigeria. *FUNAI Journal of Science & Technology*, 3, 30-47.
- [12] Huss, J. (2011) Les risques sanitaires des métaux lourds et d'autres métaux. Rapport 1 de la Commission des questions sociales, de la santé et de la famille, de l'Assemblée parlementaire de l'union européenne, Doc. 12613.
- [13] Said, B.M. (2011) Etude des impacts environnementaux des débris de démolition de la région de Boumerdes. These soutenue le pour l'obtention du grade de docteur de l'Université M'Hamed Bougara-Boumerdes, Année universitaire: 2010/2011, p 146.
- [14] Aloueimine, S.O. (2006) Methodologie de caracterisation des dechetsmenagers à Nouakchott (Mauritanie): Contribution à la gestion des Dechets et outils d'aide à la decision. These Presentee et soutenue publiquement le 13 Avril 2006 pour l'obtention du grade de Docteur de l'Université de Limoges, p 195.
- [15] Matech, F., Zaakour, F., Moustarhfer, K. and Chemsî, Z. (2014) Concentrations en elements traces metalliques dans les sols irrigues par les eaux useesversees dansL'OUED MERZEG (CASABLANCA-MAROC). *European Scientific Journal*, 10, 121-138.
- [16] Lokeshwari, H. and Chandrappa, G.T. (2006) Impact of Heavy Metal Contamination of Bellandur Lake on Soil and Cultivated Vegetation. *Current Science*, 91,622-627.
- [17] Majolagbe, A.O., Yusuf, K.A. and Duru, A.E. (2014) Trace Metals Characterisation in Environmental Media: A Case Study of Cement Production Area, Ewekoro, Southwest, Nigeria. *European Scientific Journal*, 3, 83-89. <https://doi.org/10.11648/j.ajep.20140302.17>
- [18] Tankari, D.-B.A., Guero, Y., Dan Lamso, N., Barage, M., Balla, A., Sterckeman, T., Ech Evarria, G. and Feidt, C. (2013) Evaluation des niveaux de contamination en elements traces metalliques de laitue et de chou cultives dans la vallee de GountiYenaa Niamey, Niger. *Journal of Applied Biosciences*, 67, 5326-5335. <https://doi.org/10.4314/jab.v67i0.95056>
- [19] Rapheal, O. and Adebayo, K.S. (2011) Assessment of Trace Heavy Metal Contaminations of Some Selected Vegetables Irrigated with Water from River Benue within Makurdi Metropolis, Benue State, Nigeria. *Advances in Applied Science Research*, 2, 590-601. <http://www.pelagiaresearchlibrary.com>
- [20] Bambara, L.T. (2016) Etude de la Pollution en metaux lourds dans les zones de maraichage au Burkina Faso: Cas de Ouagadougou et de Loubila. These unique de Doctorat, Université Ouaga I Professeur Joseph KI-ZERBO, p 190.
- [21] Bambara, L.T., Kabore, K., Derra, M., Zoungrana, M., Zougmore, F. andCisse, O. (2015) Assessment of Heavy Metals in Irrigation Water and Vegetables in Selected Farms at Loubila and Paspanga, Burkina Faso. *IOSR Journal of Environmental Science Toxicology and Food Technology*, 9, 99-103. <http://www.iosrjournals.org>
- [22] Andrews Obeng Affum, Shiloh Dede Osae, Edward EbowKwaansa-Ansah, Michael K. Miyittah, (2020). Quality assessment and potential health risk of heavy metals in leafy and non-leafy vegetables irrigated with groundwater and municipal-waste-dominated stream in the Western Region, Ghana. *Heliyon* 6 (2020) e05829, <https://doi.org/10.1016/j.heliyon.2020.e05829>
- [23] Prosper, D. (2007) Renforcement des capacites des exploitants maraichers: Cas du village de Nougou dans le departement de Loubila, memoirepresente a Institut International d'ingenierie en Eau et Environnement, 124.
- [24] Hema Diwan, Altaf Ahmad, and Muhammad Iqbal, 2010. Uptake-related parameters as indices of phytoremediation potential. *Biologia* 65/6: 1004—1011, 2010, Section Botany, DOI: 10.2478/s11756-010-0106-7.
- [25] LeônidasCarrijo Azevedo Melo, Evandro Barbosa da Silva, Luís Reynaldo FerracciúAlleoni, (2014). Transfer of cadmium and barium from soil to crops grown in tropical soils. *Soil Use and Management • Rev. Bras. Ciênc. Solo* 38 (6) • Dec 2014 • <https://doi.org/10.1590/S0100-06832014000600028>