

Original Research Article

Effect of foliar spray of polyamines and biostimulants on Rose (*Rosa hybrida* L.) underpolyhouse condition

ABSTRACT

Investigation was done to find out the effect of different polyamines viz., spermine and spermidine (5, 10, 20 mg/L) and biostimulants viz., 2 % Novel (banana pseudostem sap) and 2% cow urine as foliar spray on growth parameters in terms of qualitative and quantitative traits of rose var. Top Secret. The study was conducted for three years consecutively from 2015 to 2018. Plant growth in general increased with polyamines and biostimulants.

The vegetative attributes viz. plant height, number of branches and number of leaves were found to be significantly increased in rose plants given foliar spray of polyamines and biostimulants as observed during 1st year, 2nd year and, 3rd year as well as in pooled analysis. Comparing different treatments as per pooled analysis, significantly maximum plant height (118.27 cm) was found in plants treated with 20 mg/L spermidine while number of branches (6.91) were found significantly maximum in plants treated with 20 mg/L spermine. Maximum number of leaves (129.30) were observed with 20 mg/L spermine and were at par with 10 mg/L spermine and 20 mg/L spermidine.

Flowering parameters were also found to be improved by spraying of polyamines and biostimulants during each individual year as well as in pooled. Flower stem length (83.78 cm) was significantly maximum with treatment 20 mg/L spermine which was at par with 10 mg/L spermine and 2% novel. Flower stem diameter (7.96 mm) was significantly maximum with treatment 20 mg/L spermine at par with 10 mg/L spermine and followed by 2% Novel in the pooled data analysis. Bud diameter was significantly maximum (5.99 cm) with treatment of 10 mg/L spermine that was at par with 20 mg/L spermine, 20 mg/L spermidine and 2% novel while bud length (5.87 cm) and flower diameter (9.57 cm) were significantly maximum in 20 mg/L spermidine. Pigments viz., chlorophyll and anthocyanins were observed maximum in plants treated with 20 mg/L spermine followed by 2% Novel spray. Further, plants treated with 2% Novel showed maximum flower yield (16.98 flowers/plant) as well as showed maximum vase life (8.53 days).

Keywords: Rose, Polyamines, Spermine, Spermidine, Banana pseudostem sap (Novel)

Comment [h1]: Abstract should be in 100 to 200 words

Comment [h2]: Reduce your abstract to 200 words

Formatted: Space After: 0 pt

INTRODUCTION

Protected cultivation of flower crops is gaining significance for quality flower production both for National market as well as exports. Rose (*Rosa hybrida*L.) from Rosaceae family is one of the most commercial and popular cut flowers in the world and holds leading position in planting area, production rate, employment, and exports (4, 7). In fierce competitive international flower market, quality cut flower production plays important role wherein Indian roses suffer from lower prices owing to poor quality (11). From this perspective, the application polyamines can be an effective way to improve the quantitative and qualitative traits in rose (Farahiet al., 6). Polyamines viz., spermine, spermidine and putrescine are a new class of growth hormones that are ubiquitous in living organisms and involved in many plant developmental processes, including cell division, embryogenesis, reproductive organ development, root growth, as well as floral initiation and development (8,2). Significant effect of polyamines on flower quality, vase life and yield has been reported in chrysanthemum (10), Gladiolus (24) and in rose (6-5,26). Biostimulant viz., Novel is composed of banana pseudo stem sap, a patented product of Navsari Agricultural University, that has been known to have some special properties relating to various growth phenomena and is a potential source of cellulose, consisting of different phytochemicals. It is a rich source of macro and micro nutrients as well as contains growth regulators like GA₃ and cytokinin (19, 5). Novel has been found to effectively influence good and qualitative production in different horticultural crops (21, 5, 18). Cow urine has also been known to have presence of some growth promoting auxins (29) along with N, P, K, Ca, Mg, S and others (20) and influence plant growth in flower crops (15, 23). Influence of polyamines and biostimulants like banana pseudo stem sap and cow urine on cut flower crops under protected conditions has not been extensively studied. Hence, this experiment was planned with the basic objective to study the influence of poyamines (spermine and spermidine) and biostimulants (Novel and cow urine) as foliar spray on vegetative and flowering parameters in rose plants.

MATERIALS AND METHODS

The experiment was conducted consecutively for three years during 2015-16 to 2017-18 under naturally ventilated polyhouse at greenhouse complex, Department of Floriculture and Landscape architecture, Navsari Agricultural University, Navsari, Gujarat, India. The

Comment [h3]: 1.You should start by 1. It seems like you copy in your thesis and paste here without changing numeration

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

research work was carried out on one year old plants of Hybrid Tea rose variety 'Top Secret' that was cultivated on raised bed. Foliar spray treatments consisted of spermine (5, 10 and 20 mg/L), spermidine (5, 10 and 20 mg/L), 2% Novel i.e. banana pseudostem sap, 2% cow urine and water as control. Plants were given foliar spray twice at fifteen days interval after pruning starting in the second week of November of each year. Traits were measured as following: plant height with scale, number of branches and number of leaves by counting, length of flower stem by scale, flower stem diameter, bud diameter and flower diameter by *Vernier Calipers* and number of flowers per plant and vase life (days to wilting) were counted. Bud diameter was measured at tight bud stage while flower diameter was measured at fully opened flower stage. Number of flowers per plant was counted based on harvest flowers every month after regular pinching and bending practices. Besides, total chlorophyll content from leaves was determined by following the method of Yoshida *et al.*, (28) and anthocyanins from flowers was estimated based on method of Lees and Francis (13) and expressed in mg/g of fresh weight.

Formatted: Highlight

Formatted: Highlight

The statistical analysis was done by adopting the standard procedures of Panse and Sukhatme (13), analysis was done following statistical NAU software for CRD design and the results were interpreted.

Formatted: Highlight

RESULTS AND DISCUSSION

The vegetative attributes *viz.* plant height, number of branches and number of leaves were found to be significantly increased in rose plants given foliar spray of polyamines and banana pseudo stem sap as observed during 1st year, 2nd year, 3rd year as well as pooled analysis. Comparing different treatments, significantly maximum plant height (116.33, 120.27, 118.20 and 118.27 cm) was found in plants treated with 20 mg/L spermidine, significantly maximum number of branches (6.73, 6.80, 7.20 and 6.91) and number of leaves (126.67, 131.77, 129.47 and 129.30)(Table 1) were observed with 20 mg/L spermine and were at par with 10 mg/L spermine and followed by 20 mg/L spermidine and 2% Novel as seen respectively during the three years as well as pooled analysis.

Polyamines, a new class of plant growth regulators plays role as Hormonal Second Messengers (HSM) (8) and thereby known to implicate biological processes like protein synthesis (27, 2). Besides, spermine and spermidine seem to mediate sensing signal mechanisms and nitrogen assimilation and carbon metabolism in different plant organs and

Formatted: Space After: Auto

Formatted: Underline

Formatted: Highlight

thereby promote plant growth. Foliar application of polyamines significantly promoted vegetative growth in chrysanthemum (10) and gladiolus (16). Besides, Novel i.e. banana pseudostem sap has also been reported to enhance vegetative growth in different horticulture crops (5).

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Flowering parameters also were found to be improved by spraying of polyamines and banana pseudostem sap (Novel) during individual year as well as in pooled. Flower ~~stem length~~ stem length (81.00, 84.27, 86.07 and 83.78 cm) and flower stem diameter (8.17, 8.27, 7.43 and 7.96 mm) were significantly maximum with treatment of 20 mg/L spermine, at par with 10 mg/L spermine and followed by 2% Novel in the pooled data analysis. Further, pigments viz., chlorophyll (3.70, 4.02, 4.18 and 3.96 mg/g) and anthocyanins (2.17, 2.33, 2.42 and 2.30 mg/g) were observed maximum in plants treated with 20 mg/L spermine which is followed by 2% Novel and 20 mg/L spermidine. Exogenous application of polyamines in carnation has been also shown to retard chlorophyll loss and senescence (12). Further, increase in the chlorophyll content in leaves was also observed with foliar application of polyamines in gladiolus (16) and in chrysanthemum (10). Polyamines being involved in plant biological process may have promoted better plant growth in form of flower stalk length as also envisaged by other researchers in cut flowers (16, 6).

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Flower size, yield and vase life were highly influenced by foliar application of spermine and Novel. Bud diameter was maximum (6.03, 6.10, 5.83 and 5.99 cm) (Table ????) with treatment of 10 mg/L spermine, which was at par with 20 mg/L spermine and 20 mg/L spermidine while maximum flower diameter (9.80, 9.87, 9.03 and 9.57 cm) (Table ???) was found with 20 mg/L spermidine, followed by 2 % Novel and 20 mg/l spermine. Plants treated with 2% Novel showed maximum flower yield (15.17, 18.30, 17.47 and 16.98) (Table ???) as well as showed maximum vase life (8.50, 8.57, 8.53 and 8.53 days) (Table ???) followed by 20 mg/L spermine treatment. Novel i.e. banana pseudo stem sap has been known to possess nutrients like N and K along with gibberellins and cytokyanins which may have influenced flower quality and yield. Foliar application of Novel i.e. banana pseudostem sap has been found to increase yield in spinach (22), onion (21) marigold (9) and mango (18). Polyamines are known to be associated with the physiology of flowering metabolite synthesis (3, 25) and correlation between polyamines and flowering processes has been observed earlier in *Arabidopsis* (1) and in chrysanthemum (10). Beneficial effects on flower parameters have also been reported with polyamines in gladiolus (16), carnation (14) and with banana pseudostem sap in marigold (9).

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Conclusion

In conclusion, the foliar application of 20 mg/L spermine as well as 2% Novel i.e. banana pseudostem sap twice at fifteen days interval after pruning has been found to be highly beneficial in improving flower quality and yield in cut flowers of rose under protected cultivation.

Comment [h4]: Rewrite

Comment [h5]: Remind the objective of your study
Main results
Concluding words

Formatted: Strikethrough

REFERENCES

1. Applewhite, P.B., Kaur-Sawhney, R. and Galston, A.W. 2000. A role of spermidine in the bolting and flowering of *Arabidopsis*. *Physiol. Plant.* **79**: 635-640.
2. Bais, H.P. and Ravishankar, G.A. 2002. Role of polyamines in the ontogeny of plants and their biotechnological applications. *Plant Cell Tissue Organ Cult.* **69**: 1-34.
3. Cabanne, F., Dalebroux, M.A., Martin, J. and Martin, C. 1981. Hydroxycinnamic acid amides and ripening to flower of *Nicotianatabacum* L. var. Xanthi. *Physiol. Plant.* **53**: 399-404.
4. Chaudhary, U.C., Singh, A., Ahlawat, T.R. and Palagani, N. 2016. Influence of nitrogen on growth parameters and leaf nutrient composition of rose cv. Samurai under protected conditions. *The Bioscan*, **3**: 1377-1380.
5. Desai, C.S., Patel, J.M., Pawar, S.L., Usadadia, V.P., Naik, V.R. and Savani, N.G. 2016. Value added products form banana pseudostem, Soil and water management research unit, Navsari Agricultural University, Navsari,
6. Farahi, H.F., Khalingi, A., Kholdbarin, B., Akbar, M.M., Eshghi, S., Kavooosi, B. and Aboutalebi, A. 2012. Influence of exogenous spermidine on quality properties and vase life of rose (*Rosa hybrida* cv. Dolcvita). *Ann. Biol. Res.* **3**(10): 4758-4763.
7. Farahi, M.H. and Jahroomi, A.A. 2018. Effect of Pre-harvest Foliar Application of Polyamines and Calcium Sulfate on Vegetative Characteristics and Mineral Nutrient Uptake in *Rosa hybrida*. *J. Ornam. Plant.* **8** (4): 241-253.
8. Galston, A.W., Kaur-Sawhney, R., Altabella, T. and Tiburcio A.F. 1997. Plant polyamines in reproductive activity and response to abiotic stress. *Acta. Bot.* **110**: 197-207.

9. Jadhav, P.B., Singh, A., Mangave, B.D., Patil, N.B., Patel, D.J., Dekhane, S.S. and Kireeti, A. 2014. Effect of organic and inorganic fertilizers on growth and yield of African marigold (*Tagetes erecta* L.) Cv. PusaBasantiGaiinda. *Ann. Biol. Res.* **5**(9): 10
10. Kandil, M.M., El-Saad, M.B., Mona, H.M., Afaf, M.H. and Iman, M.E. 2011. Effect of putrescine and uniconazole treatments on flower characters and photosynthetic pigments of *Chrysanthemum indicum* L. plant. *J. Am. Sci.* **7** (3): 399-408.
11. Kumar, M.S., Ponnuswami, V., Jawaharlal, M. and Kumar, R.A. 2012. Effect of plant growth regulators on growth, yield and exportable quality of cut roses. *The Bioscan.* **7**(4): 733-738.
12. Lee, M.M., Lee, S.H. and Park, K.Y. 1997. Effects of spermine on ethylene biosynthesis in cut carnation (*Dianthus caryophyllus* L.) flowers during senescence. *J. Plant Physiol.* **151**: 68-73.
13. Lees, D.H. and Francis, F.J. 1972. Standardization of pigment analysis in cranberries. *Hortscience.* **7**: 83.
14. Mahagoub, M.H., El-Ghorrab, M. and Bekheta, M. 2006. Effect of some bioregulators on the endogenous phytohormones, chemical composition, essential oil and its antioxidant activity of carnation (*Dianthus caryophyllus* L.). *J. Agric. Sci. Mansoura Univ.*, **31** (7): 4229-4245.
15. Mankar, S.S., Haldankar, P.M., Rangwala, A.D. and Joshi, G.D. 2003. Effect of growth regulators and organic compound on growth, flowering and yield of China aster (*Callistephus chinensis*) var. "Ostrich Plume Mixed". *South Indian Hort.* **52** (1-6): 210-212.
16. Nahed, A.A.G., Taha, L. and Ibrahim, S.M.M. 2009. Some studies on the effect of putrescine, ascorbic acid and thiamine on growth, flowering and some chemical constituents of gladiolus plants at Nubaria. *Ozean J. Appl. Sci.* **2**(2): 1943-2429.
17. Panse, V.G. and Sukhatme, P.V. 1985. Statistical methods for agricultural workers. Indian Council of Agricultural Research, New Delhi.
18. Patel, S., Sharma, H. and Vaja, S. 2017. Effect of Banana Pseudostem Sap and *P. fluorescens* on Mango Diseases in Field Condition. *Int.J.Curr.Microbiol.Appl.Sci.* **6**(11): 514-521
19. Patil, A.P. 2008. Nutrient, banana pseudostem sap and sucker management in ratoon banana under drip irrigation. M.Sc. Thesis, ASPEE College of Horticulture and Forestry, Navsari Agricultural University, Navsari.

20. Phrimantoro. 1995. Cow urine as element of plant hara Stockholm. Environment Institute, EcoSanRes Series, 2009-1, website <http://agrotechnopark.blogspot.in/2011/08/cow-urine-as-element-of-plant-hara.htm>.
21. Salunkhe, J.R., Patel, A.M., Patil, R.G. and Pisal, R.R. 2013. Effect of banana pseudostem sap as liquid fertilizer in onion. *Indian J. Agric. Res.* **43**(7): 258-262.
22. Satashiya, K.F., Patel, K.G., Patel, D.P. and Patel, A.M. 2013. Comprehensive banana pseudostem sap and fertilizer application on yield and nutrient uptake by amaranths and spinach for manipulation soil health and sustainable agriculture. *Ecol. Environ. Conserv.* **19** (04): 1139-1142.
23. Shah, H.P., Singh, A., Patel, D.P. and Tandel, M.B. 2018. Influence of Inorganic and Organic fertilizers on growth and yield of *Dendrobium* Orchid cv. Sonia 17. *Int. J. Curr. Microbiol. Appl. Sci.* **7**(11): 299-304
24. Sivaprakasam, G., Singh, V.P. and Arora, A. 2009. Physiological and molecular analysis of effect of spermine on senescing petals of gladiolus. *Indian J. Plant Physiol.* **14** (4):384-391.
25. Slocum, R.D. and Galston, A.W. 1985. Changes in polyamines associated with past fertilization and development in tobacco ovary tissue. *Plant Physiol.* **79**: 336-343.
26. Tatte, S., Singh, A. and Ahlawat, T.R. 2015. Effect of polyamines on postharvest quality and vase life of rose var. Samurai. *The Bioscan*, **10**(2): 675-678.
27. Van den Broeck, D., Van Den, S.D., Van Montague, M. and Caplan, A. 1994. A group of chromosomal proteins is specifically released by spermine and loses DNA-binding activity upon phosphorylation. *Plant Physiol.* **106**: 559-566.
28. Yoshida, S., Forno, D.S., Cock, J.H. and Gomez, K.A. 1971. Laboratory manual for physiological studies of rice. IRRI, Philippines. pp.1- 43.
29. Zhang, Y. 2000. The origin and summary of the theory of the cloning of mammal. *Sci. and Tech. Review.* **2**:10-13.

Table 1: Effect of foliar spray of polyamines and biostimulants in Rose (*Rosa hybrida* L.) underpolyhouse condition on Plant height (cm), Number of Branches and Number of Leaves.

Sr.No.	Treatments	Heightof plant (cm)				Numberof branches				Numberof leaves			
		1 st year	2 nd year	3 rd year	pooled	1 st year	2 nd year	3 rd year	pooled	1 st year	2 nd year	3 rd year	Pooled
1	T1: 5 mg/L Spermine	108.33	113.43	105.33	109.03	3.93	4.10	3.87	3.97	116.33	120.10	119.93	118.79
2	T2: 10 mg/L Spermine	116.00	119.37	109.73	115.03	6.03	6.17	6.73	6.31	125.33	128.23	128.93	127.50
3	T3: 20 mg/L Spermine	112.67	117.73	110.07	113.49	6.73	6.80	7.20	6.91	126.67	131.77	129.47	129.30
4	T4: 5 mg/L Spermidine	90.67	95.87	101.20	95.91	3.73	3.83	4.27	3.94	111.00	114.90	118.27	114.72
5	T5: 10 mg/L Spermidine	108.33	116.07	111.93	112.11	4.93	5.10	4.40	4.81	119.00	122.43	119.67	120.37
6	T6: 20 mg/L Spermidine	116.33	120.27	118.20	118.27	4.43	4.67	5.53	4.88	122.33	125.87	126.90	125.03
7	T7: 2% Novel	102.00	107.27	104.47	104.58	4.73	4.90	4.93	4.86	119.00	121.67	118.80	119.82
8	T8: 2% Cow urine	98.00	101.67	101.67	100.44	3.17	3.23	4.20	3.53	110.00	112.40	121.53	114.64
9	T9: control	98.33	102.63	104.80	101.92	2.60	2.67	3.13	2.80	106.33	109.13	117.80	111.09
T	S.Em ±	2.79	2.30	2.17	1.37	0.18	0.17	0.27	0.13	3.57	1.96	2.18	1.51
	C.D. 0.05	8.29	6.84	6.45	4.07	0.52	0.50	0.81	0.39	10.62	5.83	6.48	4.49
Y	S.Em ±	-	-	-	0.82	-	-	-	0.07	-	-	-	0.90
	C.D. 0.05	-	-	-	2.35	-	-	-	0.20	-	-	-	2.58
T X Y	S.Em ±	-	-	-	2.46	-	-	-	0.20	-	-	-	2.70
	C.D. 0.05	-	-	-	NS	-	-	-	0.57	-	-	-	NS
	CV%	4.57	3.61	3.50	3.82	6.81	6.38	9.58	8.43	5.27	2.82	3.09	3.76
					3.95				7.52				3.89

Comment [h6]: Did you counted manually or there's a scientific method?

Table 2: Effect of foliar spray of polyamines and biostimulants in Rose (*Rosa hybrida* L.) under polyhouse conditions on flower stem length (cm), flower stem diameter (mm) and bud length (cm)

Sr.No.	Treatments	Flower stem length (cm)				Flower stem diameter (mm)				Bud length (cm)			
		1 st year	2 nd year	3 rd year	pooled	1 st year	2 nd year	3 rd year	pooled	1 st year	2 nd year	3 rd year	pooled
1	T1: 5 mg/L Spermine	68.00	69.67	77.07	71.58	5.60	6.00	7.10	6.23	4.77	4.80	5.07	4.88
2	T2: 10 mg/L Spermine	78.67	82.37	80.27	80.43	7.93	7.97	7.17	7.69	5.07	5.13	5.23	5.15
3	T3: 20 mg/L Spermine	81.00	84.27	86.07	83.78	8.17	8.27	7.43	7.96	5.50	5.67	5.50	5.56
4	T4: 5 mg/L Spermidine	68.00	70.83	76.07	71.63	5.57	6.33	6.83	6.24	4.60	4.70	5.20	4.83
5	T5: 10 mg/L Spermidine	73.00	74.57	73.80	73.79	6.90	7.07	7.13	7.03	5.30	5.33	5.73	5.46
6	T6: 20 mg/L Spermidine	75.67	77.87	78.07	77.20	7.37	7.43	7.23	7.34	5.83	5.90	5.87	5.87
7	T7: 2% Novel	77.67	80.70	81.07	79.81	7.47	7.67	7.20	7.44	5.37	5.40	5.23	5.33
8	T8: 2% Cow urine	59.00	64.27	72.20	65.16	5.53	5.67	6.87	6.02	5.47	5.43	5.53	5.48
9	T9: control	52.67	56.70	69.20	59.52	4.70	5.07	6.77	5.51	4.90	4.97	5.20	5.02
T	S.Em ±	2.38	1.78	1.93	1.66	0.21	0.20	0.30	0.14	0.11	0.12	0.26	0.09
	C.D. 0.05	7.08	5.28	5.72	4.93	0.62	0.60	NS	0.42	0.33	0.35	NS	0.27
Y	S.Em ±	-	-	-	0.49	-	-	-	0.08	-	-	-	0.06
	C.D. 0.05	-	-	-	1.41	-	-	-	0.23	-	-	-	NS
T X Y	S.Em ±	-	-	-	1.47	-	-	-	0.25	-	-	-	0.18
	C.D. 0.05	-	-	-	4.22	-	-	-	0.72	-	-	-	NS
	CV%	5.86	4.19	4.33	6.75	5.52	5.16	7.39	6.02	3.66	3.89	8.36	5.28
					3.44				6.22				6.04

Table 3: Effect of foliar spray of polyamines and biostimulants in Rose (*Rosa hybrida* L.) under polyhouse conditions on bud diameter (cm), flower diameter (cm), number of flowers per ~~plant~~ plant and vase life (days)

Sr. No.	Treatments	Bud diameter (cm)				Flower diameter (cm)				Number of flowers/ plant				Vase life (days)			
		1 st year	2 nd year	3 rd year	pooled	1 st year	2 nd year	3 rd year	pooled	1 st year	2 nd year	3 rd year	pooled	1 st year	2 nd year	3 rd year	pooled
1	T1: 5 mg/L Spermine	5.03	5.13	5.53	5.23	7.13	7.23	8.13	7.50	11.37	11.87	14.20	12.47	6.10	6.20	7.80	6.70
2	T2: 10 mg/L Spermine	6.03	6.10	5.83	5.99	7.97	8.03	8.17	8.06	13.03	14.40	14.87	14.10	8.40	8.43	8.60	8.48
3	T3: 20 mg/L Spermine	5.80	5.83	5.90	5.84	8.67	8.73	8.80	8.73	14.67	16.37	16.80	15.94	8.50	8.47	8.33	8.43
4	T4: 5 mg/L Spermidine	4.80	4.87	5.30	4.99	7.77	7.80	7.90	7.82	11.03	11.30	14.33	12.22	5.67	5.73	6.80	6.07
5	T5: 10 mg/L Spermidine	5.10	5.17	5.63	5.30	8.10	8.23	8.30	8.21	12.37	12.87	14.73	13.32	8.23	8.30	8.07	8.20
6	T6: 20 mg/L Spermidine	5.87	5.90	5.97	5.91	9.80	9.87	9.03	9.57	13.70	15.13	15.60	14.81	8.60	8.63	8.27	8.50
7	T7: 2% Novel	5.37	5.50	5.73	5.53	9.00	9.10	8.77	8.96	15.17	18.30	17.47	16.98	8.50	8.57	8.53	8.53
8	T8: 2% Cow urine	4.97	5.00	5.43	5.13	8.60	8.63	8.30	8.51	12.77	13.50	14.73	13.67	6.43	6.67	7.80	6.97
9	T9: control	4.57	4.60	5.47	4.88	7.07	7.10	7.97	7.38	11.37	12.13	14.40	12.63	6.00	6.07	7.13	6.40
T	S.Em ±	0.22	0.23	0.30	0.16	0.28	0.26	0.26	0.17	0.39	0.34	0.44	0.26	0.24	0.22	0.23	0.21
	C.D. 0.05	0.64	0.67	NS	0.48	0.84	0.78	NS	0.51	1.16	1.01	1.29	0.77	0.72	0.66	0.69	0.62
Y	S.Em ±	-	-	-	0.08	-	-	-	0.08	-	-	-	0.12	-	-	-	0.05
	C.D. 0.05	-	-	-	0.23	-	-	-	NS	-	-	-	0.34	-	-	-	0.14
T X Y	S.Em ±	-	-	-	0.24	-	-	-	0.25	-	-	-	0.36	-	-	-	0.14
	C.D. 0.05	-	-	-	NS	-	-	-	NS	-	-	-	1.03	-	-	-	0.40
	CV%	7.06	7.34	9.04	8.59	5.97	5.46	5.39	6.27	5.27	4.23	4.94	5.58	5.69	5.17	5.09	8.08
					7.55				5.25				4.40				3.11

Table 4: Effect of foliar spray of polyamines and biostimulants on in Rose (*Rosa hybrida* L.) underpolyhouse conditions on Chlorophyll and Anthocyanin content

Sr.No.	Treatments	Chlorophyll (mg/g)				Anthocyanin (mg/g)			
		1 st year	2 nd year	3 rd year	pooled	1 st year	2 nd year	3 rd year	pooled
1	T1: 5 mg/L Spermine	2.86	2.97	2.73	2.85	1.7	1.68	1.82	1.73
2	T2: 10 mg/L Spermine	3.36	3.47	3.46	3.43	1.78	1.88	2.03	1.90
3	T3: 20 mg/L Spermine	3.70	4.02	4.18	3.96	2.17	2.33	2.42	2.30
4	T4: 5 mg/L Spermidine	3.11	3.04	3.15	3.10	1.68	1.79	1.78	1.75
5	T5: 10 mg/L Spermidine	3.15	3.26	3.34	3.25	1.83	1.89	1.94	1.89
6	T6: 20 mg/L Spermidine	3.36	3.50	3.58	3.48	1.98	2.10	2.19	2.09
7	T7: 2% Novel	3.49	3.67	3.86	3.67	2.12	2.21	2.31	2.21
8	T8: 2% Cow urine	3.06	3.30	3.33	3.23	1.81	1.87	1.90	1.86
9	T9: control	2.19	2.43	2.50	2.37	1.68	1.74	1.70	1.70
T	S.Em ±	0.14	0.12	0.10	0.099	0.059	0.093	0.10	0.071
	C.D. 0.05	0.43	0.37	0.30		0.17	0.27	0.30	
Y	S.Em ±	-	-	-	0.096	-	-	-	0.061
	C.D. 0.05	-	-	-		-	-	-	
T X Y	S.Em ±	-	-	-	0.032	-	-	-	
	C.D. 0.05	-	-	-		-	-	-	0.020
	CV%	7.99				5.50			