

# Study of the effect of three extraction processes on the phytoactives of *Alchornea cordifolia* (Schumach. & Thonn.) Müll. Arg. (Euphorbiaceae) and evaluation of their antioxidant and analgesic activities

## Abstract

**Aims:** An important aspect of the valorization of plants of interest is the search for phytoactives. This work aims to show the effect of extractive processes (maceration, decoction, Soxhlet extraction) both on the phytoconstituents of a plant species and on their intrinsic properties.

**Methodology:** *Alchornea cordifolia*, a botanical species growing in Côte d'Ivoire, was chosen for this purpose. Phytophenols, flavonoids and condensed tannins were quantified by spectrophotometry. Antioxidant (DPPH and FRAP) and analgesic activity were also evaluated.

**Results:** The quantitative analysis of total phytophenols, total flavonoids and condensed tannins of macerated, decocted and Soxhlet extracts of the plant organs (leaves, stems, roots), highlighted their respective contents, varying proportionally according to the extraction conditions. The same observation was made at the end of the estimation of the antioxidant (DPPH and FRAP tests) and analgesic activities of these different extraction fractions.

**Conclusion:** This study has demonstrated the effect of three extraction processes on some phytoactives quantified of *A. cordifolia*

**Key words:** *Alchornea cordifolia*, phytoactives, DPPH, FRAP, analgesic activity.

## Introduction

Species of the family of Euphorbiaceae, *Alchornea cordifolia* is a plant used in traditional medicine, in decoction or maceration for its analgesic, anti-inflammatory, antimicrobial, antiparasitic, veinotonic, antioxidant, antidiabetic and astringent properties [1; 2]. Biological and pharmacological activities of this plant can be attributed to the phytoactives it contains, which prevent or slow down oxidation by neutralizing free radicals [3; 4; 5]. Their extraction from a plant matrix by means of a process appropriate is an initial and necessary step before any analysis. It depends on several factors, in particular the nature of the solvent, the temperature, the method and the extraction time [6]. Optimizing the extraction yield of phytoactives is necessary since the effectiveness of extracts depends on their chemical composition [7]. *A. cordifolia* is a perennial shrub, bushy, erect or climbing, which can reach 4 m in height. Its leaves are simple, alternate with the blade cordate at the base and acuminate at the top. Its flowers are greenish-white in color, and its small fruits are capsular with three compartments containing red seeds [8; 9]. Our study was initiated with the intention to understand how traditional extraction processes have an impact on phytoactives and their antioxidant and analgesic activities. Thus, we focused our attention on the phytophenols of *A. cordifolia*.

## 2. Materials and methods

### 2.1. Plant material

The plant material consists of the leaves, stems and roots of *Alchornea cordifolia*, crescent on the NANGUI ABROGOUA University site (Abidjan / Ivory Coast) (5° 20' 11" north, 4° 01' 36" west). After identification with respect to the species in the herbarium of the Center National de Floristique (CNF) of the Félix HOUPOUËT-BOIGNY University (Cocody / Abidjan) and authentication (N°AD 224) [10], the organs were cleaned, dried under permanent air conditioning (20°C, 15 days) and reduced to powder using an electric grinder.

### 2.2. Methods

#### 2.2.1. Preparations of extracts

Three extraction processes were used: maceration, decoction and Soxhlet extraction. The extracts were filtered using a funnel and a Büchner flask and then reduced with a BÜCHI type EL-131 rotary evaporator.

### **2.2.2. Quantification of total phytophenols, total flavonoids and condensed tannins**

#### ***Assay of total phytophenols (PT)***

The PT contents are determined by adding to 1 ml of alcoholic extract, 1.5 ml of sodium carbonate ( $\text{Na}_2\text{CO}_3$ ) (17%, m/v) and 0.5 ml of Folin-Ciocalteu reagent (0.5N). The mixture is incubated for 30 min in the dark and the absorbance read at 760 nm with a UV-visible spectrophotometer (Spectro AL 800). Calibration with gallic acid (0-1000  $\mu\text{g}/\text{ml}$ ) is carried out under the same conditions and the contents are expressed in milligrams of gallic acid equivalent per gram of dry matter (mg EAG/g DM) [11; 12].

#### ***Assay of total flavonoids (TF)***

The FT contents were determined by adding to 100  $\mu\text{l}$  of Neu's reagent, 2 ml of alcoholic plant extract or quercetol (0.05 mg/ml). The absorbance of the mixture is read at 404 nm (UV-visible spectrophotometer (Spectro AL 800)) and the contents are determined according to formula (1) [13].

$$\text{TF (\%)} = ((0,005 \times \text{Aext} / \text{Aq}) \times \text{Cext}) \times 100 \quad (1)$$

**Ext:** Absorbance of the extract; **Aq:** Absorbance of quercetol; **Cext:** Concentration of the extract.

The proportions of TF determined are subsequently expressed in mg EAG/g DM relative to the TP contents.

#### ***Dosage of condensed tannins (TC)***

To 400  $\mu\text{l}$  of alcoholic extract are added 3 ml of a methanolic solution of vanillin (4%, m/v) and 1.5 ml of concentrated hydrochloric acid (HCl). After 15 min at room temperature, the absorbance of the mixture is read at 500 nm (UV-visible spectrophotometer (Spectro AL 800)). A calibration line is drawn with catechin (0-300  $\mu\text{g}/\text{ml}$ ), prepared under the same conditions. The TC contents obtained are expressed in milligram catechin equivalent per gram of dry extract (mg EC/g DM) [14].

### **2.2.3. Evaluation of antioxidant activity**

#### ***DPPH test***

To 0.4 ml of alcoholic plant extract are added 1.2 ml of a methanolic solution of DPPH (0.03 mg/ml). The mixture is adjusted with 2.4 ml of methanol (MeOH), and the absorbance read at 517 nm every 3 min for 30 min against a blank. Plant extracts and ascorbic acid (vitamin C, reference antioxidant) are prepared at concentrations (0.128; 0.0357; 0.0014; 0.001 and 0.00071 mg/ml) [15]. The percentage reductions (PR) of DPPH are determined from equation (2):

$$\text{PR (\%)} = ((1 - (\text{Ae} / \text{Ab})) \times 100) \quad (2)$$

**Ae:** Absorbance of the extract; **Ab:** Absorbance of DPPH

The effectiveness of the samples against DPPH was assessed by graphical determination of the  $\text{CR}_{50}$  (concentration that reduces 50% of DPPH) [16; 17].

#### ***FRAP test***

The FRAP reagent used is a mixture of sodium acetate buffer (300 mM, pH = 3.6), TPTZ (2, 4, 6-Tri (2-pyridyl)-s-triazine at 10 mM prepared in a solution of HCl at 40 mM) and iron trichloride ( $\text{FeCl}_3$ ) (20 mM) in a volume ratio of 10:1:1. The freshly prepared FRAP reagent is previously heated to 37°C in a water bath. 100  $\mu\text{l}$  of Trolox (0.187; 0.375; 0.75 and 1.5 mM) (reference antioxidant) and the samples to be tested are prepared at 0.25 mg/ml, and added to 3 ml of the FRAP reagent. The absorbance of the complex (TPTZ- $\text{Fe}^{3+}$ ) is read at 593 nm after 4 min. The results are expressed in mM Trolox equivalent with the calibration line constructed from the Trolox [18; 19].

### **2.3.4. Assessment of analgesic activity**

The formaldehyde test is used to assess the effectiveness of plant extracts on pain. The control batch of mice received distilled water while the other batches were treated with the extracts (60 and 120 min) of *A. cordifolia* at 100 mg/kg, acetyl salicylic acid (aspirin) (100 mg/kg) and tramadol

hydrochloride (trabar) (30 mg/kg). Male and female mice of the species *Mus musculus* of mass between 18 and 22 g, aged 5 to 6 weeks were used. They come from the Physiology, Pharmacology and Phytotherapy laboratory the Natural Sciences Training and Research Unit (UFR-SN) of the UNA. One hour before the injection of 20 µl of the formaldehyde solution (2.5%) into the right hind paw, the mice received the extracts and the reference products orally. Immediately after the injection of the formaldehyde solution, the number of licks of the treated paw is counted during the first five minutes, then between 15-30 min [20; 21]. The analgesic effect is determined by the formula (3):

$$\% \text{ Inhibit} = ((\text{NI control} - \text{NI treated}) / \text{NI control}) \times 100 \quad (3)$$

**NI treated** =number of licks of the paw of the mice of the control group;

**NI control** =number of licks of the paw of the mice of the treated group.

## 2.4. Statistical analyzes

The results are expressed as mean ± standard deviation. One-way analysis of variance (ANOVA) was used to compare total phytophenol contents, antioxidant activities, and formaldehyde-induced pain inhibition using Graph Pad Prism 5 software. When a significant difference exists between the means ( $P < 0.05$ ), the multiple comparison test of the means is carried out (Tukey's test) to determine the level of difference between the different groups of extracts [22] (Westlake, 1971).

## 3. Results and discussion

### 3.1. PT levels

The PT contents of *A. cordifolia* (Table 1) were determined from the linear regression equation for gallic acid (expressed in mg EAG / g DM) taken as standard ( $y = 0.0232 x + 0.0002$ ;  $R^2 = 0.9983$ ). The proportions are variable with significant maximum levels ( $p < 0.001$ ) observed with the Soxhlet extracts of leaves (120 min:  $183.57 \pm 0.01$ ). The maximum PT content in the stems was obtained by maceration (120 min:  $137.28 \pm 0.52$ ) and that of the roots by decoction ( $77.53 \pm 0.28$ ). Whatever the organ, TP levels in macerated extracts increase over time (60 min to 120 min). In addition, the roots have significantly ( $p < 0.001$ ) the lowest phytophenol contents for any of the extraction methods.

**Table 1: PT levels in the organs of *A. cordifolia***

Organ	Time (mins)	PT content (mg EAG/g DM)		
		Macerated	Decocted	Soxhlet extract
Leaves	60	$108.22 \pm 0.15$	$129.26 \pm 0.02$	$155.12 \pm 0.05$
	90	$109.09 \pm 0.01$	$118.91 \pm 0.08$	$175.03 \pm 0.01$
	120	$111.5 \pm 0.15$	$114.6 \pm 0.4$	$183.57 \pm 0.01$
Stems	60	$120.12 \pm 1.49$	$121.50 \pm 0.12$	$112.02 \pm 0.15$
	90	$125.72 \pm 0.45$	$125.12 \pm 0.15$	$133.05 \pm 0.15$
	120	$137.28 \pm 0.52$	$113.40 \pm 0.14$	$117.45 \pm 0.15$
Roots	60	$64.86 \pm 0.26$	$73.4 \pm 0.15$	$48.66 \pm 0.83$
	90	$71.67 \pm 0.15$	$77.53 \pm 0.28$	$52.53 \pm 0.65$
	120	$72.28 \pm 0.14$	$70.55 \pm 0.40$	$61.41 \pm 0.68$

The results demonstrated that the PT increase when the extraction time increases. Indeed, a high extraction time allows a prolonged contact between the solutes and the solvent, resulting in a large mass transfer [23]. In addition, there is a decrease in PT content in the decoctions and Soxhlet extracts, probably due to heating, which dehydrates the cell walls allowing easy penetration of the

extraction solvent, and the release of solutes [24]. The increase in temperature promotes the extraction of phytophenols by increasing the diffusion coefficient, the solubility of solutes in the extractor and the decrease in its viscosity. [25]. Studies have shown that increased extraction time and prolonged exposure to high temperatures potentially increase the loss of phytophenols [26; 27].

### 3.2. TF content

The TF contents (in mg EAG/g DM) estimated (Table 2) are in variable proportions, depending on the organ, the process and the extraction time. Whatever the extraction process, TF contents in the leaves are significantly higher ( $p < 0.001$ ) than those of the stems and roots.

**Table 2: TF contents of *A. cordifolia* organs**

Organ	Time (mins)	TF content (mg EAG/g DM)		
		Macerated	Decocted	Soxhlet extract
Leaves	60	28.52 ± 0.13	42.05 ± 0.25	44.54 ± 0.35
	90	31.21 ± 0.25	35.82 ± 0.01	56.24 ± 0.22
	120	33.92 ± 0.22	33.75 ± 0.13	64.38 ± 0.82
Stems	60	10.46 ± 0.14	13.09 ± 0.01	11.7 ± 0.94
	90	10.94 ± 0.152	13.57 ± 0.14	20.13 ± 0.39
	120	12.22 ± 0.38	9.79 ± 0.08	13.29 ± 0.51
Roots	60	10.07 ± 0.08	12.09 ± 0.01	11.27 ± 0.12
	90	11.23 ± 0.14	19.51 ± 0.09	12.83 ± 0.62
	120	12.24 ± 0.30	21.16 ± 0.08	17.88 ± 0.06

The TF contents of macerated extracts increase with time (leaves: 33.92 ± 0.22 mg EAG/g DM; stems: 12.22 ± 0.38 mg EAG/g DM; roots: 12.24 ± 0, 30 mg EAG/g DM at 120 mins). Concerning the roots, the TF contents of the decoctions and Soxhlet extracts increase when the extraction time is prolonged. However, in the decoctions of the leaves and stems, we note the influence of the temperature and the extraction time on the different TF contents. Indeed, flavonoids are both thermo and photosensitive [28].

### 3.3. TC contents

The catechin calibration line ( $y = 0.004 x + 0.006$ ) was used to determine the TC contents of macerated, decocted and Soxhlet extracts. Table 3 summarizes the results obtained. Levels that vary proportionally are more significant in macerated leaves and stems. The TC of macerated extracts increase when the extraction time is prolonged (Leaves: 60 min 31.67 ± 0.62; 90 min 34.17 ± 0.8 and 120 min 36.25 ± 0.95). Whatever the organ, the macerated TC are more significant ( $p < 0.05$ ) than decoctions and Soxhlet extracts. The leaves and stems contain significantly ( $p < 0.001$ ) more TC than roots. However, in decoctions and Soxhlet extracts, the levels drop when the extraction time is extended. The decreases in TC recorded could also be due to their degradation under the effect of heat. Indeed, some phenolic phytochemicals seem to be heat-sensitive, therefore can be degraded by heating. Heating for a prolonged extraction time can lead to the decomposition of target phytochemicals contained in the extracts [29; 30].

**Table 3: TC levels in the organs of *A. cordifolia***

Organ	Time (min)	TC content (mgEC /g DM)		
		Macerated	Decocted	Soxhlet extract
Leaves	60	31.67 ± 0.62	22.5 ± 0.63	10.0 ± 0.62
	90	34.17 ± 0.8	14.17 ± 0.01	28.75 ± 0.30
	120	36.25 ± 0.95	12.92 ± 0.36	12.92 ± 0.40
Stems	60	32.13 ± 0.13	22.21 ± 0.19	17.25 ± 0.01
	90	33.88 ± 0.07	21.46 ± 0.31	22.25 ± 0.13
	120	35.88 ± 0.01	21.08 ± 0.07	19.0 ± 0.08
Roots	60	12.67 ± 0.07	7.38 ± 0.03	3.83 ± 0.07
	90	15.88 ± 0.13	8.04 ± 0.14	4.58 ± 0.26
	120	16.42 ± 0.19	7.88 ± 0.01	6.21 ± 0.14

### 3.4. Antioxidant activity

The antioxidant profile of macerations, decoctions and Soxhlet extracts of *A. cordifolia* was evaluated by spectrophotometry against the DPPH radical and by the FRAP method.

#### 3.4.1 DPPH antioxidant activity

The DPPH antioxidant potential of the extracts evaluated by spectrophotometry made it possible to determine the percentage reductions (PR) of the DPPH radical by the extracts of *A. cordifolia* (Figure 1). The PR are in variable proportions, and have made it possible to graphically determine the concentrations which reduce 50% of the DPPH (CR<sub>50</sub>). This quantitative parameter of the effectiveness of an analyte against DPPH is the concentration necessary to reduce 50% of the initial concentration of the DPPH radical. The lower its value, the more pronounced the antioxidant activity [16; 17]. Irrespective of the extraction method, the CR<sub>50</sub> values of the leaf extracts decrease when the extraction time increases up to 120 min; indicating that the antioxidant activity is enhanced. The prolonged extraction time has a positive impact on the antioxidant activity of the leaf extracts (macerated CR<sub>50</sub> = 0.0017 at 120 min; Decocted CR<sub>50</sub> = 0.0012 at 120 min; Soxhlet extract CR<sub>50</sub> = 0.0012 at 120 min).

The Soxhlet extracts showed greater antioxidant activity than the decoctions, however this difference is not significant. As observed in the leaves, the CR<sub>50</sub> of the stem and root extracts decrease with an increase in the extraction time. However, beyond 90 min, an increase in the CR<sub>50</sub> of decoctions and Soxhlet extracts of stems and roots was observed, corresponding to a drop in antioxidant potential. The antioxidant activity of the extracts could be linked to the presence of quantified phenolic phytochemicals. Indeed, most antioxidants of natural origin have in their molecular structures, hydroxyphenolic groups responsible for the antioxidant character [31; 32]. The results obtained show the influence of temperature and extraction time on the antioxidant profile of the extracts. The decoctions and Soxhlet extracts showed higher antioxidant activity than the macerated ones.

Several studies have shown that decoctions and Soxhlet extracts of plants show greater antioxidant activity than macerated ones [33; 34; 35; 36]. A rise in temperature allows a reinforcement of the antioxidant activity up to a threshold beyond which the antioxidant activity decreases; which would probably be due to the degradation of antioxidant molecules [37; 38]. The extraction time has an influence on the antioxidant activity because a prolongation of the contact time of the extractor and the drug would improve the diffusion of the antioxidant compounds [37; 39]. On the other hand, a prolonged extraction time can expose antioxidant phytochemicals to degradation, polymerization and/or oxidation [25; 39].

T1 (3): 60 min extract after 3 min incubation with DPPH; T1 (15): 60 min extract after 15 min incubation with DPPH; T1 (30): 60 min extract after 30 min incubation with DPPH  
T2 (3): 90 min extract after 3 min incubation with DPPH; T2 (15): 90 min extract after 15 min incubation with DPPH; T2 (30): 90 min extract after 30 min incubation with DPPH;  
T3 (3): 120 min extract after 3 min incubation with DPPH; T3 (15): 120 min extract after 15 min incubation with DPPH; T3 (30): 120 min extract after 30 min incubation with DPPH;

**Figure 1: CR<sub>50</sub> of *A. cordifolia* organ extracts as a function of time and extraction method**

### 3.4.2. FRAP antioxidant activity

The evaluation of the FRAP antioxidant profile was carried out with the extracts of the leaves, stems and roots of *A. cordifolia* by determining their potential for reducing Fe<sup>3+</sup> to Fe<sup>2+</sup> (**Figure 2**). The presence of Fe<sup>2+</sup> ions was evaluated by measuring the blue color of the reaction medium [40]. The FRAP activities of the extracts, estimated at 0.25 mg/ml, were expressed in mM TE/g MS from the Trolox calibration line. The FRAP antioxidant activity of vitamin C, under the same conditions, is 6.45 mM TE/g DM. Leaves and stems showed significantly ( $p < 0.001$ ) the highest FRAP antioxidant activities. However, for leaves, strong FRAP antioxidant activities were recorded in Soxhlet extracts (5.64 mM TE/g MS). In stems, macerations and decoctions were more active than Soxhlet extracts. The maximum activity of the macerated was observed after 120 min while that of the decoctions and extracts with Soxhlet is perceived at 90 min. Overall, FRAP antioxidant activity varies very little depending on extraction method and time. Statistical analysis reveals a significant difference ( $p < 0.001$ ) between the antioxidant activity of vitamin C and those of stem and root extracts. However, this difference is moderately significant ( $p < 0.05$ ) in macerated leaves and decoctions. There is no significant difference between the Soxhlet extracts of the leaves and that of vitamin C.

### Figure 2: FRAP antioxidant profiles of *A. cordifolia* organ extracts

The macerated organs reduce the  $\text{Fe}^{3+}$  ions to  $\text{Fe}^{2+}$ . Some studies revealed that the methanol maceration of the leaves of *A. cordifolia* obtained in 72 h of extraction, showed a FRAP antioxidant activity of 533.33  $\mu\text{M Fe(II) / g}$  against 1005  $\mu\text{M Fe(II) / g}$  for vitamin C [41]. Antioxidants, for the most part, are good reducers. The reducing power of a compound can therefore serve as a meaningful indicator of its potential antioxidant activity [42; 43].

### 3.5 Analgesic activity

The formaldehyde test was used to evaluate the analgesic activity of macerated, decocted and Soxhlet extracts of *A. cordifolia*. Figure 3 presents the results translated into percentage inhibition (%). The 60 min and 120 min extracts were administered at a dose of 100 mg/kg of MC, and compared with the effect of aspirin and trabar on the pain caused. All *A. cordifolia* extracts significantly inhibited phases 1 and 2 of formaldehyde-induced pain, with a remarkable effect in the second phase close to the effect of aspirin (82%). The effect of macerated increases when the extraction time increases (FM2 76.85%; TM2 61.57%; RM2 60.19%). However, a decrease in the effect of decoctions of leaves, roots and Soxhlet extracts of roots was observed.

### Figure 3: Analgesic profiles of *A. cordifolia* organ extracts

FM1 and FM2: macerated leaves obtained respectively in 60 min and 120 min; FD1 and FD2: decoctions of the leaves obtained respectively in 60 min and 120 min; FS1 and FS2: Soxhlet extracts of the leaves obtained respectively in 60 min and 120 min

TM1 and TM2: macerated stems obtained respectively in 60 min and 120 min; TD1 and TD2: stem decoctions obtained respectively in 60 min and 120 min; FS1 and FS2: Soxhlet extracts of stems obtained respectively in 60 min and 120 min

RM1 and RM2: macerated roots obtained respectively in 60 min and 120 min; RD1 and RD2: root decoctions obtained respectively in 60 min and 120 min; RS1 and RS2: Soxhlet extracts of the roots obtained respectively in 60 min and 120 min

This decrease would be due to the thermal degradation of the phytochemicals, responsible for the activity. Overall, leaf and root decoctions exhibited significantly ( $p < 0.001$ ) more effective analgesic activity; whereas that of the Soxhlet extracts is more significant in the stems ( $p < 0.001$ ). The effect of time and the extraction process on the analgesic activity of the extracts is proven. In phase 2 (Figure 3), aspirin inhibited the activity of cyclo-oxygenase which leads to the formation of prostaglandins. These metabolites of arachidonic acid cause inflammation, swelling, pain and fever [44]. In phase 1 (Figure 3), the trabar highlighted local inflammatory pain, which is influenced by inflammatory mediators (serotonin, histamine and prostaglandins) [21]. Similar to the analgesic effect of aspirin, all extracts inhibited in varying proportions phase 2 of the response to formaldehyde. The analgesic effect of the extracts in phase 1 could be related to the inhibition of cyclo-oxygenase. Several studies have reported the analgesic effects of flavonoids, coumarins, tannins and phenolic acids [45; 46]. The presence of phytochemicals in *A. cordifolia*, including phytochemicals, total flavonoids and quantified condensed tannins, seems to attest to the observed analgesic properties.

## Conclusion

The implementation of the present study demonstrated the manifest and palpable effect of three extraction processes (maceration, decoction, Soxhlet extraction) on some quantified phytoactives (polyphenols, flavonoids, condensed tannins) of *A. cordifolia* and their activities. antioxidant (interactions with DPPH and  $Fe^{3+}$  ions) and analgesic. The deductions resulting from this study remain dependent on our experimental choices.

## references

1. Kerharo J, Adam JG. Pharmacopée Sénégalaise traditionnelle - Plantes médicinales et toxiques. Paris Edition Vigot frères, 1974. French.
2. Tra Bi FH, Irié GM, N'gaman KCC, Mohou BHC. Etudes de quelques plantes thérapeutiques utilisées dans le traitement de l'hypertension artérielle et du diabète : deux maladies émergentes en Côte d'Ivoire. Sciences & Nature. 2008; 5(1): 39-48. French.
3. Lamikanra A, Ogundaini AO, Ogungbamila FO. Antibacterial constituents of *Alchornea cordifolia* leaves. Phytotherapy research. 1990; 5(4): 198-200.
4. Okoye FBC, Odimegwu CD, Anyasor NC, Ajaghaku DL, Gugu TH, Osadebe PO. et al. A new antioxidant and antimicrobial compound isolated from *Alchornea cordifolia* leaves. Afric J Pharm Res Develop. 2015; 7(2): 87-94.
5. Noundou XS, Krause RWM, Vuuren VSF, Ndinteh DT, Olivier DK. Antibacterial effects of *Alchornea cordifolia* (Schumacher and Thonn.) Müll. Arg extracts and compounds on gastrointestinal, skin, respiratory and urinary tract pathogens. J Ethnopharm. 2015. 179: 76-82.
6. Liorach R, Thomas-Barberan FA, Ferreres F. Lettuce and chicory by products as a source of antioxidant phenolic extracts. J Agric. Food Chem. 2004; 52: 5109-5116.
7. Santos SS, Deolindo CTP, Esmerino LA, Genovese MI, Fuita A, Marques MB. et al. Effects of time and extraction temperature on phenolic composition and functional properties of red rooibos (*Aspalathus linearis*). Food Res Inter. 2016; 89: 476-487.
8. Adjanohoun E, Aké Assi L. Contribution au recensement des plantes médicinales de Côte d'Ivoire. Ministère de la Recherche Scientifique, Centre national de floristique (CNF) de l'Université Nationale de Côte d'Ivoire, Abidjan. 1979; 358 pages. French.
9. Hawthorne W, Jongkind C. Woody Plants of Western African Forests, A Guide to the Forest trees, Shrubs and Lianes from Senegal to Ghana. Kew Publishing Royal Botanic Gardens. 2006: 734-874.
10. Malan DF, Danho FRN, Kouakou LK. Medicinal plants and traditional healing practices in Ehotile people, around the aby lagoon (eastern littoral of Côte d'Ivoire), J Ethnopharm. 2015; 1-18.
11. Singleton VL, Ortofer R, Lamuela-Raventos RM. Analysis of total phenol and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. Methods in enzymology Orlando Academic Press. 1999; 152-178.

12. N'Gaman KCC. Étude phytochimique et effet d'extraits de *Gmelina arborea* Roxb. (Verbenaceae) de Côte d'Ivoire sur la stabilité osmotique d'érythrocytes. Thèse Université Abobo-Adjamé Côte d'Ivoire. 2013; 152 pages. French.
13. Hariri EB, Sallé G, Andary C. Involvement of flavonoids in the resistance of two poplar cultivars to mistletoe (*Viscum album* L). *Protoplasma*. 1991; 162(1): 20-26.
14. Heilmer D, Vignini P, Dini MG, Vincieri FF, Romani A. Antiradical activity and polyphenol composition of local Brassicaceae edible varieties. *Food Chem*. 2006; 99: 464-469.
15. Espin JC, Soler-Rivas C, Wichers HJ. Characterization of the total free radical scavenger capacity of vegetable oils and oil fractions using 2,2-Diphenyl-1-picrylhydrazyl radical. *J Agric Chem*. 2000; 48 (3): 648-656.
16. Etekpó DS, N'Gaman-Kouassi CC, Mamyrbekova-Békro JA, Békro YA. Antioxidant profiles of alcoholic tinctures from *Heterotis rodundifolia* (SM.) Jacq. Fél. (Melastomaceae) by DPPH radical trapping. *Europ J Biomed Pharma Sci*. 2018; 10 (5): 39-45.
17. Tanoh SK, N'gaman-Kouassi CC, Boa D, Mamyrbekova-Békro AJ, Békro YA. Activité antioxydante des extraits bruts hydroéthanoliques et hydroacétoniques des organes de quatre plantes de Côte d'Ivoire médicinales. *Nature & Technol*. 2019; 22: 1-7. French.
18. Benzie IF, Strain J. The Ferric Reducing Ability of Plasma (FRAP) as a Measure of Antioxidant Power: The FRAP Assay. *Anal Biochem*. 1996; 239: 70-76.
19. Gong J, Huang J, Xiao G, Chen F, Lee B, You Y. et al. Antioxidant Capacities of Fractions of Bamboo Shaving Extract and Their Antioxidant Components. *Molecules*. 2016; 21(996):1-14.
20. Hunskaar S, Hole K. (1987). The formalin test in mice: dissociation between inflammatory and non-inflammatory pain. *Pain*; 30: 103-114.
21. Ouédraogo N, Lompo M, Sawadogo RW, Tibiri A, Hay AE, Koudou J. et al. Étude des activités anti-inflammatoire, analgésique et antipyrétique des décoctés aqueux des feuilles et des racines de *Pterocarpus erinaceus* Poir. (Fabaceae). *Phytotherapie*. 2012; 10(5): 286-292. French.
22. Westlale WJ. A one-sided version of the Turkey-Duckworth test. *Technometrics*. 1971; 13(4): 901-903.
23. Tambunan AP, Bahtiar A, Tjandrawinata RR. Influence of extraction parameters on the yield, phytochemical, TLC-Densitometric quantification of quercetin, and LC-MS profile, and how to standardize different batches for long term from *Ageratum conyzoides* L. leaves. *Pharmac J*. 2017; 9(6): 767-774.
24. Dahmoune F, Boulekbache L, Moussi K, Aoun O, Spigno G, Madani K. Valorization of *Citrus limon* residues for the recovery of antioxidants: evaluation and optimization of microwave and ultrasound application to solvent extraction. *Indus Crops Prod*. 2013; 50: 77-87.
25. Chew KK, Ng SY, Thoo YY, Khoo MZ, Wan Aida WM, Ho CW. (2011). Effect of ethanol concentration, extraction time and extraction temperature on the recovery of phenolic compounds and antioxidant capacity of *Centella asiatica* extracts. *Inter Food Res*. 2011; 18: 571-578.
26. Tan PW, Tan CP, Ho CW. Effect of extraction solvent system, time and temperature on total phenolic content of henna (*Lawsonia inermis*) stems. *Inter Food Res J*. 2013; 20(6): 3117-3123.
27. Kankara SS, Muskhazli M, Nulit R, Rusea G. Effect of drying methods, solid-solvent ratio, extraction time and extraction temperature on phenolic antioxidants and antioxidant activity of *Guiera senegalensis* J. F. Gmel (Combretaceae) leaves water extract. *Am J Phytomed Clin Therap*. 2014; 2(12): 1378-1392.
28. Ioannou I, Hafsa I, Hamdi S, Charbonnel C, Ghoul M. Review of the effects of food processing and formulation on flavonol and anthocyanin behavior. *J Food Eng*. 2012; 111: 208-217.
29. Wang L, Weller LC. Recent advances in extraction of nutraceuticals from plants. *Food Sci Tech*. 2006; (17): 300-312.

30. Khan MK, Abert-Vian M, Fabiano-Tixier AS, Dangles O, Chemat F. Ultrasound-assisted extraction of polyphenols (flavanone glycosides) from orange (*Citrus sinensis* L.) peel. *Food Chem.* 2010; 119: 851-858.
31. Huang D, Ou B, Prior RL. The chemistry behind antioxidant capacity assays. *J Agric Food Chem.* 2005; 53: 1841-1856.
32. Popovici C, Saykova I, Tylkowski B. Evaluation de l'activité antioxydante des composés phénoliques par la réactivité avec le radical libre DPPH. *Rev Gen Indus.* 2009; 4: 25-39. French
33. Kalia K, Sharma K, Singhi HP, Singh B. Effects of extraction methods on phenolic contents and antioxidant activity in aerial parts of *Potentilla atrosanguinea* Lodd. and quantification of its phenolic constituents by RP-HPLC. *J Agric Food Chem.* 2008; 56: 10129-10134.
34. Mohseni S, Sani AM, Tavakoli M, Raoufi AM. Effect of extraction conditions on antioxidant activities of *Echinops persicus*. *J Essential oil Bearing Plants.* 2017; 20(6): 1633-1644.
35. Hmidani A, Bouhlali EDT, Khouya T., Ramchoun M., Zegzouti Y.F., Benlyas M. et al. Effect of extraction methods on antioxidant and anticoagulant activities of *Thymus atlanticus* aerial part. *Sci Afric J.* 2019; 5: 1-7.
36. Senhaji S, Lamchouri F, Toufik H. Phytochemical content, antibacterial and antioxidant potentiel of endemic plant *Anabasis aretioïdes* Coss. & Moq. (Chenopodiaceae). *Biomed Res Inter.* 2020; 6: 1-16.
37. Ahmed MI, Xu X, Sulieman AA, Mahdi AA, Na Y. Effect of extraction conditions on phenolic compounds and antioxidant properties of koreeb (*Dactyloctenium aegyptium*) seeds flour. *J Food Measur Charact.* 2019; 14: 799-808.
38. Duy NQ, Pham TN, Binh ML, Thuanh M, Van TTN, Lam TD. et al. Effect of extraction conditions on antioxidant activities of Roselle (*Hibiscus sabdariffa* L.) extract. *Materials Sci forum.* 2020; 977: 201-206.
39. Gope PS, Dutta AK, Makhnoon S, Banik S, Siddiquee MA, Kabir Y. Effect of solvent and extraction time on the antioxidant properties of *Citrus macroptera*. *International J Drug Dev Res.* 2014; 6(2): 28-35.
40. Hinneburg I, Dorman HJD, Hiltunen R. Antioxidant activities of extracts from selected culinary herbs and spices. *Food Chem.* 2006; 97(1): 122-129.
41. Omorede IE, Nkeiruka E, Kissinger O. Analytical comparison of the phytochemical composition and antioxidant activity of methanol extracts derived from *Alchornea cordifolia* and *Corchorus olitorius*. *J Phytomed Therap.* 2020; 19(1): 364-374.
42. Jeong SM, Kim SY, Kim DR, Jo SC, Nam KC, Ahn DU. et al. Effects of heat treatment on the antioxidant activity of extracts from citrus peels. *J Agric Food Chem.* 2004; 52: 3389-3393.
43. Kumaran A, Karunakaran RJ. In vitro antioxidant activities of methanol extracts of five *Phyllanthus* species from India. *Lebensmittel-Wissenschaft und Tech.* 2007; 40: 344-352.
44. Vane JR, Botting RM. The mechanism of action of Aspirin. *Thrombosis Res.* 2003; 110 (5-6): 255-258.
45. Ribeiro D, Freitas M, Tomé SM, Silva AMS, Laufer S, Lima JLFC et al. Flavonoids inhibit COX-1 and COX-2 enzymes and cytokine/chemokine production in human whole blood. *Inflammation.* 2014; 38(2): 852-870.
46. Lago JHG, Arruda TCA, Mernak M, Barrosa KH, Martins MA, Tibério LFLC. et al. Structure-activity association of flavonoids in lung diseases. *Molecules.* 2014; 19(3): 3570-3595.