

EFFECT OF ORGANIC, INORGANIC AND BIOFERTILIZERS ON GROWTH, YIELD AND QUALITY OF CLUSTER BEAN (*Cyamopsis tetragonaloba* L.)

ABSTRACT

Cluster bean (*Cyamopsis tetragonaloba* L. Taub] popularly known as "Guar" is an important legume crop mainly grown under rainfed condition in arid and semi-regions of Rajasthan during *kharif* season. The experiment was conducted in the horticulture Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj (UP) During July 2021- October 2021. The experiment was laid out in RBD with 9 Treatments with 3 replications. The results revealed that T8 (N20P50K50 Vermicompost 5 t ha⁻¹ + PSB 5 kg ha⁻¹ + Rhizobium 5 kg ha⁻¹) – performed the best in terms in Plant Height (140.22), Number of Branches (21.39), Length of Pods (25.00), Width of Pods (3.70), Weight of Pods (4.10), Number of cluster per plant (21.60), Number of Pods per plant (89.17), Number of seeds per pod (11.80), Number of Pod yields per plant (196.87), Number of pod yield per plot (68.03), TSS (12) B:C Ratio (2.5:1). Therefore, the treatment T8 (N20P50K50 Vermicompost 5 t ha⁻¹ + PSB 5 kg ha⁻¹ + Rhizobium 5 kg ha⁻¹) is the best when compared to other treatments. As, the highest benefit cost ratio was observed in T8 (N20P50K50 Vermicompost 5 t ha⁻¹ + PSB 5 kg ha⁻¹ + Rhizobium 5 kg ha⁻¹). i.e., (2.5:1) which states that is economically profitable compared to all other treatments.

Keywords: Cluster Bean, Growth, yield, Quality.

INTRODUCTION

Cluster bean (*Cyamopsis tetragonaloba* L. Taub] popularly known as "Guar" is an important legume crop mainly grown under rainfed condition in arid and semi-regions of Rajasthan during *kharif* season. It is very hardy and drought tolerant crop. Its deep penetrating roots enable the plant to utilize available moisture more efficiently and thus offer better scope for rainfed cropping. The crop also survives even at moderate salinity and alkalinity conditions. There is no other legume crop so hardy and drought tolerant as Cluster bean, which is especially suited for soil and climate of Rajasthan. (Kherawat *et al.*, 2013). The pods of Cluster bean are as rich in food value as that of Cluster Bean. According to (Aykroyd 2007), the composition of Cluster bean is 81.0 (g) moisture, 10.8 (g) carbohydrate, 3.2 (g) protein, 1.4 (g) of fat, 1.4 (g) of minerals, 0.09 (mg) thiamine, 0.03 (mg) riboflavin, 47 I.U. vitamin C, 316 I.U. vitamin A (100 g⁻¹) of edible portion. Nitrogen is required for the synthesis of chlorophyll and amino acids, which contribute to the protein building blocks and consequently plant growth. Cluster bean cultivars with nitrogen applied had higher crude protein, crude fibre, ash percentage, carbs, leaf area per plant, dry matter, and green fodder yields (Ayubet *al.*, 2010). Phosphorus is the second most critical nutrient to add to the soil in order to maintain plant growth and crop yield (Singh *et al.*, 2009). It boosts rhizobium activity and promotes the production of root nodules. As a result, it aids in the

Comment [L1]: italic

Comment [L2]: italic

Comment [L3]: Avoid words that are in the title

Comment [L4]: *al.*, 2013

Comment [L5]: Aykroyd (2007)

Comment [L6]: Insert space

Comment [L7]: Delete space

Comment [L8]: Idem L7

Comment [L9]: italic

fixation of more nitrogen from the atmosphere in root nodules. **Handbook of Agriculture by ICAR, (2010).** Potassium benefits ascribed to k include resistance of plant against pest, disease, and stresses caused by drought, salinity, sodicity in assuring improving crop quality characteristics (Kherawat *et al.*, 2013). Farmyard manure the organic manure like FYM is the oldest and cheapest source ornutrient being popular from the ancient times. Application of FYM on field enrich the soil fertility is an old practice unlike chemical fertilizers which contain only one, two or three plant nutrients. FYM seems to act directly for increasing the crop yields either by acceleration of respiratory process with increasing cell permeability and hormonal growth action or by combination of all these processes. FYM provides plants both macro and micronutrients. A well decomposed FYM contains 0.5% N, 0.2% P₂O₅ and 0.5% K₂O. **Handbook of Agriculture by ICAR, (2010).** Vermicompost are products derived from the accelerated biological degradation of organic wastes by earthworms and microorganisms. Earthworms consume and fragment the organic wastes into finer particles by passing them through a grinding gizzard and derive their nourishment from microorganisms that grow upon them. These materials contained mineral contents (% dry weight) ranging from 2.2–3.0 N, 0.4-2.9 P, 1.7-2.5 K, and 1.2-9.5 Ca compared to those of a commercial plant growth (Arancon *et al.*, 2005). *Rhizobium* the use of biofertilizers are more eco-friendly in nature. They can play a significant role in fixing atmospheric nitrogen; biofertilizer enrich soil fertility and improves soil fertility. Of these biofertilizers, *rhizobium* inoculants specific for different leguminous crop in the most important in India. The largest contribution of biological nitrogen fixation to agriculture is derived from the symbiosis between legumes and *Rhizobium* species.

Comment [L10]: delete

Comment [L11]: idem L7

Comment [L12]: 0.5% N, 0.2% P₂O₅ and 0.5% K₂O

Comment [L13]: delete

Comment [L14]: delete

MATERIALS AND METHODS

The experiment was conducted at Experimental Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj (UP) during 2021 during Kharif season in India. The experiment material consist of MDU 1 variety of cluster bean, which is released from Tamil Nadu agricultural university, Tamil Nadu. The soil of the experiment field was alkaline, sandy loam and pH of 7.2. The pit was dug with 30×30×30 cm³ and applied dose of FYM is 20 tonnes/ha and N, P, K is 20 kg, 50kg and 50kg/ha respectively. The experiment was laid out in randomized block design with three replications consisting of 8 treatment combinations, inorganic fertilizers, two kind of organic manures (FYM and vermicompost) and biofertilizers (*Rhizobium* and PSB) alone, some treatments are comprising of organic manures with biofertilizers and different quantity levels of inorganic fertilizers. The plot size was 45cm×30cm spacing rows and plants. Statistical analysis of variance was performed on the data collected throughout the experiment. The observation were recorded for Plant height(cm), Number of Branches per plant, Length of pods, Width of pods, Weight of pods, Number of Cluster per plant, Number of Pods per plant, Number of Seeds per pod, Number of pod yield per plant (g), Number of pod yield per plot (kg), Total soluble solid TSS (°Brix), economics were analysed statistically. The significance of the treatments was determined using the 'F' test at a level of significance of 5%.

Comment [L15]: tonnes ha⁻¹

Comment [L16]: 50kg ha⁻¹

Comment [L17]: italic

Comment [L18]: height (cm)

3. RESULTS AND DISCUSSION

Growth Parameters

The data on growth parameters in different treatment combinations was recorded (Table 1). The maximum height (33.50, 48.83 and 140.22 cm) at 30, 60, 90 DAT was recorded on 80% RDF Chemical fertilizers + 20% Vermicompost + PSB+ **Rhizobium**. While the minimum Plant height recorded in Absolute Control (23.83, 38.00 and 94.30 cm). It might be due to improved soil physical, chemical and biological properties. Higher availability of all plant nutrients resulted in the improved plant characters like plant height. These findings are in conformity with the findings of **Kumar et al. (2004) and Ashwini (2005) in french bean**.

Comment [L19]: italic

Comment [L20]: delete

The maximum Number of Branches (15.50, 17.67 and 21.39) at 30, 60, 90 DAT was recorded on 80% RDF Chemical fertilizers + 20% Vermicompost + PSB+ **Rhizobium**. While the minimum Number of Branches recorded in Absolute Control (6.18, 8.17 and 8.94). Increase in the number of branches per plant due to PSB inoculation may be attributed to the conversion of unavailable phosphorus to available forms particularly during the early crop growth phase which would have helped in the absorption of all major and minor nutrients required for the plant to put forth early vigour in vegetative phase and helps to increase number of branches similar observations were also recorded by **Prasad et al. (2013)** in cowpea.

Comment [L21]: italic

Comment [L22]: delete

Yield Parameters

The maximum Length of Pods (25.00) at 90 DAT was recorded on 80% RDF Chemical fertilizers + 20% Vermicompost + PSB+ **Rhizobium**. While the minimum Length of Pods recorded in Absolute Control (8.83). Increase in Length of pods is due to better assimilation of photosynthates and better portioning into developing pod cluster have taken place and improving yield attributing characters like pod length and diameter. Similar results are in accordance with the **Ashwini (2005) in French Bean**. Width of Pods (3.70) at 90 DAT was recorded on 80% RDF Chemical fertilizers + 20% Vermicompost + PSB+ **Rhizobium**. While the minimum Width of Pods recorded in Absolute Control (2.50). At various amounts of NPK, Vermicompost, PSB, and **Rhizobium**, the mean value of Width of Pods was determined to be significant. Enhanced yield attributing features such as pod width have occurred due to improved photosynthate assimilation and better portioning into growing pod clusters. **Nirmala and Vadivel (1978)** made similar observations in cucumber. Weight of Pods (4.10) at 90 DAT was recorded on 80% RDF Chemical fertilizers + 20% Vermicompost + PSB+ **Rhizobium**. While the minimum Weight of Pods recorded in Absolute Control (2.50). At various amounts of NPK, Vermicompost, PSB, and **Rhizobium**, the mean value of Weight of Pods was determined to be significant. Enhanced yield attributing features such as pod width have occurred due to improved photosynthate assimilation and better portioning into growing pod clusters. **Gandhi and Sivakumar (2010)** made similar observations in cucumber.

Comment [L23]: italic

Comment [L24]: italic

Comment [L25]: italic

Comment [L26]: italic

Comment [L27]: italic

Number of Cluster per plant (21.60) at 90 DAT was recorded on 80% RDF Chemical fertilizers + 20% Vermicompost + PSB+ Rhizobium. While the minimum Number of Cluster per plant recorded in Absolute Control (14.42). PSB could be the greater availability of nutrients in the soil and resulted better growth and development which might be attributed to better mobilization of phosphorus and increased allocation of photosynthates towards the economic parts and also hormonal balance on the plant system Ramana et al., (2011). Number of Pods per plant (89.17) at 90 DAT was recorded on 80% RDF Chemical fertilizers + 20% Vermicompost + PSB+ Rhizobium. While the minimum Number of Pods per plant recorded in Absolute Control (67.40).

Comment [L28]: italic

Comment [L29]: insert space

Comment [L30]: delete

Comment [L31]: italic

The maximum Number of Seeds per pods (11.80) at 90 DAT was recorded on 80% RDF Chemical fertilizers + 20% Vermicompost + PSB+ Rhizobium. While the minimum Number of Seeds per pods recorded in Absolute Control (7.93). The significant improvement in yield attributes of cluster bean was due to fact that rhizobium inoculation increased the root volume through better root development, nodulation, more nutrient availability resulting in vigorous plant growth which in turn results in better flowering, pod formation and increased pod yield. Since, PSB helps in reducing phosphorus fixation by its chelating effect and also solubilized the fixed phosphorus leading to more uptakes of nutrients and reflected in better yields. Similar findings were found by Mishra and Baboo (2002).

Comment [L32]: italic

Comment [L33]: italic

The maximum Number of Pod yields per plant (196.87) at 90 DAT was recorded on 80% RDF Chemical fertilizers + 20% Vermicompost+ PSB+ Rhizobium. While the minimum Number of Pod yields per plant recorded in Absolute Control (156.47). PSB could be due to the greater availability of nutrients in the soil and resulted better growth and development which might be attributed to better mobilization of phosphorus and increased allocation of photosynthates towards the economic parts and also hormonal balance on the plant system These finding are in conformity with the findings of Ramana et al., (2011). The maximum Number of Pod yields per plot (68.03) at 90 DAT was recorded on 80% RDF Chemical fertilizers + 20% Vermicompost + PSB+ Rhizobium. While the minimum Number of Pod yields per plot recorded in Absolute Control (46.47). The Enhancing yield attributing features such pod width have occurred due to improved photosynthate assimilation and better portioning into growing pod clusters. Gandhi and Sivakumar (2010) made similar observations in cucumber.

Comment [L34]: italic

Comment [L35]: delete

Comment [L36]: delete

Comment [L37]: italic

The maximum TSS (Brix) (12) at 90 DAT was recorded on 80% RDF Chemical fertilizers + 20% Vermicompost + PSB+ Rhizobium. While the minimum TSS (Brix) recorded in Absolute Control (8.66). The increased protein content attributed to increase in N content and uptake by the crop due to phosphorus application. Increase of protein content of cluster bean over due to the seed inoculation increased the N content of pods which ultimately reflects TSS content of pods. Similar findings were found by Tripathi and Edward (2017) in guar.

Comment [L38]: italic

Table 1 : Plant characteristics

Comment [L39]: Table 1: Plant characteristics.

Plant Height			Number of Branches			Length of pods	Width of pods	Weight of pods	No. of cluster plant ⁻¹	No. of pods plant ⁻¹	No. of seeds pod ⁻¹	Pod yield plant ⁻¹	Pod yield plot ⁻¹ (kg)	TSS
30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS									
23.83	38	94.3	6.18	8.17	8.94	8.83	2.5	2.5	14.42	67.4	7.93	156.5	46.47	8.66
25.67	41.28	100.7	7.5	10.39	17.99	12.5	2.55	2.55	15.37	71.43	8	169.9	47.27	9.33
27.56	44.33	103.9	8.5	12.39	18.05	12.51	2.59	2.59	16.02	73.22	8.06	176.5	48.43	10.4
26.5	38.07	122.8	7.5	15.06	21	16.75	2.89	3.1	19.39	82.71	9.33	184.6	57.32	10.83
27.5	37.6	122.3	12	15.1	20.13	16	2.8	2.9	18.66	79.13	9.26	180.5	56.64	10.8
28.06	40.63	110.8	7.78	12.22	18.67	14.1	2.6	2.6	16.89	78.42	8.4	178.1	49.47	10.5
26.89	38.39	122.2	6.6	14.94	19.67	15	2.7	2.8	17.15	78.46	8.46	179.1	50.55	10.6
30.83	45.17	130.3	12	15.1	21.39	21	2.9	3.5	19.82	85.34	11.46	192.1	67.33	11
33.5	48.83	140.2	15.5	17.67	21.39	25	3.7	4.1	21.6	89.17	11.8	196.9	68.03	12
S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
0.71	2.31	8.45	0.53	0.78	1.09	1.00	0.18	0.19	1.12	4.63	0.442	8.93	0.36	0.65
2.01	4.40	5.03	4.58	4.76	4.60	4.96	4.91	5.07	4.91	4.78	3.79	3.92	4.51	4.91
0.342	1.111	4.076	0.25	0.376	0.523	0.484	0.084	0.09	0.52	2.23	0.213	4.3	0.17	0.31

Comment [L40]: Inform meaning in the table footer

CONCLUSION

The result from the present investigation concluded that the treatment T₈ which received RDF Chemical fertilizers + 20% Vermicompost + PSB + *Rhizobium* was found superior Plant height, Number of branches, Number of days of first picking, Average fruit width, Fruit yield per plant (kg), Number of seeds per fruit, Average fruit length, Number of clusters plant, Number of pods in a cluster, TSS°(Brix), net income of 84750.00 Rs./ha and B:C ratio of 2.5.

Comment [L41]: italic

Comment [L42]: Rs ha⁻¹

REFERENCES

Comment [L43]: Do a general review of the references, the majority are outside the standards

Anupama, Dolkar, R., Vadodaria, J.R. and Verma, P. (2016). Effect of different varieties of cluster bean (*Cyamopsis tetragonoloba* (L.)Taub.) and biofertilizers on growth, yield and quality parameters. *Ecology, Environment Conservation*, 22: pp. (S457-S462).

Comment [L44]: .

Comment [L45]: italic

Comment [L46]: journal data is missing

Ayub, M., Ali S.A., Tahir M., Tahir sh., Tanveer, A., and Siddiqui, M. H. (2013) Evaluating the role of phosphorus solubilizing bacterial inoculation and phosphorus application on forage yield and quality of Cluster bean (*Cyamopsis tetragonoloba* L.) *International journal of modern agriculture*, 2 (1) ISSN 2305- 7246.

Comment [L47]: .

Comment [L48]: ???

Ayub, M., Khalid, M., Tariq, M., Nadeem, M.A., and Naeem, M., (2011) Effect of different seeding densities and nitrogen levels on growth, forage yield and quality attributes of Cluster bean (*Cyamopsis tetragonoloba* Tuab.) *Journal of Agricultural Technology*. vol.7 (5): 1409-1416.

Comment [L49]: .

Comment [L50]: ???

Ayub, M., Nadeem, M. A., Naeem, M., Tahir, M., Tariq, M. and Ahmad, W. (2012) Effect of different levels of P and K on growth, forage yield and quality of Cluster bean (*Cyamopsis Tetragonolobus* L.) *The journal of animal and plant sciences* 22(2): 2012,479-483.

Comment [L51]: .

Ayub, M., Tahir, M., Nadeem, M. A., Zubair, M. A., Tariq, M., and Ibrahim, M. (2010) Effect of nitrogen applications on growth, forage yield and quality of three Cluster bean varieties. *Pak. j. life soc. Sci.* 8 (2): 111-116.

Arya, M.P.S and Singh, R. (1996). Effect of N, P and k on the nodulation, growth and yield characters of horse gram (*Macrotyloma uniflorum* (Lam) Verdc). *Legume Research*.19(2):65-69

Atiyash R., Edward C., Metzger J, Lee S. and Arancon N. (2002). The influence humic acids derived from earthworms – produced organic waste on plant growth, Bioresource technology, 84, pp7-14

Baliah, T.N. and Muthulakshmi, B. (2017). Effect of microbially enriched vermicompost on growth and biochemical characteristics of okra (*Abelmoschus esculentus* L. Moench) *Advances in Plants and Agriculture Research*, 6(5): 0022

Babiskar, V.S., Damame, H.S., Raj and Shette, P.G (2012). Growth, yield and quality of clusterbean(*Cyamopsis tetragonoloba* L.Taub.) as influenced by organic fertilizers and different levels of sulphur for vegetable purpose. *Crop research*.43(1,2&3):52-54

Comment [L52]: Insert space

Comment [L53]: Insert space

Bhathal, S. and Kumar, R. (2016) Response of integrated nutrient management on growth, yield and yield attributing characters of Cluster bean (*Cyamopsis tetragonaloba* L. TAUB.) under irrigated conditions of Amritsar. *International Journal in Management and Social Science*, vol.04 Issue-05.

Comment [L54]: Insert space

Comment [L55]: Idem L45

Deshmukh, R. P., Nagre, P. K., Wagh, A. P. and Dod, V. N. (2014) Effect of different bio-fertilizers on growth, yield and quality of Cluster bean *Indian Journal of Advances in Plant Research* (IJAPR) 1 (2): -39-42; ISSN: 2347-8918.

Comment [L56]: Delete space

Comment [L57]: .

Comment [L58]: italic

Comment [L59]: ???

Fisher, R.A., (1955) Statistical methods and scientific induction. *Journal of the royal Statistical society series*, 17,69-78.

Comment [L60]: ???

Gireesh, S.P., and Malabasari, T. A. (2014) Effect of major nutrient and picking stage on seed yield and quality of Cluster bean (*Cyamopsis tetragonaloba* L.). *ISSN:2319-9857*

Comment [L61]: journal data is missing

Gandhi, A. and Sivakumar, K. (2010). Impact of vermicompost carrier based bioinoculants on the growth, yield and quality of rice (*Oryza sativa* L.) cv. nlr 145. *The Ecoscan*. 4: 83- 88.

Comment [L62]: ???

Indirabai, W. P. S. and Suja, P. S. R. (2009). Vermicompost of kitchen waste by the earthworm, *lampitoma uritiifor* kitchen garden. *The Ecoscan*. 3: 231-234

Comment [L63]: ????

Iqbal, M., Khan, A. G., Hassan, A. and Islam, K. R. (2013) Tillage and nitrogen fertilization impact on irrigated Cluster bean yields, and soil chemical and physical properties under semiarid climate *Journal of Sustainable Water. Sc. and Mang.* 1(3): 90-98.

Comment [L64]: *Journal of Sustainable Watershed Science and Management*

Comment [L65]: ???

Comment [L66]: .

Comment [L67]: .

Jackson, M. L., (1958) Soil Chemical Analysis Prentice Hall of India Pvt Ltd., New Delhi

Comment [L68]: ???

Kherawat, B. S., Munnalal, Agarwal, M., Yadav, H. K. And Kumar, S. (2013) Effect of applied potassium and manganese on yield and uptake of nutrients by Cluster bean (*Cyamopsis tetragonaloba*). *Journal of Agricultural Physics* 13 (1):22- 2

Comment [L69]: .

Comment [L70]: *Environment and Ecology*

Comment [L71]: ???

Kumawat, P. D. and Khangarot, S. S (2007) Response of sulfur, phosphorus and Rhizobium inoculation on content and uptake of nutrients of Cluster bean [*Cyamopsis tetragonaloba* L. Taub] *Environ and Eco.* 20 (4):803-805

Comment [L72]: ???

Kumar, M., Sinha, K. K. and Roysherma, R. P. (2004). Effect of organic manure, NPK and boron application on the productivity of French bean in sandy loam soil of north Bihar. *Indian Journal of Pulse Research*. 17:42-44.

Meena R., Jat N. L. and Meena N. L. (2002) Effect of phosphorus and biofertilizer (*Rhizobium* and PSB) on yield and quality of Cluster bean (*Cyamopsis tetragonaloba* L. Taub). *Ann. Agric. Res.*, 23(2): 349-351.

Munsell, A. H. (1921) Munsell's description of his colour system, from a lecture to the American Psychological Association. *American journal of psychology* 23(2): 236 244.

- Muthuvel, P., Udayasoorian, C., Natesan, R. and Ramaswamy, P. P. (1992)** Introduction to Soil Analysis, Tamil Nadu Agricultural University Coimbatore-641002
- Olsen, S.R., Cole, S.C. and Dean, L.A. (1954)** Response of available phosphorus in soils by extraction with HNO₃, U.S.D.A. Pp. 939.
- Patel, C. S., Patel, J. B., Sathar, J. V. and Patel, P. M. (2006)** Effect of integrated nutrient management on Cluster bean (*Cyamopsis tetragonaloba* L.) seed production of cv. Pusa navbahar. *International journal of Agricultural research*, 6:206-208.
- Prabhu, T, Narwadkar, P.R. Sajindranath, A.K., Rathod, N. G. (2002).** Integrated nutrient management in coriander. *South Indian Horticulture* ;50(1):680-84.
- Ponnumani, A. and Semmalar, R. (2015).** Effect of Panchagavya on growth parameter analysis on *Capsicum frutescens*. *Species*, 14(44):88-99
- Reager, M. L., Choudhary, G. R. and Dahama, A. (2007)** Effect of weed control and phosphorus on growth and quality of Cluster bean (*Cyamopsis tetragonaloba* L.) *Ann. Agri. Res.* 24 (3):563-56
- Rajkhowa, D.J., Saitya, M. and Rakhowa, K.M. (2003).** Effect of vermicompost and levels of fertilizer on green gram. *Legume Research* ,26(1):63-65.
- Sharma, S. J., Dixit, A. K. and Sharma, A. K. (2011).** Growth and yield of capsicum (*Capsicum annum*) and garden pea (*Pisum sativum*) as influenced by organic manures and bio fertilizers. *Indian Journal of agricultural sciences*: 81:637-6.
- Sajid, M., Ahmed, I., and Rab, A. (2009)** Effect of nitrogen level on the yield and yield component of guar gum (*Cyamopsis tetragonaloba* L.). *Am, Eurasian J. Sustain Agric* 3 (1): 29-32.
- Saravanan, P., Sathish K. S. and Ignesh, A. (2013)** Effect of organic manures and chemical fertilizers on the yield and macronutrient concentrations of green gram. *International Journal of Pharmaceutical Science Invention ISSN (Online)*:2319-6718.
- Sharma, S. K. and Jain, N. K. (2012)** Effect of balanced fertilization on productivity and soil fertility status of Cluster bean. *Legume Research*, 35(1):32-35.
- Singh, B. and Kumar, R. (2016)** Effect of integrated nutrient management on growth, yield and nutrient uptake of Cluster bean (*Cyamopsis tetragonaloba*) under irrigated conditions. *Agric. Sci. Digest*, 36 (1): 35-39.
- Singh, R. (2002)** Effect of integrated nutrient management in Cluster bean (*Cyamopsis tetragonaloba*)- wheat (*Triticum aestivum*) cropping system. under western Rajasthan conditions. *Indian Journal of Agronomy*, 47(1):41-45.

Subbiah, B. V. Asija, and G. L. (1956) A rapid procedure for the estimate of Available nitrogen in soil. *Current Sci.*, 25:259-260.

Takase, M., Sam-Amoah, Owusu, L. K. and Sekyere, J. D. (2011) the effects of four sources of irrigation water on soil chemical and physical properties. *Asian Journal of Plant Sciences* 10:(1) 92-96.18.

Toth, S.J. and Prince, A.L. (1949) Estimation of cation exchange capacity and exchangeable Ca, K and Na content of soil by Flame Photometer technique. *Soil Sci.*, 67: 439-445.

Walkley, A. and Black, I.A. (1947) Critical examination of rapid method for determining organic carbon in soils, effect of variance in digestion conditions and of inorganic soil constituents. *Soil Sei*, 632:251.

Wilcox, L. V. (1950). Electrical conductivity Am. water work Assoc. Yellow River in Northwest China *Chinese J. of Eco-Agri* 18: 5: 917-922.