

Interaction Effect of Phosphorus and Boron on yield components, productivity parameters and quality traits of Lentil (*Lens culinaris* L.)

Abstract

A field trial was conducted on sandy loam soil having low status of organic carbon and accessible nitrogen, medium in accessible phosphorous and high in accessible potassium at pot house of department of Soil Science and Agricultural Chemistry of C.S.A.U.A&T, Kanpur (campus) under Indo-Gangetic Plain zone of Uttar Pradesh, amid Rabi season of 2021-22. The experiment comprised of 16 treatment combinations in Factorial randomized block design with three replications. Lentil variety Lentil Shekhar 2 (KLB-303) was grown with the recommended agronomic practices. On the premise of the comes about exuded from the present investigation, it might be concluded that application of 60 kg P₂O₅ ha⁻¹+ 2.5 kg Boron ha⁻¹ significantly recorded maximum yield components viz. no. of pod plant⁻¹ (71), Seed pod⁻¹ (1.84) and Seed Index (3.23 g). Similarly, among the productivity parameters such as grain yield (15.67 q ha⁻¹), straw yield (28.22 q ha⁻¹), biological yield (43.89 q ha⁻¹) and harvest index (35.70%) was also associated with the application of 60 kg P₂O₅ ha⁻¹+ 2.5 kg Boron ha⁻¹. Among the quality traits maximum protein content (22.40%) and protein yield (351.00 kg ha⁻¹) was also associated with the application of 60 kg P₂O₅ ha⁻¹+ 2.5 kg Boron ha⁻¹. The present experiment clearly points out the significance of balanced use of Phosphorus and Boron nutrients in Lentil for improving the yield components, yield and quality of Lentil crop.

Key Words: Boron, Lentil, Phosphorus, Protein and Yield.

INTRODUCTION

Lentil, like other pulse crops, has a unique position in cropping systems and sustainable agriculture production because it restores soil fertility. The lentil (*Lens culinaris* L. Medik) often known as masoor in India, is one of the most important Rabi season legumes. It's one of the most nutrient-dense cool-season legumes that can also withstand drought. Its origin is South-West Asia and Mediterranean region. It includes between 24 to 26% protein, 1.3% fat, 2.1% minerals, 3.2% fiber, and 57% carbohydrate (Ali *et al.*, 2017 and Singh *et al.*, 2013). It is also rich in Calcium, Iron, Vitamin B, Folate and Niacin. Red (or pink) lentils have a lower fiber content than green lentils (11% rather than 31 %). Lentils, after soybeans and hemp, have the greatest protein content of any plant, at 26% and as a result, they are an essential

element of the diet in many regions of the world, particularly in Nepal and India, where a big vegetarian population exists. Lentil uses atmospheric N₂ to meet its nitrogen needs via biological nitrogen fixing (**Badarneh and Ghawi 1994**). Several factors influence nitrogen fixation in legumes, including rhizobial strains and soil N and P availability. In the symbiotic N₂-fixation mechanism, phosphorus (P) and nitrogen (N) play distinct roles. Indirectly, symbiotic nitrogen fixation has a high P demand because it requires a lot of energy (**Schulze et al., 2006**), and energy-generating metabolism is highly dependent on energy availability (**Israel, 1987**).

Lentil ranks second only to chickpea in terms of total pulse output in India, occupying 1.59 million hectares and contributing 1.6 million tonnes to pulse production and productivity of 1.01 tonne ha⁻¹ (**Anonymous, 2020**). India ranks 1st in the world in respect of production as well as acreage of lentil followed by Turkey. In India Uttar Pradesh has largest acreage and production of lentil. In Uttar Pradesh, the area, production and yield of lentils are 495 thousand hectares, 441 thousand tonnes, and 891 kg/ha (2012-13) respectively. M.P., U.P., Bihar, West Bengal, Haryana, Rajasthan, Punjab, and Assam are the most prominent lentil-growing states.

Among the key nutrients, phosphorus is one of the most important nutrient elements in the production of Lentil in India. **Hussain et al. (2002)** found that the lentil crop responded well to phosphorus fertilizer. Phosphorus is vital in legume agriculture because it promotes grain formation and root growth, as well as nodulation, seed yield, and seed crude protein content (**Singh et al., 2014**). Plants use only a small percentage of the phosphorus provided through synthetic or chemical fertilizer, but a considerable portion of it is converted into insoluble fixed forms, making it inaccessible for crop uptake. Crop phosphorus recovery efficiency is only between 10% to 30% (**Swarup, 2002**). The overall growth and production of the crop can be boosted by increasing the availability of phosphorus.

The production of nodules on legume roots requires the presence of Boron. Lentil reproductive growth, particularly blooming, fruit, and seed set, is more susceptible to Boron shortage. Boron is essential for cell division, as well as the production of pods and seeds. Boron has an effect on N, P, and K absorption, and its lack alters the ideal balance of those three macronutrients. The B treatment had a significant impact on the N and B concentrations of lentil grain, showing that the B played a positive effect in lentil growth and production of proteins. In comparison to most other field crops, legumes have a higher need for Boron.

Boron is essential for glucose transport, cell wall metabolism, cell membrane permeability and stability, and phenol metabolism, with lignin production being the most important. It also helps to improve the quality of goods. Boron was proven to be beneficial in increasing plant growth and yield when applied to the soil or as a foliar spray.

The pulse crops have a vital role in the agricultural economy of India. Application of phosphorus in combination with boron improves the productivity as well as quality traits of lentil. Combined use of chemical fertilizers gave higher yield, owing to adequate availability of nutrients. Therefore, the present study effect of Phosphorous and boron levels on yield components, productivity parameters and quality characteristics of lentil was undertaken.

METHOD AND MATERIAL

2.1 Geographical Situation of Experimental Site:

The field experiment was conducted at pot culture house of the department Soil Science and Agricultural Chemistry of Chandra Shekhar Azad University of Agriculture & Technology, Kanpur for year 2021. Kanpur is located in the Gangetic plains of Central Uttar Pradesh, at a height of around 125.9 meters above sea level, in the central part of Uttar Pradesh's subtropical tract of North India, between latitudes of 25⁰26' to 28⁰58' North and longitudes of 79⁰3' to 80⁰34' East. The average annual rainfall is 816 mm, with most of it falling between the second and first weeks of June and mid-October, with a few showers in the winter.

2.2 Soil of the experimental material

The soil of the experimental field had originated from alluvial deposits. Soil is sandy loam in texture neutral in reaction (pH 7.28), low inorganic carbon (0.45%), available N (180.20 kg ha⁻¹), available P (11.10 kg ha⁻¹), available K (170.02 kg ha⁻¹).

2.3 Layout and Design of the Experiment

The experiment was laid out in Factorial Randomized Block design with three replications. The total numbers of unit plots were 48. The size of a unit plot was 1 m × 1 m. The width of the main irrigation channel is 1 m.

2.4 Details of Treatment

Table 1 : Treatment used for the study

Phosphorous Levels	Symbol	Boron Levels	Symbol
---------------------------	---------------	---------------------	---------------

0 kg ha ⁻¹	P₀	0 kg ha ⁻¹	B₀
30 kg ha ⁻¹	P₃₀	1.0 kg ha ⁻¹	B₁
45 kg ha ⁻¹	P₄₅	1.5 kg ha ⁻¹	B_{1.5}
60 kg ha ⁻¹	P₆₀	2.5 kg ha ⁻¹	B_{2.5}

2.5 Observation recorded

a.) Number of pods plant⁻¹

Number of pods of the five randomly selected plants was counted and their means were computed to express as number of pods plant⁻¹.

b.) Number of seeds pod⁻¹

Number of grains per pod were recorded at harvest by counting the grains of five randomly selected pods from five tagged plants of each plot and average was worked out.

c.) Seed Index (g)

One hundred seed were counted from samples of each plot and their weight was recorded as seed index (g).

d.) Grain yield (kg ha⁻¹)

After threshing and winnowing, the clean grains obtained from the produce of individual net plot, were weighed and weight was recorded as grain yield. The grain yield recorded under each plot was converted into kg per hectare (kg ha⁻¹).

e.) Straw yield (kg ha⁻¹)

Straw yield was obtained by subtracting the grain yield (kg ha⁻¹) from biological yield (Kg ha⁻¹)

f.) Biological yield (kg ha⁻¹)

Above ground harvestable total biomass produce of each plot tied in bundles was weighed with spring balance after complete sun drying. These results obtained in kg plot⁻¹ area were converted in kg ha⁻¹ by multiplying factor.

g.) Harvest Index (%)

It is the ratio expressed in percentage of economic yield in relation to biological yield. It was estimated by the formula proposed by Donald and Hamblin (1976).

$$\text{Harvest index (\%)} = \frac{\text{Economical yield (kg ha}^{-1}\text{)} \times 100}{\text{Biological yield (kg ha}^{-1}\text{)}}$$

RESULT AND DISCUSSION

Yield Attributes

The observation regarding yield attributes viz. no. of pod plant⁻¹, no. of seed pod⁻¹ and seed index (g) were shown in Table 2. Significantly higher number of pod plant⁻¹ (71.00) was observed in treatment T₁₆ with application of phosphorus 60 kg ha⁻¹ and boron 2.5 kg ha⁻¹ and treatment T₁₅ with application of phosphorus 60 kg ha⁻¹ and boron 1.5 kg ha⁻¹ was statistically at par. Minimum number of pod plant⁻¹ (31) was associated with the control treatment where only recommended dose of N & K are applied. Significantly higher number of seed pod⁻¹ (1.84) was observed in treatment T₁₆ with application of phosphorus 60 kg ha⁻¹ and boron 2.5 kg ha⁻¹ and treatment T₁₅ with application of phosphorus 60 kg ha⁻¹ and boron 1.5 kg ha⁻¹ was statistically at par. Minimum number of seed pod⁻¹ (1.54) are associated with the control treatment where no use of phosphorous and boron. Significantly higher seed index (3.22) was observed in treatment T₁₆ with application of phosphorus 60 kg ha⁻¹ and boron 2.5 kg ha⁻¹ and treatment T₁₅ with application of phosphorus 60 kg ha⁻¹ and boron 1.5 kg ha⁻¹ was statistically at par. Minimum seed index (3.02 g) was observed in the control treatment. From the (Table 2) the results were observed that increasing levels of phosphorus from 0 kg ha⁻¹ to 60 kg ha⁻¹

and boron levels from 0 kg ha⁻¹ to 2.5 kg ha⁻¹ has significant effect on all the productivity parameters viz. no. of pod plant⁻¹, no. of seed pod⁻¹ and seed index (g). The consequences of the current investigation are additionally in concurrence with the investigation of **Datta *et al.*, (2013)**, **Chauhan and Raghav (2017)** and **Yumnam *et al.*, (2018)**

Table 2: Effect of yield attributes of lentil as influenced by phosphorous and Boron Levels

Symbol	Treatment combination	No. of Pods plant ⁻¹	No. of Seed Pod ⁻¹	Seed Index (g)
T ₁	P ₀ B ₀	31	1.54	3.02
T ₂	P ₀ B ₁	34	1.57	3.05
T ₃	P ₀ B _{1.5}	36	1.59	3.07
T ₄	P ₀ B _{2.5}	39	1.60	3.08
T ₅	P ₃₀ B ₀	45	1.63	3.11
T ₆	P ₃₀ B ₁	48	1.64	3.13
T ₇	P ₃₀ B _{1.5}	50	1.66	3.14
T ₈	P ₃₀ B _{2.5}	56	1.69	3.16
T ₉	P ₄₅ B ₀	53	1.67	3.15
T ₁₀	P ₄₅ B ₁	58	1.70	3.17
T ₁₁	P ₄₅ B _{1.5}	62	1.71	3.18
T ₁₂	P ₄₅ B _{2.5}	63	1.75	3.19
T ₁₃	P ₆₀ B ₀	60	1.72	3.18
T ₁₄	P ₆₀ B ₁	66	1.78	3.20
T ₁₅	P ₆₀ B _{1.5}	68	1.81	3.22
T ₁₆	P ₆₀ B _{2.5}	71	1.84	3.23
C.D. at 5 %	P	1.42	2.62	0.10
	B	1.42	2.62	0.10
	P×B	2.92	2.83	0.22
S.E(m)±	P	0.49	0.87	0.03
	B	0.49	0.87	0.03
	P×B	0.97	0.94	0.07

Fig. 1: Effect of yield attributes of lentil as influenced by phosphorous and Boron Levels

Productivity Parameters

Significantly higher seed yield (15.67 q ha^{-1}) was observed in treatment T_{16} with application of phosphorus 60 kg ha^{-1} and boron 2.5 kg ha^{-1} and treatment T_{15} with application of phosphorus 60 kg ha^{-1} and boron 1.5 kg ha^{-1} was statistically at par. Significantly higher Stover yield (28.22 q ha^{-1}) was observed in T_{16} with application of phosphorus 60 kg ha^{-1} and boron 2.5 kg ha^{-1} and treatment T_{15} with application of phosphorus 60 kg ha^{-1} and boron 1.5 kg ha^{-1} was statistically at par. Lowest seed yield (8.19 q ha^{-1}), straw yield (20.42 q ha^{-1}) and biological yield (28.61 q ha^{-1}) was associated with the control treatment. Seed yield and stover yield that is biological yield of lentil increases with increase in phosphorus, this increase is due to significant increase in vegetative characters such as no. of pods, seed weight etc. availability of high doses of phosphorus results in higher photosynthetic activity as phosphorus is major constituent of ATP and ATP is utilized in dark reactions of photosynthesis, phosphorus increases production of carbohydrates, sugars, starch, amino acids and proteins, which enhances pod and seed yield these eventually play role in enhancing biological yield. Higher levels of boron resulted in greater uptake of nutrients by seed and stover, boron also enhances chlorophyll content in leaf and thereby bio mass and photosynthates production is increased, which are effectively transferred towards the roots for its development and to provide required energy for nutrient uptake this uptake results in higher biological yields. From the (Table 3) the results were observed that increasing levels

of phosphorus from 0 kg ha⁻¹ to 60 kg ha⁻¹ and boron levels from 0 kg ha⁻¹ to 2.5 kg ha⁻¹ has significant effect on all the productivity parameters viz. seed yield, stover yield, and harvest index. The results of the present investigation are also in agreement with the findings of Navsare *et al.* (2018), Chauhan and Raghav (2017) and Kumar *et al.* (2017).

Table 3: Effect of productivity parameters of lentil as influenced by phosphorous and Boron Levels

Symbol	Treatment combination	Seed Yield (q ha ⁻¹)	Straw Yield (q ha ⁻¹)	Biological Yield (q ha ⁻¹)	Harvest Index (%)
T ₁	P ₀ B ₀	8.19	20.42	28.61	28.62
T ₂	P ₀ B ₁	8.48	20.83	29.31	28.93
T ₃	P ₀ B _{1.5}	8.90	21.10	30.00	29.66
T ₄	P ₀ B _{2.5}	9.22	21.52	30.74	29.99
T ₅	P ₃₀ B ₀	9.89	21.80	31.69	31.20
T ₆	P ₃₀ B ₁	10.66	21.11	31.77	33.55
T ₇	P ₃₀ B _{1.5}	11.02	22.51	33.53	32.86
T ₈	P ₃₀ B _{2.5}	12.10	22.82	34.92	34.65
T ₉	P ₄₅ B ₀	11.56	22.68	34.24	33.76
T ₁₀	P ₄₅ B ₁	12.55	23.05	35.60	35.25
T ₁₁	P ₄₅ B _{1.5}	13.12	23.67	36.79	35.66
T ₁₂	P ₄₅ B _{2.5}	14.22	25.32	39.54	35.96
T ₁₃	P ₆₀ B ₀	13.83	24.82	38.65	35.78
T ₁₄	P ₆₀ B ₁	14.81	25.88	40.69	36.39
T ₁₅	P ₆₀ B _{1.5}	15.08	26.19	41.27	36.53
T ₁₆	P ₆₀ B _{2.5}	15.67	28.22	43.89	35.70
C.D. at 5 %	P	1.12	1.27	1.68	1.49
	B	1.12	1.27	1.68	1.49
	P×B	1.57	2.04	1.86	2.62
S.E(m)±	P	0.37	0.42	0.56	0.49
	B	0.37	0.42	0.56	0.49
	P×B	0.52	0.68	0.62	0.87

Fig. 2: Effect of productivity parameters of lentil as influenced by phosphorous and Boron Levels

Quality Parameters

Protein Content and Protein Yield

The observation regarding quality parameters viz. protein content and protein yield were shown in Table 3. Significantly higher protein content (22.40 %) was observed in treatment T₁₆ with application of phosphorus 60 kg ha⁻¹ and boron 2.5 kg ha⁻¹ and treatment T₁₅ with application of phosphorus 60 kg ha⁻¹ and boron 1.5 kg ha⁻¹ was statistically at par. Lowest protein content (19.00 %) in lentil grain was associated with the control treatment. Significantly higher protein yield (351.00 kg ha⁻¹) was observed in treatment T₁₆ with application of phosphorus 60 kg ha⁻¹ and boron 2.5 kg ha⁻¹ and treatment T₁₅ with application of phosphorus 60 kg ha⁻¹ and boron 1.5 kg ha⁻¹ was statistically at par. Result will also agreement with the findings of **Sentimenla *et al.* (2012)**, **Kalyani and Math (2021)** and **Yadav *et al.*, (2022)**.

Table 4: Effect of quality parameters of lentil as influenced by phosphorous and Boron Levels

Symbol	Treatment combination	Protein Content (%)	Protein Yield (kg ha⁻¹)
T ₁	P ₀ B ₀	19.00	155.61
T ₂	P ₀ B ₁	19.55	165.78
T ₃	P ₀ B _{1.5}	19.76	175.86
T ₄	P ₀ B _{2.5}	19.88	183.29
T ₅	P ₃₀ B ₀	20.08	198.59
T ₆	P ₃₀ B ₁	20.24	215.75
T ₇	P ₃₀ B _{1.5}	20.56	226.57
T ₈	P ₃₀ B _{2.5}	20.92	253.13
T ₉	P ₄₅ B ₀	20.76	239.98
T ₁₀	P ₄₅ B ₁	21.20	266.06
T ₁₁	P ₄₅ B _{1.5}	21.46	281.55
T ₁₂	P ₄₅ B _{2.5}	21.92	311.70
T ₁₃	P ₆₀ B ₀	21.80	301.49
T ₁₄	P ₆₀ B ₁	22.10	327.30
T ₁₅	P ₆₀ B _{1.5}	22.22	335.07
T ₁₆	P ₆₀ B _{2.5}	22.40	351.00
C.D. at 5 %	P	2.62	2.47
	B	2.62	2.47
	P×B	3.10	2.95
S.E(m)±	P	0.87	0.82
	B	0.87	0.82
	P×B	1.03	0.98

Fig. 3: Effect of quality parameters of lentil as influenced by phosphorous and Boron Levels

CONCLUSION

Based on one year of experiment it may be inferred that application of treatment T₁₆ (P 60 kg ha⁻¹ + B 2.5 kg ha⁻¹) gave the highest grain and stover yield value of lentil crop showed good potential for sustainable production. Maximum protein content are also associated with the same treatment combination described as above. It is strongly recommended that farmer of the U.P adopt the dose of T₁₆ [P 60 kg ha⁻¹ + B 2.5 kg ha⁻¹] doses for better crop yield. Similar results also reported by **Padbhusan and Kumar (2014)** and **Karan *et al.* (2014)**

REFERENCES

Ali A., Ahmad B., Hussain I., Ali A. and Shah F. A. (2017), Effect of phosphorus and zinc on yield of lentil. *Pure and Applied Biology*. 6(4):1397-1402.

Anonymous, (2020). Agricultural Statistics at a Glance, Government of India, Ministry of Agriculture Cooperation and Farmers Welfare, Director of Economics and Statistics.

Badarneh, D. M. D. and Ghawi, I. O. (1994), Effectiveness of inoculation on biological nitrogen fixation and water consumption by lentil under rainfed conditions. *Soil Biology and Biochemistry*, 26 (1):1-5.

Chauhan and Raghav. (2017), Effect of phosphorus and phosphate solubilizing bacteria on growth, yield and quality of chickpea (*Cicer arietinum*) *Annals of Plant and Soil Research* 19(3): 303-306.

Datta S. K., Sarkar M. A. R. and Uddin F. M. J. (2013), Effect of variety and level of phosphorus on the yield and yield components of lentil. *International Journal of Agricultural Research Innovation and Technology*. 3(1): 78-82.

Hussain, M., Shah, S. H. and Nazir, M. S. (2002), Differential genotypic response to phosphorus application in lentil (*Lens culinaris Medic*). *Pakistan Journal of Agricultural Sciences*, 39: 193-196.

Israel DW. (1987), Investigations of the role of phosphorus in symbiotic dinitrogen fixation. *Plant Physiology* 84: 835–840.

Kalyani, M. S. R., and Math, G. (2021), Quality and economics of cowpea as influenced by genotypes, phosphorus levels and liquid based PSB. *International Journal of Chemical Studies*, 9(1): 2796-2800.

Kumar, P., Kumar, R., Mohan, B., Yadav, S. K., Singh, S. and Singh, B. (2017), Effect of foliar and soil application of micronutrients and use of PSB on productivity and water use efficiency of chickpea under rain fed conditions. *International Journal of Current Microbiology and Applied Sciences* 6(8): 3074-3080.

Karan, D., & Singh, S. B. (2014). Effect of zinc and boron application on yield of lentil and nutrient balance in the soil under Indo-Gangetic Plain zones. *Journal of Agri Search*, 1(4).

Navsare R. I., Mane S. S. and Supekar, S. J. (2018), Effect of potassium and zinc solubilizing microorganism on growth, yield and quality of mungbean. *International Journal of Chemical Studies*. 6(1): 1996-2000.

Padbhushan, R. and Kumar, D. (2014), Influence of soil and foliar applied boron on green gram in calcareous soils. *Int. J. Agric. Environ. Biotechnology*. 7(1): 129-136.

Schulze, J., Temple, G., Temple, S. J., Beschow, H. and Vance, C. P. (2006), Nitrogen fixation by white lupin under phosphorus deficiency. *Annals of Botany*, 98 (4): 731-740.

Sentimenla, Singh, A.K. and Singh, S. (2012), Response of soybean to phosphorous and boron fertilization in acidic upland soils of Nagaland. *J. Indian Soc. Soil Sci.* 60: 167-170.

Singh, S., Gupta, G., Khare, E., Behal, K. K. and Arora, N. K. (2014), Phosphate solubilizing rhizobia promote the growth of chickpea under buffering conditions. *Int. J. Pure Appl. Bio-science*, 2(5): 97-106.

Swarup, A. (2002), Lessons from long term fertilizer experiments in improving fertilizer use efficiency and crop yields. *Fertilizer News*, 47 (12): 59-73.

Yadav, P., Yadav, D. D., Pandey, H. P., Yadav, A., Sachan, R., & Yadav, S. (2022). Effect of Fertility Levels and Biofertilizers on Growth Parameters, Root Architecture and Quality of Chickpea (*Cicer arietinum* L.).

Yumnam T., Luikham E., and Singh A. H, (2018), Influence of phosphorus on growth and yield of promising varieties of lentil (*Lens culinaris* L. Medik). *International Journal of Current Microbiology and Applied Sciences*. 7(8): 162- 170.