

Original Research Article

Efficacy of capsaicin rich botanicals on the aphid population in Lablab (*Dolichos lablab* L.) under organic production system

Abstract

Aim: The growing concern for environmental safety and the demand for pesticide residue –free food worldwide have evoked interest in pest management through the use of botanicals, which offers a good alternative to manage the insect pests in an eco-friendly manner. This research was carried out to study the efficiency of botanicals on the aphid population in lablab under organic production system.

Study Design: The study was carried out in Randomized Block Design (RBD) with eight treatments and three replications.

Place and Duration of Study: The study was carried out at College Orchard, Department of Vegetable Science, Horticultural College and Research Institute, Tamil Nadu Agricultural University, Coimbatore during 2021 - 2022.

Methodology: The study was performed using chilli extract with three concentrations (2.5, 5.0 and 7.5%) and compared with other botanicals such as neem oil, neem seed kernel extract, garlic extract and panchagavya. The aphid population was counted visually before and 1, 3, 5 and 7 days after spraying.

Results: The results showed that after 7 days spraying 95.62 per cent of aphid population was reduced in T2 (5 % chilli extract) followed by T3 & T1 (7.5 % & 2.5%) with a population reduction to the level of 95.04 and 91.54 per cent, respectively. This was followed by neem oil with a reduction of 83.50 per cent followed by garlic extract (79.74 %), neem extract (76.94 %) and panchagavya (62.81 %).

Conclusion: The application of chilli extract resulted in reduction of aphid population in lablab under organic cultivation. The chilli extract is non-toxic to humans and does not harm the environment and hence this can be utilized as an alternative to chemical pesticides.

Keywords: Lablab, Chilli extract powder, Capsaicin, Aphid, Botanicals, Pest population

Introduction

Indian bean (*Dolichos lablab* L.; Family: Fabaceae), also known as the Australian pea, Egyptian bean, kidney bean, dolichos bean, Lablab bean, seim bean, or chicharo is grown in Asia and Africa. It is a perennial herbaceous plant that plays a significant role in both field and home garden-grown fruit and vegetable crops. While the dried seeds are utilized in a variety of vegetable food recipes, the green pods are the main reason it is farmed. It is one of the main sources of minerals, dietary fiber, and proteins.

Pest insects have a significant negative impact on Indian bean yield. Numerous insect pests attack the crop, including the aphid *Aphis craccivora* Koch., jassids *Empoasca fabae* (Harris), *E. krameri* Ross & Moore, and *E. kerri* Pruthi; pod borer *Etiella zinckenella* (Treit.); white fly *Bemisia tabaci* (Genn.); stem fly *Ophiomyia phaseoli* (Tryon). Among them, aphids (*Aphis craccivora* Koch.), jassids (*Empoasca fabae* Harris.), and whiteflies (*Bemisia tabaci* Genn.) are three of the most prevalent sucking pests infesting Indian beans in Rajasthan's semi-arid climate. By sucking the cell sap from the vulnerable parts of the plant as well as the lower region of the leaves, both nymphs and adults harm the environment. When plants are severely infested, these pests damage every part of the plant, including the pods, which stunts growth and reduces output.

Many researchers have advocated using chemicals to control sucking pests in order to effectively reduce their population [1,2,3]. However, because the continuous and widespread use of insecticides from the same class has led to issues with resistance, negative effects on parasites and predators, residue hazards for people and animals, and environmental pollution [4]. One of the most effective control measures for synthetic chemical hazards is the use of plant-derived products. Plant products are becoming increasingly popular due to their biodegradability, low persistence and toxicity to non-target organisms, low cost, and ease of availability. There are currently about 200 plants known to have insecticidal properties. Azadirachtin, an active compound extracted from the *Azadirachta indica* A. Juss (neem) tree (Family Meliaceae) whose antiviral, antifungal, antibacterial, and insecticidal properties have been known for several years, is one of the most promising natural compounds[5]. Capsaicinoids (capsaicin, dihydrocapsaicin, nordihydrocapsaicin, homocapsaicin, and homodihydrocapsaicin) are a class of phenolic alkaloids present in chilli. Capsaicin is the active ingredient in chilli peppers and the most abundant irritant compound in hot peppers that causes humans to experience a burning sensation [6]. Capsaicin has broad-spectrum insecticidal activity against many species of insects, eg, stored product beetles (*Sitophilus zeamais* and *Tribolium castaneum*), rice grain insects (*Sitotroga cerealella*), Alfalfa weevil, *Myzus persicae*, *Bemisia tabaci* and *Plutella xylostella* [7].

Materials and Methods

The experiment was conducted at the College Orchard, Department of Vegetable science, Horticultural College and Research Institute, Tamil Nadu Agricultural University, Coimbatore. The experimental plot was ploughed 2-3 times and leveled thoroughly. During the last ploughing adequate amount of FYM were applied to the field and raised beds were prepared. Irrigation was given through drip.

The seeds were sown directly in the mainfield with a spacing of 60 × 30 cm. Recommended package of practices were followed to raise the crop except plant protective measures. Experiment was laid out in Randomized Block Design (RBD) with eight treatments and replicated thrice.

Table 1. Treatment details

| Treatments | | Concentration (%) |
|-------------------|--------------------------|--------------------------|
| T 1 | Chilli extract powder | 2.5 |
| T 2 | Chilli extract powder | 5.0 |
| T 3 | Chilli extract powder | 7.5 |
| T 4 | Neem oil | 0.5 |
| T 5 | Neem seed kernel extract | 5 |
| T 6 | Garlic extract | 3 |
| T 7 | Panchagavya | 5 |
| T 8 | Control | Untreated |

Collection and processing of plant powders

The chilli fruits having high capsaicin content (more than 1 lakh SHU) collected from the TNAU orchard, Horticultural College and Research Institute, Tamil Nadu Agricultural University, Coimbatore were dried under shade. The dried fruits were ground into powder.

Preparation of chilli extracts

The chilli powder weighed at different concentration viz., 2.5, 5.0 and 7.5 per cent and was mixed with 100ml of water. The solution was shaken for an hour in an electric shaker. Sticking agent was added to the extract and sprayed in the field

Preparation of neem seed kernel extract

Fifty grams of well pulverized neem seeds were soaked in a little amount of water overnight. One litre of suspension was produced by filtering it through a muslin cloth with additional water. Three ml of sticking agent per litre was added to the extract and carefully mixed. The extract was immediately applied to the plot.

Preparation of neem oil emulsion

The neem oil emulsion was made by thoroughly mixing the neem oil (0.5 ml) and sticking agent (1 ml). Water (1litre) was added to the mixture and thoroughly mixed.

Preparation of garlic extract

Garlic bulbs were first skinned and thoroughly mixed in sterile distilled water (1:1). Supernatant was collected after centrifugation and passed through a No. 1 Whatman filter paper and then a micropore filter (0.45 µm). The solution thus obtained was stored in a refrigerator (4°C) until use [8].

Preparation of panchagavya

Panchagavya is made up of nine ingredients such as cow dung, cow urine, milk, curd, jaggery, ghee, banana, tender coconut and water. In a wide mouthed mud vessel 5 kg of Cow dung and 500 gm of cow ghee were mixed thoroughly and kept for 3 days. After 3 days the following ingredients were added and kept for 19 days with regular mixing both in the morning and evening hours daily. The container was always kept covered with a mosquito net or cotton cloth [9].

1. Fresh cows urine – 3 liters
2. Cow's milk – 2 liters
3. Cow's curd – 2 liters
4. Jaggery – 500 gms
5. Water – 3 liters
6. Ripe banana fruit – 1 bunch
7. Tender coconut water – 3 liters

Monitoring of pests

After the seedlings emerged, the plants were checked weekly to see if there were any aphid infestations. After the infestation had occurred, the plant extracts were applied. A hand sprayer was used to apply the extracts.

Observation on aphids

Observation on population of aphids was recorded in the morning time in the shoots of five randomly selected plants of each plot. The population was recorded just before treatment and one, three and seven days after the spray.

Statistical analysis

The per cent reduction of aphid population over control was calculated based on field observations [10].

$$\text{Per cent reduction in pest population} = \left(1 - \frac{Ta \times Cb}{Tb \times Ca}\right) \times 100$$

Where,

Ta = Population in treated plots after treatment

Tb = Population in treated plots before treatment

Ca = Population in untreated plots after treatment

Cb = Population in untreated plots before treatment

The data were statistically analysed with ANOVA and means were separated using Least Significant Difference (LSD) test by using AGRES software package (version 7.01).

Results

The sucking pest infestation allowed naturally in lablab during the period of study was aphids (*Aphis craccivora*). The population of aphids ranged from 245.75 to 487.68 numbers per 5 plants before the application of botanicals. (Table 2)

First day after spraying

The aphid population in different treatments ranged between 129.42 and 222.17 after first day of treatment. The per cent reduction was found to be significantly different among the treatments. The reduction due to the application of chilli extract ranged from 44.45 to 54.68 per cent on first day after treatment. The maximum reduction was observed in 5 % chilli extract (54.68 per cent) followed by 7.5 and 2.5 % with 53.32 and 44.45 per cent respectively. This was followed by neem oil treatment with a reduction of 30.13 per cent followed by garlic extract 28.32 per cent. The minimum reduction was observed in panchagavya 21.69 per cent (Table 3).

Third day after spraying

The aphid population in different treatments ranged between 54.08 and 144 third day after treatment. The per cent reduction was found to be significantly different among the treatments. The reduction due to the application of chilli extract ranged from 77.56 to 80.68 per cent on third after treatment. The maximum reduction was observed in 5 % chilli extract (80.68 per cent) followed by 7.5 and 2.5 % with 79.73 and 77.56 per cent respectively. This was followed by neem oil with a reduction of 66.26 per cent followed by garlic extract 62.61 per cent. The minimum reduction was observed in panchagavya 48.84 per cent (Table 3).

Fifth day after spraying

The aphid population in different treatments ranged between 31.08 and 144.16 fifth day after treatment. The per cent reduction was found to be significantly different among the treatments. The reduction due to the application of chilli extract ranged from 83.74 to 89.43 per

cent on fifth day after treatment. The maximum reduction was observed in 5 % chilli extract (89.43 per cent) followed by 7.5 and 2.5 % with 83.74 and 86.94 per cent respectively. This was followed by neem oil treatment with a reduction of 68.46 per cent followed by garlic extract 72.63 per cent. The minimum reduction was observed in panchagavya 51.92 per cent (Table 3).

Seventh day after spraying

The aphid population in different treatments ranged between 11.91 and 106.75 seventh day after treatment. The per cent reduction was found to be significantly different among the treatments. The reduction due to the application of chilli extract ranged from 91.54 to 95.62 per cent on seventh day after treatment. The maximum reduction was observed in 5 % chilli extract (95.62 per cent) followed by 7.5 and 2.5 % with 95.04 and 91.54 per cent respectively. This was followed by neem oil treatment with a reduction of 83.50 per cent followed by garlic extract 79.74 percent. The minimum reduction was observed in panchagavya 62.81 per cent (Table 3).

Discussion

The result of this study established the potential of chilli extract, neem product, garlic extract and panchagavya to control *A.craccivora* on lablab. The chilli extract was significantly effective in reducing population of *A.craccivora* with the maximum reduction observed in chilli extract powder (5 %) on seventh day after treatment. Chilli extract was effective in reducing the survival of aphids. Capsaicin is responsible for the pungent principle in chilli. This might be responsible for the reduction in aphid population. Capsaicin has repellent and insecticidal properties, for example, against hemipterans [11,12]. From the results, the chilli extract showed highest efficiency at controlling aphids. The chilli extract has the ability to kill aphids and mites [13].

The efficacy of neem oil was very close to chilli extract against *A. craccivora* on lablab plants. The neem oil against jassids at 3 % reduced about 88.48 after 7 days after treatment [14]. The neem has controlled the aphids through its multiplicity of action viz., repellency, antifeedency and ovipositional deterrence [15].

The garlic extract also exhibited prominent efficacy after the neem oil. Garlic extract was the most effective at reducing aphid population by 90.98 per cent as well as leafhopper and plant hopper population by a mean percentage of 68.09 per cent [16]. Garlic is known to have anti-feedent, insecticidal and repellent properties [17] which accounted for reduction of aphid population on the garlic extract sprayed plants.

In the present study, the chilli extracts were observed to be more effective against Aphids (*A. craccivora*) followed by neem oil and garlic extract. From the results, the aphid population was reduced seven days after application of chilli extract. These showed that, aphids can be managed with this treatment.

Conclusion

The present study concluded that the plant extracts were effective against aphids. Among the different botanicals used, the chilli extract was found to be very effective and compatible in reducing aphid population in lablab. Five per cent chilli extract concentration was found to be highly effective against aphids after seven days of spraying. It may be concluded that the chilli extract can be used in controlling aphid population in lablab which is very serious sucking pest. Thus, these botanicals can be used as a better alternative to conventional insecticides for the management of pest attack. Since botanicals are environment friendly and non-toxic to human.

Table 2. Aphid population in lablab before and after spraying of botanicals

| Treatments | Mean number of aphid population | | | | |
|--------------------------|---------------------------------|--------------------------|--------------------------|--------------------------|----------------------------|
| | Before spraying | First day after spraying | Third day after spraying | Fifth day after spraying | Seventh day after spraying |
| Chilli extract(2.5%) | 249.5 | 142.58 | 54.08 | 43 | 21.33 |
| Chilli extract(5%) | 279.58 | 129.42 | 55.08 | 31.08 | 11.92 |
| Chilli extract(7.5%) | 460.58 | 222.17 | 97.92 | 65.83 | 23.17 |
| Neem oil | 292.83 | 187.75 | 105.98 | 85.17 | 51.58 |
| Neem seed kernel extract | 215.75 | 161.92 | 101.33 | 85.83 | 59.92 |
| Garlic extract | 280.5 | 174.08 | 113.33 | 86.58 | 63.33 |
| Panchagavya | 277.17 | 200.25 | 144 | 144.17 | 106.75 |
| Control | 390.33 | 401.33 | 411 | 442.42 | 459.33 |

Table 3. Per cent reduction of aphid population after spraying of botanicals

| Treatments | Per cent reduction of aphid population | | | |
|--------------------------|--|--------------------------|--------------------------|----------------------------|
| | First day after spraying | Third day after spraying | Fifth day after spraying | Seventh day after spraying |
| Chilli extract(2.5%) | 44.45 | 77.57 | 83.74 | 91.54 |
| Chilli extract(5%) | 54.69 | 80.68 | 89.44 | 95.62 |
| Chilli extract(7.5%) | 53.32 | 79.74 | 86.94 | 95.04 |
| Neem oil | 28.32 | 66.27 | 75.02 | 83.50 |
| Neem seed kernel extract | 26.40 | 60.05 | 68.47 | 76.94 |
| Garlic extract | 30.10 | 62.61 | 72.63 | 79.74 |
| Panchagavya | 21.69 | 48.85 | 51.92 | 62.81 |
| SED | 10.91 | 2.62 | 2.90 | 2.91 |
| CD(.05) | 23.77 | 5.70 | 6.33 | 6.34 |
| CD(.01) | 33.32 | 7.99 | 8.87 | 8.89 |

| | | | | |
|-----|-------|------|------|------|
| CV% | 36.11 | 4.72 | 4.71 | 4.26 |
|-----|-------|------|------|------|

SED- Standard error of a difference

CD- Critical Difference

CV- Coefficient of Variation

References

1. Garhwal SN, Verma SK, and Sharma JK. Field efficacy of different insecticide against cowpea aphid, *Aphis craccivora* Koch. Annals of Arid Zone. 1994; 33(2):159-160.
2. Dhamaniya B, Sharma JK and Kumawat KC. Bio-efficacy of insecticides against sucking insect pests of moth bean, *Vigna acontifolia*. Annals of plant protection sciences. 2005; 91-93
3. Yadav SR, Kumawat KC and Khinchi SK. Efficacy of new insecticide molecules and bioagents against sucking insect pests of cluster bean, *Cyamopsis tetragonolaba* (Linn). Journal of Protection and Environment. 2011; 115-122.
4. Sande D, Mullen J, Wetzstein M and Houston J. Environmental impacts from pesticide use: a case study of soil fumigation in Florida tomato production. *International journal of environmental research and public health*. 2011; 8(12):4649-4661.
5. Debashri M and Tamal M. A Review on efficacy of *Azadirachta indica* A. Juss based biopesticides: An Indian perspective. *Res. J. Recent Sci*. 2012; 2277:2502.
6. Antonious GF, Meyer JE and Snyder JC. Toxicity and repellency of hot pepper extracts to spider mite, *Tetranychus urticae* Koch. Journal of Environmental Science and Health, Part B. 2006; 41(8):1383-91.
7. Li B, Yang M, Shi R and Ye M. Insecticidal activity of natural capsaicinoids against several agricultural insects. *Natural Product Communications*.2019; 14(7):1934578X19862695.
8. Mobki M, Safavi SA, Safaralizadeh MH and Panahi O. Toxicity and repellency of garlic (*Allium sativum* L.) extract grown in Iran against *Tribolium castaneum* (Herbst) larvae and adults. *Archives of Phytopathology and Plant Protection*.2014; 47(1):59-68.
9. Sailaja V, Ragini NN, Kumar KD, Reddy BR and Satyanarayana SV. Effect of foliar application of Panchagavya on growth and development of leafy vegetable *Spinacia oleracea*. *Int J Agric Food Sci*.2014; 4(4):119-122.
10. Henderson CF and Tilton EW. Tests with acaricides against the brown wheat mite. *Journal of economic entomology*.1955; 48(2):157-161.
11. Bergmann EJ and Raupp MJ. Efficacies of common ready to use insecticides against *Halyomorpha halys* (Hemiptera: Pentatomidae). *Florida Entomologist*.2014; 97(2):791-800.
12. Dayan FE, Cantrell CL and Duke SO. Natural products in crop protection. *Bioorganic & medicinal chemistry*.2009; 17(12):4022-4034.

13. Tomita M and Endo H. Using Capsaicin as a Less Toxic Insecticide©. In Combined Proceedings International Plant Propagators' Society.2007; 57:728-732.
14. Dhiloo KH, Zhang YJ, Rizwan S, Ursani TJ, Chandio JI and Sidhoo M. (2016). Efficacy of different neem oil concentrations against jassid on eggplant under field conditions. European Academic Research.2016; 3(2):12170-12179.
15. Baidoo PK, Baidoe-Ansah D and Agbonu I. Effects of neem (*Azadirachta indica* A. Juss) products on *Aphis craccivora* and its predator *Harmonia axyridis* on cowpea. *American Journal of Experimental Agriculture*.2012; 2(2):198-206.
16. Khodeir IA, El-Dakhakhni TN and Youssef AE. Effect of Garlic and Eucalyptus oils in comparison to Organophosphatinsecticides against some Piercing-Sucking Faba bean insect Pests and natural enemies' populations. Egyptian Academic Journal of Biological Sciences, F. Toxicology & Pest Control.2013; 5(2):21-27.
17. Vijayalakshmi K, Subhashini B and Koul S. Plant in pests control Garlic and Onion. Centre for Indian knowledge system, Chennai, India.1999; 123.