

# Sub-soil nitrogen content as influenced by long-term manuring and its relationship with nitrogen availability and productivity of a rice-rice cropping system in Eastern India

## ABSTRACT

Investigation was made to study the impact of long-term fertilizer and manure application on the sub-soil nitrogen (N) fertility of an acidic *Inceptisol* under continuous rice-rice cropping system. For this purpose, a long-term fertilizer experiment commenced from 2005-06, *rabi* season in the Central Farm of Odisha University of Agriculture and Technology (OUAT), Bhubaneswar under aegis of ICAR, New Delhi was used. The experiment had 12 manurial treatments whose impact on sub-soil N fertility up to 75 cm depth and its relation with surface soil N fertility, N uptake and crop yield has been assessed. has been assessed after 20 cropping cycles. The initial soil was acidic (pH 5.8) with low soil organic carbon ( $4.3 \text{ g kg}^{-1}$ ) and CEC of  $3.75 \text{ cmol (p}^+) \text{ kg}^{-1}$ , mineralizable N ( $187 \text{ kg ha}^{-1}$ ) and total N ( $1280 \text{ kg ha}^{-1}$ ). After 20 cropping cycles, the soil organic carbon (SOC) content increased in all the fertilized treatments in the surface layer and with depth it decreased sharply from 15-30 cm to 30-45 cm layer. Treatments with balanced nutrition registered higher content of total N, organic N, mineralizable N and inorganic N both in surface soil and sub-soil. Not only the surface soil but also the sub-soil contents of total N, organic N and mineralizable N are significantly influenced by long-term use of fertilizer nutrients and manure under wet land intensive rice production system. Sub-soil contents of total N, organic N, mineralizable N, inorganic N, ammonium N and nitrate N have strong positive correlation with those of surface layer. SOC, total N mineralizable N and inorganic N are strongly correlated with each other in the same layer. Correlation of organic N and total N with nitrate N was however significant up to 30 cm depth and beyond that there was no correlation at 30-45 cm layer. From the results, it is concluded that application of nutrient in balanced and integrated manner (NPK + FYM) not only increased grain yield and sustainability of rice but also improved various pools of N in surface as well as sub-soil up to 75 cm depth under wet land rice-rice production system under subtropical climatic situation.

*Key words: Long-term manuring, rice-rice system, sub-soil nitrogen, productivity, sustainability,*

## 1. Introduction

Nitrogen is the one of most limiting nutrients for rice production in rice-rice cropping system. Rice crop requires large amount of N for its growth, development and grain production. Generally, rice plant removes around 14-20 kg N to produce one tonne of rough rice including straw [1]. The amount of plant mineralizable N is positively influenced by N fertilization, mineralization of soil organic matter, biological N fixation and by precipitation. Negative influences result from immobilization, crop uptake and removal, denitrification (and to some extent nitrification), volatilization, leaching, run-off and erosion [2]. The relative importance of these processes depends on environmental variables such as soil pH, top soil texture, soil profile characteristics, soil aeration, water supply and temperature, as well as human

activities such as type, amount, placement and timing of N fertilizers, crop residue management, tillage, soil compaction, drainage, irrigation and land use change.

Soil nitrogen status is one of the key parameters to assess the fertility status of soil. In most soils, N is predominantly organic in nature and only a small portion of total N is present in inorganic form. Soil organic N is the largest source of plant mineralizable N for rice representing 50 to 80% of total N assimilated by crop [3]. Soil organic N (SON) plays a key role in terms of plant nutrition through direct and indirect effects on microbial activity and nutrient availability [4]. Continuous N input increases soil total nitrogen (STN) content in the profile. Significantly higher total N content was reported in the surface soil with organic or organic–inorganic combined fertilization [5]. Continuous cropping and long-term nutrient management have influence on carbon and nitrogen content of soil of different layers and its availability to crops [6]. So far studies on relationship of soil nitrogen with crop yield are restricted to surface soil or maximum up to 30-45 cm depth. Lower layers of greater depth may also influence the surface soil nitrogen fertility and crop yield as there exists connectivity of layers because of submergence, linkage through leaching and percolation, capillary rise of soil solution, presence of oxidised and reduced zones at different layers under rice-rice system. No studies have yet been done on the content of various mineralizable pools of nitrogen in lower layers up to 75 cm or more and their relationship with upper surface soil nitrogen fertility, crop yield and sustainability. So, this investigation was conducted using 10-year-old long-term fertilizer experiment (ICAR, New Delhi) in order to get more information on the impact of repeated use of various inorganic fertilizer nutrients and organic manure on contents N of various mineralizable forms of nitrogen in sub-soil up to 75 cm and study their relationship with the contents of surface soil and crop productivity.

## **2. Materials and methods**

### **2.1 Experimental Site**

The study was conducted during *kharif*, 2015 and *rabi*, 2015-16 in the ongoing experimental field of All India Coordinated Research Project (AICRP) on Long-Term Fertilizer Experiment (LTFE) of ICAR at OUAT, Bhubaneswar, India (20°17' N, 85°49' E and 30 m above mean sea level). The location of the experimental site is characterized as sub-humid sub-tropical climate with *rabi* season from October to June and *kharif* season from July to September. The average annual rain fall is 1453 mm and the mean maximum and minimum temperatures are 31.40°C and 21.10°C, respectively. The experimental soil is a pale yellow (10YR6/8), lateritic *Inceptisol*. The initial soil properties of 0-15cm layer were sandy loam texture with clay 17%, silt 12% and sand 71%, bulk density 1.55 kg m<sup>-3</sup>, cation exchange capacity 3.75 cmol (p<sup>+</sup>) kg<sup>-1</sup>, pH 5.8, electrical conductivity 0.12 dSm<sup>-1</sup>, organic carbon 4.3 g kg<sup>-1</sup>, mineralizable N (Alkaline KMnO<sub>4</sub>)187, available P (Olsen's)19.4 and available K (NH<sub>4</sub>OAc) 43.4 kg ha<sup>-1</sup>, respectively. The DTPA extractable Fe, Mn, Zn and Cu were 33.0, 7.53, 1.80 and 3.15 mg kg<sup>-1</sup>, respectively. Exchangeable cations Ca and Mg were 2.25cmol (p<sup>+</sup>) kg<sup>-1</sup> and 0.65cmol (p<sup>+</sup>) kg<sup>-1</sup>. The hot water-soluble boron was 0.46 mg kg<sup>-1</sup>.

### **2.2 Experiment details**

The experiment consisted of 12 treatments; T<sub>1</sub>-100% PK, T<sub>2</sub> - 100% NPK, T<sub>3</sub>-150%NPK, T<sub>4</sub>- 100%NPK+Zn, T<sub>5</sub>-100%NPK+ FYM, T<sub>6</sub>-100%NPK+Lime+FYM, T<sub>7</sub>-100%NPK+B+Zn, T<sub>8</sub>-100%NPK+S+Zn, T<sub>9</sub>-100%N, T<sub>10</sub>-100%NP, T<sub>11</sub>-100%NPK+Lime and T<sub>12</sub>- Control, where 100% NPK correspond to 80-40-60 kg of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O ha<sup>-1</sup>. The experiment was laid out in randomized block design (RBD) with four replications. Rice cultivar Swarna (MTU 7029) was grown in *kharif*

season and Lalat in *rabi* season of every year. Twenty-five days old rice seedlings were transplanted at a spacing of 20 cm × 10 cm with 2-3 seedlings per hill to puddled field in both the seasons. Nitrogen (N) was applied in three splits i.e. 25% at puddling as basal, 50% topdressing at 18 days after transplanting and 25% topdressed at panicle initiation stage. Entire dose of phosphorus (P) was applied during puddling as basal and potassium (K) was applied in two splits, 50% at puddling as basal and 50% topdressing at panicle initiation (PI) stage. Entire FYM (5 t ha<sup>-1</sup>season<sup>-1</sup>) was applied at the time of puddling. FYM has been added @ 5 t ha<sup>-1</sup> in each season in T<sub>5</sub> and T<sub>6</sub>. Lime @ 1 t ha<sup>-1</sup> in each season has been applied in T<sub>6</sub> and T<sub>11</sub> at the time of land preparation. Zn has been applied as Zinc oxide 0.4% solution seedling root dipping in T<sub>4</sub>, T<sub>7</sub> and T<sub>8</sub>. Borax was foliar sprayed twice as a source of boron @0.25% solution in T<sub>7</sub>. Gypsum was applied to supply sulphur @ 30 kg ha<sup>-1</sup> in T<sub>8</sub>. Necessary uniform intercultural, water management and plant protection measures were undertaken in general until the crop was matured for harvesting. Before harvest of crop, grain yield was monitored through crop cutting.

### **2.3 Soil sample and plant analysis**

Individual soil samples from each plot were collected from surface downwards up to 75 cm depth at an interval of 15 cm i.e., 0-15, 15-30, 30-45, 45-60 and 60-75 cm after harvest of *kharif* rice crop in the year 2015 through profile digging. Immediately after collection, the soil samples were air dried, ground and passed through a 2 mm sieve and analysed for different basic soil properties by following standard laboratory procedures. Different forms of N viz., NO<sub>3</sub><sup>-</sup>-N, NH<sub>4</sub><sup>+</sup>-N, mineralizable N and total -N in soil samples were estimated adopting methods suggested by [7]. Plant samples after separating straw and grain were washed thoroughly, air dried and then oven dried at 60°C, finely pulverized and digested in a diacid mixture of HNO<sub>3</sub> and HClO<sub>4</sub> in 3:1 ratio. Nitrogen was determined by modified micro-Kjeldahl method [8]. Organic carbon (OC) in soil was determined by a wet oxidation procedure of [9].

### **2.4 Statistical Analysis**

The experiment has been laid out in Randomized Complete Block Design with 12 treatments and 4 replications. Analysis of variance has been conducted to test the overall significance of difference between the treatments.

Null hypothesis, H<sub>0</sub>: all the treatment means are identical

Alternate Hypothesis, H<sub>1</sub>: at least one pair of treatment mean differ significantly.

Test statistics for the treatment,  $F = \frac{TMS}{EMS}$

If the p value of F is <0.05, then F is considered to be significant and H<sub>0</sub> is rejected.

If F is found to be significant, then Least Significance Difference (LSD) test is conducted (at α=0.05) to test the significance of difference between each pair of treatment [10].

Simple linear regression analysis of was carried out between yield and sustainability with basic soil properties using the data analysis tool of MS Excel.

## **3. Results and Discussion**

### 3.1 Soil organic carbon

Application of various combination of inorganic fertilizer nutrients at varying levels or in combination with FYM had significant influence on soil organic carbon content after continuous rice-rice cropping system for 10 years (Table 1). In the surface soil it varied from 3.12 to 6.29 g kg<sup>-1</sup>. Non-application of fertilizer/FYM significantly decreased soil organic carbon content in control both in the surface and sub-surface soil layers, whereas, substantial build-up in the organic carbon content occurred under NPK+FYM, NPK+lime +FYM and NPK+lime treatments. The increase in SOC content under integrated use of fertilizers and organic manure treatments might have been due to direct incorporation of organic matter, better root growth and more plant residue addition resulting in increased soil organic carbon content. These results are in conformity with the finding of Singh *et al.* [11], Jadhao *et al.* [12], Ravankar *et al.* [13] and Mishra *et al.* [14] who reported the enhanced soil organic carbon status after ten years of continuous rice-wheat cropping under varying fertilizer and manure treatment in Mollisols at Pantnagar. The maximum increase in soil organic carbon content was observed with integrated use of inorganic fertilizers (N+P+K) and organic manure in rice-wheat cropping system in long-term experiment [15]. With depth the content decreased slightly up to second layer & sharply in the third layer (Fig 1)

### 3.2 Total Nitrogen

Total N content in soil as influenced by various treatments (Table 1) indicated that higher values of N content was obtained from surface soil which could be due to the presence of residues after the harvest of crop as suggested by Kushwaha *et al.* [16] and Singh *et al.* [17]. Further, lowest content found in unfertilized control (676.4, 543.4, 435.7, 221.6 and 110.2 kg ha<sup>-1</sup>, respectively under 0-15, 15-30, 30-45, 45-60 and 60-75 cm depths) is attributed to less productivity as a result of no fertilizer application because organic matter and biological fixation are the sources of N addition and both are dependent on primary productivity [17]. The total N content ranged from 676.4 to 1712.4 kg ha<sup>-1</sup> in surface soil and 110.2 to 1277.6 kg ha<sup>-1</sup> in sub-surface soil. Increase in total N was noted with concomitant increase in fertilizer levels from N alone (100% N) to optimal (100% NPK) and further to super optimal (150% NPK) level, which again due to increase in primary productivity [18]. The highest total N content was registered with 100% NPK+FYM (1712.4 kg ha<sup>-1</sup>) treatment followed by 100% NPK + lime + FYM (1691.3 kg ha<sup>-1</sup>) in the surface soil. The larger increase in total N in these plots was due to addition of extra N through FYM and greater amount of biomass to soil. Total soil N was also found to enhance in the treatment of INM (100% NPK + FYM), which is useful for N management under intensified rice -rice cropping in the subsequent years for better supply of N to the crops after its mineralization [19].

**Table 1. Soil organic carbon (g kg<sup>-1</sup>), total N (kg ha<sup>-1</sup>) and organic N (kg ha<sup>-1</sup>) in surface and sub surface soils under different treatments as influenced by long-term manurial practices**

Treatments	SOC (g kg <sup>-1</sup> )					Total N (kg ha <sup>-1</sup> )					Organic N (kg ha <sup>-1</sup> )				
	0-15 cm	15-30 cm	30-45 cm	45-60 cm	60-75 cm	0-15 cm	15-30 cm	30-45 cm	45-60 cm	60-75 cm	0-15 cm	15-30 cm	30-45 cm	45-60 cm	60-75 cm
100% PK	4.20	3.93	1.22	1.13	0.68	1019.6	812.5	523.5	240.8	122.7	952.4	786.8	502.3	227.1	114.7
100% NPK	4.67	4.27	1.28	1.11	0.80	1300.3	933.5	623.5	343.8	233.6	1218.2	904.8	601.2	329.9	225.4
150% NPK	5.05	4.35	1.32	1.18	0.98	1562.8	1235.8	986.7	565.8	207.7	1466.7	1203.3	964.0	552.0	199.1
100% NPK + Zn	4.92	4.52	1.49	1.19	0.85	1212.5	945.1	743.2	444.9	224.5	1124.2	912.8	721.4	431.1	216.1
100% NPK +FYM	6.29	5.77	2.12	2.08	1.50	1712.4	1277.6	945.8	589.6	366.3	1593.6	1221.9	912.4	570.4	354.5
100% NPK +lime +FYM	6.05	5.47	2.07	1.76	1.21	1691.3	1232.6	908.7	587.3	346.7	1579.8	1177.6	877.5	569.1	335.6
100% NPK +B +Zn	4.97	4.54	1.56	1.23	0.87	1377.5	965.3	713.6	388.5	292.6	1289.9	936.9	693.4	375.8	284.5
100% NPK S+ Zn	5.21	4.83	1.70	1.56	1.02	1408.6	1076.6	856.3	376.2	235.9	1320.5	1042.5	831.7	361.3	227.8
100%N	4.22	3.67	1.06	0.92	0.70	1011.2	850.6	712.5	322.5	145.8	936.6	826.7	692.9	311.2	138.5
100% NP	4.47	3.89	1.14	0.98	0.77	1057.1	876.8	744.2	334.6	154.5	980.3	849.6	724.7	322.8	147.1
100 % NPK +Lime	5.86	5.38	1.76	1.55	1.15	1415.7	1120.6	854.8	434.6	251.5	1331.1	1078.9	828.5	420.4	242.4
Control	3.12	2.69	0.92	0.86	0.62	676.4	543.4	435.7	221.6	110.2	623.2	524.6	421.1	212.0	104.0
CD <sub>(0.05)</sub>	0.57	0.48	0.20	0.17	0.11	49.3	39.4	32.3	27.8	21.3	49.5	39.1	32.6	28.1	21.2

Similar to soil organic C content, the total N content was higher in surface soil as compared to subsurface soil. Continuous cropping without fertilization and manuring also resulted in depletion of total nitrogen both in surface and sub-soil.

### 3.3 Organic N

Data on organic N presented in table 1 revealed that it varied from 623.2 kg ha<sup>-1</sup> to 1593.6 kg ha<sup>-1</sup> in surface soil and 104.0 kg ha<sup>-1</sup> (60-75 cm) to 1221.9 (15-30 cm) kg ha<sup>-1</sup> in sub-soil. Organic carbon contributed 92.1 to 97.7 % of total N. In control it constituted 92.1% on the surface layer, 96.5% in 15-30 cm, 96.6% in 30-45 cm, 95.7% in 45-60 cm and 94.4% at 60-75 cm layer. Highest organic N was recorded in high yielding NPK+FYM treatment where it constitutes 93.1, 95.6, 96.5, 96.7 and 96.8% at the respective surface and sub-soil layers. The content almost reduced to half in 30-45 cm layer and then there was a sharp fall with the depth.

### 3.4 Mineralizable N

Mineralizable N is easily oxidizable N from organic matter. The initial (2005) mineralizable N value of the soil (0-15 cm) was 187.0 kg ha<sup>-1</sup>. Improvement in mineralizable N values in both surface and sub-surface soil layers was noted in all the treatments (Table 2).

The mineralizable N content ranged from 154.4 to 291.3 kg ha<sup>-1</sup> in surface soil and from 8.2 (60-75 cm) to 119.3 (15-30 cm) kg ha<sup>-1</sup> in sub-surface soil. Increase in mineralizable N indicates that increase in easily oxidizable fraction of organic matter which means increase in availability of N to plant [16]. However, due to addition of fertilizer doses from 100% to 150%, N content was correspondingly improved indicating an impact of fertilizer application on enrichment of N pools [17]. The highest mineralizable N content registered in 100% NPK+FYM (291.3 kg ha<sup>-1</sup>) treatment followed by 100% NPK +lime +FYM (275.5 kg ha<sup>-1</sup>) could be resulted due to addition of larger amount N by paddy root and stubble biomass as well as addition of extra N through FYM each year and a part of it contributed to total N [20]. Cropping without any nutrient input led to perceptible decline in soil mineralizable N from the initial N content due to more removal of N through biomass and less root and stubble addition. In relation to the initial status, the mineralizable N declined by 21.1% in surface soils with continuous cropping in the control plot. Relatively low content of mineralizable N in sub-surface layer is due to less incorporation of biomass and poor biological activities [4].

### 3.5 Inorganic N

Inorganic N content presented in the table 2 indicated similar trend as total N. The inorganic N (NH<sub>4</sub><sup>+</sup>-N+NO<sub>3</sub><sup>-</sup>-N) content of soil was significantly higher at 0-15 cm depth as compared to that at 15-30 cm depth, which could be due to the presence of crop residues, FYM, more microbial activity and inorganic fertilizers in the surface layer [21]. The inorganic N content varied from 53.2 to 118.8 kg ha<sup>-1</sup> in surface soil and from 6.2 (60-75 cm) to 55.7 (15-30 cm) kg ha<sup>-1</sup> in sub-surface soil layer. The addition of different doses of fertilizers and FYM contributed towards relative increase in inorganic N content in surface and sub-surface layers. Similar, findings were reported by Khandagle *et al.* [22]. The highest inorganic N content was recorded in 100% NPK+FYM (118.8 kg ha<sup>-1</sup>) treatment followed by 100% NPK+lime+FYM (111.5 kg ha<sup>-1</sup>) and the lowest inorganic N content was obtained in control plot (53.2 kg ha<sup>-1</sup>) without fertilizer application, which directly or indirectly affected the plant growth. Increase in inorganic fraction of N is due to increase in organic carbon in soil which provided more energy source to microbes and also more site for the retention of NH<sub>4</sub><sup>+</sup>-N as well NO<sub>3</sub><sup>-</sup>-N which were replaceable with

KCl [16]. Further, lowest content was found in control (53.2 kg ha<sup>-1</sup>) which directly or indirectly affected normal biological activities.

**Table 2. Inorganic N (kg ha<sup>-1</sup>) and mineralizable N (kg ha<sup>-1</sup>) in surface and sub surface soils under different treatments as influenced by long-term manurial practices**

Treatments	Inorganic (NH <sub>4</sub> <sup>+</sup> -N+NO <sub>3</sub> <sup>-</sup> -N) N (kg ha <sup>-1</sup> )					Mineralizable N (kg ha <sup>-1</sup> )				
	0-15 cm	15-30 cm	30-45 cm	45-60 cm	60-75 cm	0-15 cm	15-30 cm	30-45 cm	45-60 cm	60-75 cm
100% PK	67.2	25.7	21.2	13.7	8	180.8	61.1	29.2	18.8	10.6
100% NPK	82.1	28.7	22.3	13.9	8.2	220.4	64.5	30.2	19.5	11.2
150% NPK	96.1	32.5	22.7	13.8	8.6	257.6	75.7	33.2	20.4	12.2
100% NPK + Zn	88.2	32.3	21.8	13.8	8.4	235.7	75.6	33.8	20.1	12.0
100% NPK +FYM	118.8	55.7	33.4	19.2	11.8	291.3	119.3	50.7	25.5	15.6
100% NPK +lime +FYM	111.5	54.9	31.2	18.2	11.1	275.5	116.3	40.1	22.9	14.6
100% NPK +B +Zn	87.6	28.4	20.2	12.7	8.1	238.3	70.1	30.2	18.6	11.5
100% NPK S+ Zn	88.1	34.1	24.6	14.9	8.1	232.8	76.9	35.2	21.3	12.7
100%N	74.5	23.9	19.6	11.3	7.3	205.1	55.1	28.2	17.9	10.9
100% NP	76.8	27.2	19.5	11.8	7.4	215.2	64.8	29.2	17.8	10.6
100 % NPK +Lime	84.6	41.7	26.3	14.2	9.1	217.9	92.7	35.6	21.3	12.9
Control	53.2	18.9	14.6	9.6	6.2	154.4	49.1	25.3	15.6	8.2
CD (0.05)	3.4	2.5	1.8	1.2	0.8	11.1	7.2	4.3	2.8	1.8

The content of NH<sub>4</sub><sup>+</sup> - N was recorded highest under NPK + FYM (80.9 kg ha<sup>-1</sup>) after 20<sup>th</sup> cropping cycle (Table 3). It ranged from 38.4 to 80.9 kg ha<sup>-1</sup> in the surface soil and its content decreased with depth up to 75 cm.

**Table 3. NH<sub>4</sub><sup>+</sup>-N (kg ha<sup>-1</sup>) and NO<sub>3</sub><sup>-</sup> - N (kg ha<sup>-1</sup>) in surface and sub surface soils under different treatments as influenced by long-term manurial practices**

Treatments	NH <sub>4</sub> <sup>+</sup> -N (kg ha <sup>-1</sup> )					NO <sub>3</sub> <sup>-</sup> -N (kg ha <sup>-1</sup> )				
	0-15 cm	15-30 cm	30-45 cm	45-60 cm	60-75 cm	0-15 cm	15-30 cm	30-45 cm	45-60 cm	60-75 cm
100% PK	47.2	17.2	12.9	8.4	4.9	20	8.5	8.3	5.3	3.1
100% NPK	57.6	19.4	15.3	9.8	5.9	24.5	9.3	7	4.1	2.3
150% NPK	65.9	23.4	16.8	9.9	6.3	30.2	9.1	5.9	3.9	2.3
100% NPK + Zn	56.3	20.7	13.0	9.1	5.9	31.9	11.6	8.8	4.7	2.5
100% NPK +FYM	80.9	37.9	23.9	13.1	7.8	37.9	17.8	9.5	6.1	4
100% NPK +lime +FYM	73.7	37.7	21.3	11.8	7.3	37.8	17.2	9.9	6.4	3.8
100% NPK +B +Zn	56.7	19.2	12.8	8.5	5.5	30.9	9.2	7.4	4.2	2.6
100% NPK S+ Zn	53.5	21.4	15.2	9.5	5.1	34.6	12.7	9.4	5.4	3
100%N	44.7	13.6	12.4	6.4	4.2	29.8	10.3	7.2	4.9	3.1
100% NP	51.6	16.9	13.1	8.2	5.2	25.2	10.3	6.4	3.6	2.2
100 % NPK +Lime	56.7	30.1	18.4	9.7	6.6	27.9	11.6	7.9	4.5	2.5
Control	38.4	12.2	10.8	7.4	4.8	14.8	6.7	3.8	2.2	1.4
CD (0.05)	2.2	1.8	1.6	1.2	1.1	1.8	1.0	0.75	0.70	0.62

Mineralization of FYM released appreciable amounts of ammonical N, which could contribute to exchangeable NH<sub>4</sub><sup>+</sup> -N [23]. As the organic carbon content is directly related with CEC, the N released through mineralization of organic matter and inorganic fertilizers applied could be retained in the exchange sites. Combined application of NPK + FYM also ensured higher NO<sub>3</sub><sup>-</sup> -N (37.9 kg ha<sup>-1</sup>) than control (14.8 kg ha<sup>-1</sup>) which could be due to increased microbial activity and resultant

enhanced nitrification process with a concomitant reduction in leaching losses (Fig 2). Increase in the rate of inorganic N added also enhanced the  $\text{NO}_3^-$ -N content possibly due to the conversion of applied mineral N via nitrification process. The  $\text{NO}_3^-$ -N content varied from 1.4 to 17.8 kg ha<sup>-1</sup> in the sub-soil and the lowest was recorded at 60-75 cm layer. The drop in  $\text{NO}_3^-$ -N was more sharp up to 45 cm than the deeper layers beyond 45 cm where it was almost constant.

### 3.6 Correlation among various pools of N and SOC of different layers

Soil organic carbon of all the layers strongly correlated with total N, mineralizable N, inorganic N, ammonium N and nitrate N of respective layers (Table 4). Among various N pools there was strong correlation of organic N with total N, mineralizable N, inorganic N and ammonium N of respective layers. Total N also had strong positive correlation with inorganic N and ammonium N of all the layers. Correlation of organic N and total N with nitrate N was however significant up to 30 cm depth and beyond that there was no correlation at 30-45 cm layer ( $r= 0.511$  &  $0.524$ ).

**Table 4. Correlation coefficients (r) between different nitrogen fractions and SOC at different depths**

<b>Total N</b>					
	<i>0-15 cm</i>	<i>15-30 cm</i>	<i>30-45 cm</i>	<i>45-60 cm</i>	<i>60-75 cm</i>
SOC	0.941**	0.902**	0.716**	0.748**	0.880**
Organic N	0.999**	0.999**	0.999**	0.999**	0.999**
Inorganic N	0.951**	0.845**	0.748**	0.793**	0.883**
<b>Mineralizable N</b>					
SOC	0.883**	0.919**	0.899**	0.956**	0.946**
Organic N	0.935**	0.832**	0.714**	0.829**	0.888**
<b>Inorganic N</b>					
SOC	0.920**	0.914**	0.934**	0.937**	0.926**
Organic N	0.945**	0.829**	0.734**	0.786**	0.878**
Total N	0.951**	0.845**	0.748**	0.793**	0.883**
Mineralizable N	0.988**	0.997**	0.944**	0.964**	0.954**
<b>NH<sub>4</sub><sup>+</sup> - N</b>					
SOC	0.844**	0.913**	0.869**	0.902**	0.895**
Organic N	0.926**	0.839**	0.737**	0.812**	0.849**
Total N	0.931**	0.855**	0.750**	0.813**	0.852**
Mineralizable N	0.949**	0.992**	0.930**	0.924**	0.842**
<b>NO<sub>3</sub><sup>-</sup> - N</b>					
SOC	0.854**	0.843**	0.829**	0.761**	0.680*
Organic N	0.847**	0.734**	0.511	0.548	0.646*
Total N	0.853**	0.751**	0.524	0.557*	0.650*
Mineralizable N	0.917**	0.927**	0.699**	0.801**	0.819**

### 3.7 Correlation of contents of various nitrogen pools of bottom layers with those of surface layer

It was also observed that the correlation of lower layers with surface layer in different fractions of N was found positively significant. The r value in the table 5 indicated that the correlation between SOC of surface layer with lower layers was highly significant ( $r=0.997^{**}$ ,  $0.958^{**}$ ,  $0.933^{**}$  and  $0.949^{**}$ ). Depending on type of

soil, cropping system and agronomic management practices and type of nutrient, the relationship of contents of N of surface layer with that of bottom layers may differ. Under wet land rice-rice production system, correlation of lower layer contents with upper layer has been worked out. All the form of N, such as total N, inorganic N, mineralizable N in lower had strong relationship on that of surface layer.

**Table 5. Correlation of lower layer contents of SOC and various fractions of N with that of surface layer**

N-fractions	Correlation coefficient (r)			
	15-30 cm	30-45 cm	45-60 cm	60-75 cm
SOC	0.997**	0.958**	0.933**	0.949**
Total N	0.985**	0.956**	0.975**	0.941**
Organic N	0.985**	0.952**	0.973**	0.941**
Inorganic N	0.905**	0.880**	0.906**	0.927**
Ammonium N	0.910**	0.869**	0.939**	0.930**
Nitrate N	0.844**	0.778**	0.769**	0.754**
Mineralizable N	0.842**	0.837**	0.864**	0.914**

Like SOC, the surface layer content of N of various forms like total N ( $r=0.985^{**}$ ,  $0.956^{**}$ ,  $0.975^{**}$  and  $0.941^{**}$ ), organic N ( $r= 0.985^{**}$ ,  $0.952^{**}$ ,  $0.973^{**}$  and  $0.941^{**}$ ), inorganic N ( $r= 0.905^{**}$ ,  $0.880^{**}$ ,  $0.906^{**}$  and  $0.927^{**}$ ), ammonium N ( $r=0.910^{**}$ ,  $0.869^{**}$ ,  $0.939^{**}$  and  $0.930^{**}$ ) and Nitrate N ( $r=0.844^{**}$ ,  $0.778^{**}$ ,  $0.769^{**}$  and  $0.754^{**}$ ) were positively correlated with those of bottom layers. The results of correlation studies of SOC, organic N and inorganic N content of different layers (0-15 cm, 15-30 cm, 30-45 cm, 45-60 cm and 60-75 cm) were found to strongly correlated with the total N content of the respective layers.

### 3.8 Grain yield and nitrogen uptake

The fertilizer treatments comprising fertilizer alone or combination with FYM/soil amendment had significant effect on grain yield of rice (Table 6) in *kharif* season. Perusal of results in grain yield revealed that there was a significant variation among the treatments. The highest grain yield was recorded in the treatment, 100%NPK+FYM which was at par with 100%NPK+FYM+lime. This may be attributed to better utilization of applied nutrients through the activities of greater population of soil micro-organisms which caused more nutrient transformation and also release of nutrients from organic sources that influenced more nutrient availability to the crop plants as well as the potential for higher production. Moreover, organic manures also supply growth promoting substances like enzymes and hormones [24]. Similar results were reported by Kandeshwari *et al.* [25]. Hence, integrated use of organic and inorganic fertilizers can make important contribution for increasing and sustaining rice production. This was also evidenced by studies of Jeyajothi and Nalliah Durai Raj [26] and Nayak *et al* [27].

**Table 6. Effect of long-term manurial practices on grain yield, N uptake and yield sustainability of rice after 10 years of continuous cropping with rice-rice system**

Treatments	Khariif season Grain yield ( $q\ ha^{-1}$ )	Total N uptake ( $kg\ ha^{-1}$ )	Sustainable Yield Index (SYI) of the system
T-1	100% PK	32.80	0.43
T-2	100% NPK	41.80	0.45
T-3	150% NPK	46.60	0.47
T-4	100% NPK + Zn	45.05	0.45

T-5	100% NPK +FYM	51.58	97.6	0.54
T-6	100% NPK +lime +FYM	49.15	93.1	0.53
T-7	100% NPK +B +Zn	45.45	74.5	0.46
T-8	100% NPK S+ Zn	45.38	75.2	0.44
T-9	100%N	35.50	48.1	0.38
T-10	100% NP	40.90	68.3	0.41
T-11	100 % NPK +Lime	43.60	75.8	0.45
T-12	Control	21.95	25.2	0.20
	CD <sub>(0.05)</sub>	4.83	-	-

Super optimal dose of NPK i.e., 150% NPK produced significantly higher yield (46.6 q ha<sup>-1</sup>) than recommended dose i.e., 100% NPK (41.80 q ha<sup>-1</sup>). The lowest grain yield (21.95 q ha<sup>-1</sup>) was recorded in control. The results of twenty cycles of long-term fertilizer experiment indicated that application of 100% N alone had increased the yield by 61.7 per cent over control. The supplementation of P with N (100% NP) enhanced the yields by 15.2 per cent over 100% N treatment. Application of Zn, Zn +S and Zn + B did not have any significant effect on grain yield.

Total N uptake (Table 6) by rice crop varied from 25.2 to 97.6 kg ha<sup>-1</sup>. The highest N uptake was found in 100% NPK+FYM, followed by 100% NPK+lime+FYM and the lowest N uptake (25.2 kg ha<sup>-1</sup>) was observed in control plot. Similar findings were also reported by Dwivedi *et al.* [20] who found maximum N uptake with the treatment of integrated use of organic manure and recommended fertilizer dose as compared to control plot.

### 3.9 Correlation of various pools of N of different layers with N uptake, grain yield and sustainability

The correlation coefficient values of N fractions with grain yield, N uptake and soil organic carbon content of surface soil are presented in table 7.

**Table 7. Correlation of various pools of N of different layers with N uptake, grain yield and sustainability**

<b>Total N</b>					
	<i>0-15 cm</i>	<i>15-30 cm</i>	<i>30-45 cm</i>	<i>45-60 cm</i>	<i>60-75 cm</i>
Grain Yield	0.934**	0.923**	0.875**	0.859**	0.848**
N uptake	0.964**	0.966**	0.915**	0.925**	0.836**
Sustainability	0.902**	0.888**	0.756**	0.776**	0.786**
<b>Mineralizable N</b>					
Grain Yield	0.947**	0.773**	0.750**	0.827**	0.885**
N uptake	0.958**	0.824**	0.783**	0.850**	0.892**
Sustainability	0.876**	0.765**	0.733**	0.837**	0.881**
<b>Inorganic N</b>					
Grain Yield	0.924**	0.763**	0.788**	0.784**	0.780**
N uptake	0.943**	0.811**	0.819**	0.814**	0.824**
Sustainability	0.881**	0.765**	0.830**	0.847**	0.831**
<b>Organic N</b>					
Grain Yield	0.931**	0.923**	0.870**	0.857**	0.848**
N uptake	0.963**	0.966**	0.911**	0.924**	0.835**
Sustainability	0.900*	0.886**	0.748**	0.771**	0.784**

\*\*Significant at 1% level of significance; \*Significant at 5% level of significance

Correlation of total N, inorganic N and organic N with N uptake, grain yield and sustainability revealed that there exists a strong positive correlation of all the forms of N of all the layers up to 75 cm with N uptake, grain yield and sustainability.

This indicates that lower layer N forms have a strong bearing on the N uptake, grain yield and yield sustainability under low land rice-rice production system, although the treatments differ with respect to contents. This is justified by the facts that there exists a strong positive correlation of sub-soil contents various pools of N with those of top soil respectively.

#### 4. Conclusion

Not only the surface soil but also the sub-soil contents of total N, organic N and mineralizable N are significantly influenced by long-term use of fertilizer nutrients and manure under wet land intensive rice production system. Sub-soil contents of total N, organic N, mineralizable N, inorganic N, ammonium N and nitrate N have strong positive correlation with those of surface layer. SOC, total N mineralizable N and inorganic N are strongly correlated with each other in the same layer. Correlation of organic N and total N with nitrate N was however significant up to 30 cm depth and beyond that there was no correlation at 30-45 cm layer ( $r = 0.511$  &  $0.524$ ).

From the results, it is concluded that application of nutrient in balanced and integrated manner (NPK + FYM) not only increased grain yield of rice but also improved various fractions of N in surface as well as sub-soil.

#### References

1. Sahrawat KL. Macro and micronutrients removed by upland and lowland rice cultivars in West Africa. *Communications in Soil Science and Plant Analysis*. 2000, 31: 717-723.
2. Hofman G, Cleemput OV. Soil and Plant Nitrogen, International Fertilizer Industry Association Paris. 2004, 1-31
3. Eagle AJ, Bird JA, Hill JE, Horwath WR, Van Kessel, C. Nitrogen dynamics and fertilizer N use efficiency in rice following straw incorporation and winter flooding. *Agronomy Journal*. 2001, 93: 1346-1354.
4. Sawarkar SD, Singh M, Khamparia NK, Wanjari RH, Thakur R. Long-term impact of fertilizer and manure on soil quality, crop productivity and sustainability under soybean-wheat system in Vertisol. *Research Bulletin of All India Coordinated Research Project on Long-Term Fertilizer Experiment*, Department of Soil Science and Agricultural Chemistry, Jawaharlal Nehru Krishi Vishwa Vidyalyaya, Jabalpur (M.P.). 2015, 1-63.
5. Li K, Wang C, Li X, Li H, Dong M, Jin S, Liu L, Zhu C, Xue R. Long-term effect of integrated fertilization on maize yield and soil fertility in a calcareous fluvisol. *Arch. Agron. Soil Sci.* 2020, 67: 1400-1410.
6. Winters E, Simonson RW. The subsoil, *Advances in Agronomy*. III. 1951
7. Bremner JM. Inorganic forms of nitrogen. In *Methods of Soil Analysis*. Part 2 (C.A. Black, Ed.) Monograph No. 9. American Society of Agronomy. Madison Wisconsin, USA. 1965 :1179-1237.
8. Jackson ML. *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd., New Delhi. 1973
9. Walkley A, Black IA. An examination of the Degtjareff method for determination of soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Science*. 1934, 37: 29-38.
10. Gomez KA, Gomez A A. *Statistical Procedures for Agricultural Research*, 2nd ed. Wiley - Interscience, New York. 1984
11. Singh NP, Sachan RS, Pandey PC, Bisht PS. Effect of a decade long-term fertilizer and manure application on fertility and productivity of rice-wheat system in a *Mollisols*. *Journal of Indian Society of Soil Science*. 1999, 47(1): 72-80.

12. Jadhao SD, Mali DV, Kharche VK, Singh M, Bhoyar SM, Kadu PR, Wanjari RH, Sonune BA. Impact of continuous manuring and fertilization on changes in soil quality under sorghum-wheat sequence on a *Vertisols*. *Journal of the Indian society of Soil science*. 2019, 67 (1): 55-64.
13. Ravankar HN, Gajbhiye NN, Sarap PA. Effect of organic manures and inorganic fertilizers on yield and availability of nutrients under sorghum- wheat sequence. *Indian Journal of Agriculture Research*. 2005, 39: 142-145.
14. Mishra B, Sharma A, Singh, SK, Prasad J, Singh BP. Influence of continuous application of amendments to maize-wheat cropping system on dynamics of soil microbial biomass in Alfisol of Jharkhand. *Journal of the Indian Society of Soil Science*. 2008, 56: 71-75.
15. Beri V, Sidhu BS, Bahl, GS, Bhat AK. Nitrogen and phosphorus transformations as affected by crop residue management practices and their influence on crop yield. *Soil Use and Management*. 1995, 11(2), 51-54.
16. Kushwaha S, Sawarkar SD, Thakur R, Khamparia NK, Singh M. Impact of Long-Term Nutrient Management on Soil N dynamics under Soybean – Wheat Cropping Sequence on a *Vertisol*. *Journal of the Indian Society of Soil Science*. 2017, 65 (3): 274-282.
17. Singh M, Wanjari RH, Dwivedi A, Dalal R. Yield response to applied nutrients and estimates of N<sub>2</sub> fixation in 33-year-old soybean–wheat experiment on a *Vertisol*. *Experimental Agriculture*. 2012, 48: 311-325.
18. Thakur R, Sawarkar SD, Vaishya UK, Singh M. Impact of continuous use of inorganic fertilizers and organic manure on soil properties and productivity under soybean-wheat intensive cropping of a *Vertisol*. *Journal of the Indian Society of Soil Science*. 2011, 59: 74-81.
19. Katkar RN, Sonune BA, Kadu PR. Longterm effect of fertilization on soil chemical and biological characteristics and productivity under sorghum (*Sorghum bicolor*)–wheat (*Triticum aestivum*) system in *Vertisol*. *Indian Journal of Agricultural Sciences*. 2011, 81:734-739.
20. Dwivedi AK, Singh M, Kauraw DL, Wanjari RH, Chauhan S. Impact of fertilizer and manure use for three decades on crop productivity and sustainability, and soil quality under soybean-wheat system on a *Vertisol* in Central India. *Research Bulletin of All India Coordinated Research Project on Long-Term Fertilizer Experiment*, Department of Soil Science and Agricultural Chemistry, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur (M.P.). 2007.
21. Soudi B, Sbai A, Chiang CN. Nitrogen mineralization in semiarid soils of Morocco: rate constant variation with depth. *Soil Science Society of America Journal*. 1990, 54, 756-761.
22. Khandagle A, Dwivedi BS, Dwivedi AK, Panwar S, Thakur RK. Nitrogen Fractions under Long-Term Fertilizer and Manure Applications in Soybean-Wheat Rotation in a *Vertisol*, *Journal of the Indian Society of Soil Science*. 2020, 68 (2): 186-193.
23. Dhakshinamoorthy M, Singh M, Malarvizhi P, Selvi D, Bhaskaran A. Soil quality, crop productivity and sustainability as influenced by long- term fertilizer application and continuous cropping of finger millet-maize-cowpea sequence in swell-shrink soil., *Research Bulletin No.3*, AICRP on LTFE to study changes in soil quality, crop productivity and sustainability, IISS, Bhopal. 2005, 66-70.
24. Yadav BK, Lourdura AC. Effect of Organic Manures and Panchagavya Spray on Yield attributes, Yield and Economics of Rice (*Oryza sativa* L.). *Crop Research*. 2006. 31(1): 1-5.

25. Kandeshwari MS, Jeyaraman and Thavaprakash N. Evaluation of integrated nutrient management practices under system of rice intensification. *In. International Symposium on 100 years of Rice Science and Looking Beyond.* 2012, 2: 523-524.
26. Jeyajothi R, Nalliah Durai Raj S. Influence of Integrated Nutrient Management Practices on Yield and Yield Attributes, and Economics of Transplanted Rice (*Oryza sativa* L.) in South Zone of Tamil Nadu. *International Journal for Innovative Research in Science & Technology.* 2015, 1(11): 462-467.
27. Nayak AK, Gangwa B, Shukla AK, Mazumdar SP, Kumar A, Raja R, Kumar A, Kumar V, Rai PK, Mohan U. Long-term effect of different integrated nutrient management on soil organic carbon and its fractions and sustainability of rice-wheat system in Indo Gangetic Plains of India. *Field Crops Research.* 2012 127: 129-139.