

## Original Research Article

### **Efficacy of selected insecticides and botanicals against the infestation of gram pod borer, *Helicoverpa armigera* Hub. on chickpea, *Cicer arietinum* L.**

#### **Abstract**

The present research was conducted at the Central Research Field of Department of Agricultural Entomology, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj during the *Rabi* season of 2021-22. The treatments used during the study were *Metarhizium anisopliae*, *Beauveria bassiana*, Karanj Oil, Neem Oil, Spinosad, Emamectin benzoate + Indoxacarb and Indoxacarb. The results revealed that highest percent reduction in larval population was observed in Emamectin benzoate 5% SG + Indoxacarb 14.5% SC (73.57%) and was most successful in bringing down the larval population. It is then followed by Spinosad 45% SC (65.03%), Indoxacarb 14.5% SC (59.02%), *Beauveria bassiana* ( $1 \times 10^8$  cfu) (54.87%), *Metarhizium anisopliae* ( $1 \times 10^8$  cfu) (50.92%), Neem Oil 2% (45.81%). While, Karanj Oil 2% prove to be the least effective treatment in reducing the larval population (37.53%).

**Key Words:** Benefit Cost Ratio, Biopesticides, Insecticides, Pod borer, Treatments and Yield

#### **Introduction**

Chickpea, *Cicer arietinum* (L.) belongs to the family Leguminosae (Fabaceae) (Sanjayrao, 2013) which is basically a diploid plant with  $2n=2x=16$  no. of chromosomes. Chickpea has its origin from South-eastern part of Turkey and which later spreaded to different regions of the world. Chickpea or Bengal gram, commonly known as 'Chana' is also considered to be the third most important pulse crop around the world.

However, India is also considered to be the highest producer of chickpea, thus attaining the title of 'king of pulses' (Sai *et al.*, 2021). Some of the most important chickpea producing countries are India, Turkey, Iran, Ethiopia, Myanmar, Pakistan, Australia, Canada, Mexico etc. It has been estimated that the cultivated area under the chickpea is around 10.91 million ha with an annual production and productivity of

approximately 9.78 million tones and 929 kg/ha respectively (**Chandrasekhara et al., 2017**). Chickpea can be used both as nutritional leguminous food for human and feed for the cattle. Moreover, the chickpea consumption can be linked to many health benefits such as affecting the markers of both metabolic syndrome and cardiovascular diseases in human as wells as animals (**Wallace et al., 2016**).

Among the major pests attacking chickpea viz. *Helicoverpa armigera*, *Spodoptera litura*, *Plusia orichalchea*, *Agrotis ipsilon*, *Aphis cracivorea*, *Bemisia tabaci* etc., *Helicoverpa armigera* (Gram pod borer) causes more damage and yield losses to the crop (**Singh et al., 2006**). The larval stage of gram pod borer is the most damaging one. The larva feeds on the flowers, pods, developing seeds and leaves of the chickpea plant. Gram pod borer causes an yield loss of about 200 million US dollar to chickpea and pigeon pea in India (**Rummana et al., 2010**).

## Materials and Methods

The present experiment was carried out in Randomised Block Design replicated thrice at the Central Research Field of department of Entomology of SHUATS, Prayagraj during *rabi* 2021-22. The chickpea variety

used in the experiment was 'Pusa 362'. The larval population count of *Helicoverpa armigera* on chickpea was accomplished by observing 5 randomly selected plants per plot at weekly intervals in each treatment. The mean larval population was worked out and the transform value was taken to analyze the data as per standard statistical methods. In order to know the most effective treatment in reducing the larval population percent larval reduction was worked out.

$$\text{Percent Larval Reduction} = \frac{C - T}{C} \times 100$$

Where,

C= Controlled Plot

T= Treated Plot

The desired concentration of insecticidal spray solution for each treatment was freshly prepared each and every time at the site of experiment, just before the start of spraying operations.

The spray solution of desired concentration was prepared by adopting the following formula:

$$V = \frac{(C \times A)}{\% a.i.}$$

Where,

V= Volume of a formulated pesticide required

C= Concentration required

A= Volume of the total solution to be prepared  
% a.i. = Given Percentage strength of a formulated pesticide.

## Result and Discussion

The results obtained on the larval population of *Helicoverpa armigera* on third day after 1<sup>st</sup> spray revealed that Emamectin benzoate + Indoxacarb (220gm/ha + 0.2 ml/l) proved to be most effective in reducing the larval population by 71.59%, followed by Spinosad (0.5ml/l) 58.83%, Indoxacarb (1ml/l) 50.61%, *Beauveria bassiana* (2gm/l) 45.84%, *Metarhizium anisopliae* (5gm/l) 38.7%, Neem oil (2%) 37.57% and Karanj oil (2%) 29.32%. On seventh day after 1<sup>st</sup> spray, Emamectin benzoate + Indoxacarb showed the maximum larval reduction by 77.05%, followed by Spinosad (69.44%), Indoxacarb (65.10%), *Beauveria bassiana* (57.58%), *Metarhizium anisopliae* (54.32%), Neem oil (47.62%) and Karanj oil (38.90%). On fourteenth day after 1<sup>st</sup> spray, Emamectin benzoate + Indoxacarb showed the maximum larval reduction by 62.77%, followed by Spinosad (52.71%), Indoxacarb (49.5%), *Beauveria bassiana* (48.8%), *Metarhizium anisopliae* (42.52%), Neem oil (37.17%) and Karanj oil (31.81%).

The results obtained from the third day after 2<sup>nd</sup> spray revealed that Emamectin benzoate + Indoxacarb (220gm/ha + 0.2 ml/l) proved to be most effective in reducing the larval population by 72.83%, followed by followed by Spinosad (66.60%) (0.5ml/l), Indoxacarb (58.62%) (1ml/l), *Beauveria bassiana* (57.49%) (2gm/l), *Metarhizium anisopliae* (53.51%) (5gm/l), Neem oil (48.52%) 2% and Karanj oil (39.33%) 2%. Seventh day after 2<sup>nd</sup> spray revealed that Emamectin benzoate + Indoxacarb proved to be most effective in reducing the larval population by 83.79%, followed by followed by Spinosad (74.67%), Indoxacarb (69.7%), *Beauveria bassiana* (64.57%), *Metarhizium anisopliae* (63.38%), Neem oil (55.46%) and Karanj oil (45.38%). Fourteenth day after 2<sup>nd</sup> spray revealed that Emamectin benzoate + Indoxacarb proved to be most effective in reducing the larval population by 73.49%, followed by followed by Spinosad (67.57%), Indoxacarb (60.6%), *Beauveria bassiana* (54.79%), *Metarhizium anisopliae* (52.80%), Neem oil (46.91%) and Karanj oil (40.17%).

In case of the mean of 1<sup>st</sup> spray, the results revealed that Emamectin benzoate + Indoxacarb was most effective in reducing the larval population by 70.54%, followed by followed by Spinosad (60.42%),

Indoxacarb (55.07%), *Beauveria bassiana* (50.75%), *Metarhizium anisopliae* (45.17%), Neem oil (41.29%) and Karanj oil (33.4%).

In case of the mean of 2<sup>nd</sup> spray, the results revealed that Emamectin benzoate + Indoxacarb was most effective in reducing the larval population by 76.71%, followed by followed by Spinosad (69.64%), Indoxacarb (62.97%), *Beauveria bassiana* (59.28%), *Metarhizium anisopliae* (56.57%), Neem oil (50.29%) and Karanj oil (41.63%).

The results obtained from the overall mean of the 1<sup>st</sup> and 2<sup>nd</sup> sprays revealed that, Emamectin benzoate + Indoxacarb was most effective in reducing the larval population by 73.62%, followed by followed by Spinosad (65.02%), Indoxacarb (56.23%), *Beauveria bassiana* (55.01%), *Metarhizium anisopliae* (50.87%), Neem oil (45.80%) and Karanj oil (37.53%).

Thus, from table 1 it is clear that all the treatments used in the experiment were found to be significantly superior over control. These results are similar to the findings of **Sai et al., (2021)** where the highest yield was recorded in Emamectin benzoate (5% SG) + Indoxacarb (14.5%). These results were supported by **Chandrasekhar et al., (2017)** where highest

reduction in larval population of gram pod borer (72.12%) was observed with spinosad 45 SC @ 0.5 ml/l. The results reported by **Kumar et al., (2014)** proved that Indoxacarb 14.5% SC was the best insecticide followed by spinosad 45 SC. Similar findings were also recorded by **Goutham and Tayde (2017)** which showed lowest percent infestation of gram pod borer was recorded in chloropyriphos (2.39%), followed by NSKE (2.94%), Neem oil (3.36%) Neem leaf extract (3.58%) and Garlic extract (3.90%), Pongamia oil (4.45%) and Papaya leaf extract (4.61%). **Choudhary et al., (2017), Harshita et al., (2018), Regmi et al., (2018)** and **Singh et al., (2017)** also supports the results.

**Table 1:** Effect of selected insecticides and biopesticides against *Helicoverpa armigera* (Hub.) on chickpea during *Rabi* season of 2021-22

		Percent reduction of larval population of <i>H. armigera</i>										
		1 <sup>st</sup> spray				2 <sup>nd</sup> spray						
Treatments		3DAS*	7DAS	14DAS	Mean	3DAS	7DAS	14DAS	Mean	1 <sup>st</sup> Spray Mean	2 <sup>nd</sup> Spray Mean	Overall Mean
T <sub>1</sub>	<i>Metarhizium anisopliae</i> (1x10 <sup>8</sup> cfu)	38.81	54.32	42.52	45.17	53.51	63.38	52.80	56.57	45.17	56.57	50.87
T <sub>2</sub>	<i>Beauveria bassiana</i> (1x10 <sup>8</sup> cfu)	45.84	57.58	48.80	50.75	57.49	64.57	54.79	59.28	50.75	59.28	55.01
T <sub>3</sub>	Karanj Oil 2%	29.32	38.90	31.81	33.40	39.33	45.38	40.17	41.63	33.4	41.63	37.53
T <sub>4</sub>	Neem Oil 2%	37.57	47.62	37.17	41.29	48.52	55.46	46.91	50.29	41.29	50.3	45.80
T <sub>5</sub>	Spinosad 45% SC	58.83	69.44	52.71	60.42	66.60	74.67	67.57	69.64	60.4	69.64	65.02
T <sub>6</sub>	Emamectin benzoate 5% SG + Indoxcarb 14.5% SC	71.59	77.05	62.77	70.54	72.83	83.79	73.49	76.71	70.54	76.71	73.62
T <sub>7</sub>	Indoxcarb 14.5% SC	50.61	65.10	49.5	55.07	58.62	69.7	60.6	62.97	49.5	62.97	56.23
T <sub>0</sub>	Control	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
F-test		*S	S	S	S	S	S	S	S	S	S	S
CV		5.71	5.42	7.66	7.89	6.09	4.96	6.68	4.16	7.89	4.16	5.87
Cd( 0.05 )		4.16	4.87	5.45	6.15	5.34	4.97	5.80	3.81	6.15	3.81	6.65

## Conclusion

From the analysis of the present findings, it can be concluded that among all the treatments used in the experiment, Emamectin benzoate (220 gm/ha) + Indoxacarb (0.2 ml/l) has proved to be the most effective in reducing the larval population of *Helicoverpa armigera*, being followed by Spinosad (0.5ml/l), Indoxacarb (1ml/l), *Beauveria bassiana* (2gm/l), *Metarhizium anisopliae* (5gm/l), Neem oil (2%) and Karanj oil (2%). Thus, Karanj Oil 2% prove to be the least effective treatment in reducing the larval population.

## References

- Adsure, P. S. and Mohite, B. P. (2015).** Efficacy Of Entomopathogenic Fungi Against Gram Pod Borer, *Helicoverpa Armigera* (Hub.) On Chickpea. *Journal of Global Biosciences*, **8** (4):3154-3157.
- Bhushan, S., Singh, R. P., and Shanker, R. (2011).** Bioefficacy of neem and Bt against pod borer, *Helicoverpa armigera* in chickpea. *Journal of Biopesticides*, **4**(1): 87 – 89.
- Chandrasekhar, G.V. (2017).** Efficacy of Certain Chemicals and Neem Products against *Helicoverpa armigera* (Hubner) on chickpea (*Cicer arietinum* L.). *Journal of Biopesticides*, **8** (4):3154-3157.
- Chaukikar, K., Bhowmick, A. K., Das, S. B., Marabi, R. S., and Tomar, V. S. (2017).** Bioefficacy of emamectin benzoate against *Helicoverpa armigera* Hubner and its natural enemies on chickpea (*Cicer arietinum*) crop. *International Journal of Bio-resource and Stress Management*, **8**(5), 716-720.
- Dagar, V. S., and Kumar, S. (2018).** Emamectin benzoate: Potential larvicide and antifeedant agent against cotton Boll worm *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Journal of applied and natural science*, **10**(2), 564-571.
- Goutham, K. (2018).** Comparative efficacy of selected botanicals and chloropyriphos against gram pod borer [*Helicoverpa armigera* (Hubner)] on chickpea [*Cicer arietinum* (L.)]. *Journal of Pharmacognosy and Phytochemistry*, **7**(1), 296-298.
- Jadhav, K. U., Chavan, A. P., More, S. A., Kulkarni, S. R., and Karande, R. A. (2021).** To study the efficacy of molecule combinations against gram pod borer (*Helicoverpa armigera* , Hubner) in chickpea (*Cicer arietinum* L.). *Journal of Entomology and Zoology Studies* 2021; **9**(6): 164-170
- Japur, K., Naveena, R., and Hosamani, A. C. (2016).** Bio-efficacy of emamectin benzoate 5-wg against

- pigeon pea pod borer, *Helicoverpa armigera* (Hub.) Under field condition in steppe climate of HK region. *J. Env. Bio-Sci*, **30**(1), 165-167.
- Kambrekar, D. N., Kulkarni, K. A., Giraddi, R. S., Kulkarni, J. H., and Fakrudin, B. (2009).** Management of chickpea pod borer, *Helicoverpa armigera* (Hubner) through Nuclear Polyhedral Virus isolates. *Precision agriculture*, **10**(5), 450-457.
- Kumar, P., Kumar, S., Rana, R., and Sachan, S. K. (2014).** Comparative efficacy of novel insecticides and bio-pesticides on larval population density of gram pod borer (*Helicoverpa armigera* Hubner) on chickpea. *Journal of Plant Development Sciences* Vol, **6**(2), 335-338.
- Kumar, A., Tripathi, M. K., Chandra, U., and Veer, R. (2019).** Efficacy of botanicals and bio-pesticide against *Helicoverpa armigera* in chickpea. *Journal of Entomology and Zoology Studies*, **7**(1), 54-57.
- Lakshmikanth, R., and Kumar, A. (2018).** Comparative efficacy of selected chemicals and Biopesticides against gram pod borer [*Helicoverpa armigera* (Hubner)](Lepidoptera: Noctuidae) on cowpea [*Vigna unguiculata* (L.) Walp.]. *Journal of Pharmacognosy and Phytochemistry*, **7**(3), 3307-3309.
- Lavanya, V., and Kumar, A. (2022).** Efficacy of certain chemicals against gram pod borer [*Helicoverpa armigera* (Hubner)] on chickpea (*Cicer arietinum* L.). *The Pharma Innovation Journal*; **11**(3): 1293-1297
- Maurya, O., and Kumar, H. (2018).** Growth of chickpea production in India. *Journal of Pharmacognosy and Phytochemistry*, **7**, 1175-1177.
- Mahmood, M. T., Akhtar, M., Ahmad, M., Saleem, M., Aziz, A., Rasool, I., & Amin, M. (2021).** An update on biology, extent of damage and effective management strategies of chickpea pod borer (*Helicoverpa armigera*). *Pakistan Journal of Agricultural Research*, **34**(1), 91.
- Narayan, S. R. P., Singh, P. S., and Singh, S. K. (2015).** Evaluation of certain new insecticides against *Helicoverpa armigera* (Hubner) on chickpea (*Cicer arietinum* L.). *Journal of Experimental Zoology*, **18**(1), 227-231.
- Nithish, A., Joshi, B., Jayaram, C. S., and Kariyanna, B. (2015).** Bio-efficacy of Some Newer Insecticides against Gram Pod Borer (*Helicoverpa armigera*) in Pigeonpea under Field Conditions. *Research Journal of Agricultural Sciences*, **6**(3), 563-566.

- Nishad, R. N., Singh, R. B., Singh, A. K., Singh, S. P., and Yadav, S. K. (2017).** Effect of various indigenous botanical seed protectants on seed quality parameters of chickpea seed during ambient storage. *Journal of Pharmacognosy and Phytochemistry*, 423-426.
- Olle, M., Tsahkna, A., Tähtjärv, T., and Williams, I. H. (2015).** Plant protection for organically grown potatoes—a review. *Biological Agriculture & Horticulture*, **31**(3), 147-157.
- Prasad, J. (2008).** Efficacy of *Metarhizium anisopliae* and *Beauveria bassiana* against *Helicoverpa armigera* in Chickpea, under Field Conditions in Nepal. *Journal of Plant Development Sciences* Vol, **6**(2), 335-538
- Patil, S. B., Goyal, A., Chitgupekar, S. S., Kumar, S., and El-Bouhssini, M. (2017).** Sustainable management of chickpea pod borer. A review. *Agronomy for Sustainable Development*, **37**(3). *Journal of Scientific and Industrial Research*, **43**(2), 321-224
- Rummana A, Choudhury A, Parvez Q, Sayed HP and Galib (2010).** Antifeedent Response of Two Medicinal Plants against *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) on chickpea, *Cicer arietinum* Middle-East Journal of Scientific Research.; **5**(5):329-335.
- Reza, M.E., Ali, M.R., Islam, M.S., Islam, M.J. and Roy, H.P. (2016)** Ecofriendly management of chickpea pod borer. *Eco-friendly Agricultural Journal*. **9**(06): 29-34.
- Rani, D. S., Kumar, S. P., Venkatesh, M. N., Sri, C. H. N. S., and Kumar, K. A. (2018).** Bio efficacy of insecticides against gram pod borer, *Helicoverpa armigera* in Redgram. *Journal of Entomology and Zoology Studies*, **6**(2), 3173-3176.
- Singh, S.S. and S. K. Yadav. (2006).** Evaluation of chickpea varieties for their resistance against gram pod borer, *Helicoverpa armigera*. *Indian Journal of Entomology*, **68**(4): 321-324.
- Singh, H., Singh, H. R., Yadav, R. N., Yadav, K. G., and Yadav, A. (2009).** Efficacy and economics of some bio-pesticide in management of *Helicoverpa armigera* (HUB) on chickpea. *Pestology*, **33**(7), 36-37.
- Sanjayrao, J. A. (2013).** Studies on validation of molecular markers linked to wilt resistance, polymorphic information content (PIC) and transferability of functional markers in chickpea (*Cicer arietinum* L.) (Doctoral dissertation, University of

- Agricultural Sciences, GKVK).  
*The Bioscan*, **8**(4):1625-1930.
- Shah, J., Inayatullah, M., Sohail, K., Shah, S., Shah, S., Iqbal, T., and Usman, M. (2013).** Efficacy of botanical extracts and a chemical pesticide against tomato fruit worm, *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Sarhad Journal of Agriculture*, **29**(1), 93-96.
- Singh, P., Singh, R., Kumar, S., Kumar, V. and Kumar, S. (2015).** Bioefficacy of certain new insecticides against larval population of gram pod borer, *Helicoverpa armigera* (Hubner) in chickpea. *The Ecoscan*, **7**, 315-318.
- Sai, K. P., Tayde, A. R., Reddy, R. D., and Srivastava, V. K. (2021).** Management of gram pod borer *Helicoverpa armigera* (Hubner) in chickpea with bio-pesticides and combination. *Journal of Entomology and Zoology Studies*, **9**(1), 561-563.
- Visalakshmi, V., Rao, G. V. R. and Rao, P. A. (2005).** Integrated pest management strategy against *Helicoverpa armigera* Hubner in chickpea. *Indian Journal of Plant Protection*, **33**(1): 17-22.
- Wakil, W., Ashfaq, M., Kwon, Y. J., and Ghazanfar, M. U. (2009).** Trends in integrated pest management strategies for the control of *Helicoverpa armigera* (Hübner) caterpillars on chickpea (*Cicer arietinum* L.). *Entomological Research*, **39**: 84–88.
- Wakil, W., Ghazanfar, M. U., Riasat, T., Qayyum, M. A., Ahmed, S., and Yasin, M. (2013).** Effects of interactions among *Metarhizium anisopliae*, *Bacillus thuringiensis* and chlorantraniliprole on the mortality and pupation of six geographically distinct *Helicoverpa armigera* field populations. *Phytoparasitica*, **41**(2), 221-234.
- Wallace, T. C., Murray, R., and Zelman, K. M. (2016).** The nutritional value and health benefits of chickpeas and hummus. *Nutrients*, **8**(12), 766.