

Original Research Article

Assessment of projected climate change impact on green gram productivity through DSSAT model under South Gujarat environmental condition

ABSTRACT

An investigation on climate change impact on green gram was conducted at Agronomy farm, NAU, Navsari, Gujarat during the *rabi* seasons of 2021-22. The calibrated and validated DSSAT CROPGRO simulation model was used to determine the productivity of green gram under various climate change scenarios. In this experiment, the effects of CO₂ concentration, solar radiation, and maximum and minimum temperatures on green gram productivity were examined individually and in combination. According to the sensitivity of the DSSAT CROPGRO model, under the conditions of projected climate change, the best case scenario for its favourable impact on green gram yield was a rise in CO₂ concentration (600 ppm), along with solar radiation up to a certain level (1 unit), which may nullified the adverse effect of increases in temperature on green gram productivity.

Keywords: DSSAT, Crop modelling, Sensitivity analysis, Climate change, Green gram

1. INTRODUCTION

In the present century, climatic change [has](#) become a serious problem all over the world. IPCC (2021) has projected [that](#) an increase in global mean temperature of 1.5 °C was achieved before 2040 and according to Representative Concentration Pathways (RCP, 8.5), the global mean temperature is reported to increase by about 4.3 °C and CO₂ concentration is projected to increase up to 1200 ppm by end of the 21st century.

It is apparent that climate change will have an impact on a weather parameter and these weather parameters such as temperature, solar radiation, and rainfall has been determining the productivity of crop of a given locality. Looking at the projected climate change scenario, it is necessary to assess the productivity of green gram crops in relation to expected climate change conditions. The CROPGRO module of the DSSAT crop simulation model is a very useful tool to evaluate the expected yield of the crop in relation to various projected climate change scenarios because this tool can reduce the expense and time consuming of field experimentation (Kumar *et al.*, 2017).

Dhage and Patil (2020) studied the effect of elevated temperature on green gram yield was evaluated by using the CROPGRO model. The result showed that a rise in temperature by 1-2 °C had a negative impact on seed yield (declined by -3.56%). The sensitivity of the CROPGRO model

simulated the chickpea yield to incremental units of CO₂ resulting in a gradual increase in seed yield (Patil *et al.*, 2018).

Comment [d1]: Please mention the objectives of your study.

2. MATERIALS AND METHODS

2.1 General information

The field experiment was conducted during *rabi* season 2021-22 at Agronomy farm, NAU, Navsari, India situated at latitude 20°57' N, longitude 72°52' E and at an altitude of 12 m above mean sea level. The GBM cultivar of green gram was sown on three different dates (27th October (D₁), 11th (D₂) and 26th November (D₃)). The monthly means of weather data during the crop growing season (October to February) are presented in table 1.

Table 1: Monthly mean of weather condition during the growing season (October to March)

Month	Maximum temperature (°C)	Minimum temperature (°C)	Bright sunshine hours (hr.)	Solar radiation (MJ m ⁻² day ⁻¹)	Relative humidity (%) (morning)
October	33.6	17.1	9.0	19.13	73
November	33.7	18.2	7.4	15.98	69
December	29.2	14.7	6.2	13.55	86
January	28.9	14.0	7.2	15.37	91
February	31.1	13.7	9.3	20.03	78
March	36.3	17.3	8.5	20.70	67

2.2 Calibration and validation of DSSAT CROPGRO model for green gram

The first step in using the CROPGRO model to determine crop productivity under anticipated climatic circumstances is to perform model calibration and validation. The model was calibrated by using the past three years weather, soil, yield data of green gram and calculates a genetic coefficients of *cv.* GBM-1 by GLUE genetic coefficient estimator which is an inbuilt option in the model, followed by a manual method until the desired agreement made between simulated and observed values is reached. The calibrated genetic coefficient of *cv.* GBM-1 is presented in table 2. The model was validated with field experimental data (2021-22) and the accuracy of the model was tested with normalized root mean square error (nRMSE), which in our case was 8.02 percent.

Table 2: Calibrated of genetic coefficients of GBM-1 cultivar of green gram

Parameter	GBM-1
EM-FL (Time between emergence to flower appearance)	33.8
FL-SD (Time between first flower and first seed (R5))	18.1
FL-SH (Time between first flower and first pod (R3))	10.5

FL-LF (Time between first flower and end of leaf expansion)	13.0
SD-PM (Time between first seed and physiological maturity)	19.77
LFMAX (Maximum leaf photosynthesis rate at 30 °C, 350 ppm CO ₂ and high light)	0.99
WTPSD (Maximum weight per seed (g))	0.560
SLAVR (Specific leaf area of cultivar under standard growth conditions (cm ² /g))	130
SDPDV (Average seed per pod under standard growing condition ([seed]/pod))	10.5
SFDUR (Seed filling duration for pod cohort at standard growth condition)	24.3
PODUR (Time required for cultivar to reach final pod load under optimal condition)	18.0
XFRT (Maximum fraction of daily growth that is partitioned to seed + shell)	1.00
THRSH (The maximum ratio of (seed/seed+shell) at maturity)	78.0
SDPRO (Fraction protein in seed (g[protein]/g [seed]))	0.240
SDLIP (Fraction oil in seed (g[oil]/g[seed]))	0.055

2.3 Climate change study through DSSAT CROPGRO model

After validation CROPGRO model, sensitivity analysis (response of model to change input parameters) of CROPGRO model was used to study the impact of maximum temperature, minimum temperature, bright sunshine hours, CO₂ and solar radiation on the seed yield of green gram. The simulated value of seed yield under expected climate conditions was compared with a corresponding value of seed yield under normal weather conditions to measure the yield variation (%) under given climatic conditions. After calibration and validation of the model, the following step had been used for the climate change study: (1) Open DSSAT model. (2) Open the management file and select environment and go to environmental modification option. (3) Add input levels of a climatic parameter that you want to see the actual climatic parameter changes up to this level. (4) Go to the treatments option and assign environmental modification as treatment. (5) Refresh the model and run it.

3. RESULTS AND DISCUSSION

3.1 Effect of temperature on seed yield

The impact of increased and decreased temperatures (± 5 °C) on seed yield variation under the different sowing dates is presented in table 3. The results showed that the increase in maximum temperature had a negative impact (yield declined from -5.1 to -49.5 %) and the decrease in maximum temperature had a positive impact on the seed yield (rises from 4.7 to 35.2 %) due to the fact that decrease in temperature will increase duration of different phenological stages, which provides a much larger period for photosynthesis (Butterfield and Morison, 1992).

The sensitivity of CROPGRO model simulated the seed yield gradually increased when the minimum temperature was decreased up to -3 °C (Table 3), then yield was decreased with temperature decrease because the minimum temperature goes close to the base temperature of green gram where plant growth not possible. [Also/the](#) seed yield decreased with the [increase in](#) minimum temperature [increased](#) up to +5 °C because a higher rate of respiration during the night time was caused by the higher minimum temperature resulting in a comparatively higher loss of

photosynthates (Shamim *et al.*, 2010). The above presented result was good confirmation ~~of to~~ the findings of Kumar *et al.* (2017) and Yadav *et al.* (2016).

3.2 Effect of CO₂ on seed yield

The sensitivity of the CROPGRO model showed the positive impact of elevated CO₂ concentration on seed yield (from 17.52 to 49.27 %) (Table 3). The rise of seed yield under the increase in the concentration of CO₂ may be due to plant assured more entry of CO₂ in plant system and resulting in ~~plant exhibiting~~ the higher rate of photosynthesis. However, since CO₂ is a greenhouse gas, an increase in CO₂ concentration is responsible for the rising earth surface temperature. In this situation, ~~a the~~ combined studies of CO₂ and maximum temperature and its impact on seed yield ~~is are~~ necessary. The results of the combined experiments showed that the seed yield dropped (from 11.87 to -7.70%) as ~~combinedly both increase~~ CO₂ concentration and maximum temperature increased. It has been concluded that the seed yield ~~rises increased more as with~~ increased CO₂ concentration alone but seed yield was decreased with rises in both CO₂ concentration and temperature.

Table 3: % change of seed yield due to varying temperature, CO₂ and combined effect of CO₂ and maximum temperature under different sowing date

Change in climatic parameters	% Change yield from base seed yield		
	27/10/2021 (D ₁) (1376 kg ha ⁻¹ base yield)	11/11/2021 (D ₂) (1449 kg ha ⁻¹ base yield)	26/11/2021 (D ₃) (1461 kg ha ⁻¹ base yield)
MaxT (°C)			
-5	35.2	24.3	18
-4	31	28.4	22.1
-3	20.3	18.2	15.5
-2	14.5	16.1	11.2
-1	5.6	8.7	4.7
+1	-9.4	-5.1	-10.0
+2	-20.9	-15.2	-14.3
+3	-28.7	-22.0	-26.5
+4	-40.7	-32.9	-35.5
+5	-49.5	-41.9	-45.5
MinT (°C)			
-5	-5.6	-23.8	-26.2
-4	-0.2	-9.1	-13.0
-3	1.9	-6.0	-7.6
-2	5.4	1.3	-5.0
-1	4.7	3.3	4.0
+1	-2.9	-1.1	1.9
+2	-10.0	-8.8	-1.9
+3	-16.1	-13.5	-3.8
+4	-19.7	-22.2	-9.7

Comment [d2]: Mention the expanded form of the abbreviations in the text.

+5	-25.1	-25.9	-12.8
CO₂ concentration (Base value 380 ppm)			
450	19.1	17.5	17.5
500	30.6	27.6	27.7
550	40.6	36.9	36.9
600	49.2	44.6	44.6
CO₂ (ppm) + MaxT (°C) (Combine effect of MaxT and CO₂)			
450 + 1	8.5	11.8	5.3
500 + 2	4.8	9.9	11.0
550 + 3	2.3	8.8	2.2
600 + 4	-7.7	1.3	-3.4

3.3 Combined effect of solar radiation, temperature and CO₂ on seed yield

The seed yield variation (%) due to the combined effect of radiation, CO₂ and temperature is presented in table 4. The results showed that at a CO₂ concentration of 600 ppm, the green gram yield could be increased to a maximum of 9.96 to 69.77 percent by either increasing radiation and temperature by 3 units or decreasing it by 2 units. However, at a concentration of 450 and 500 ppm, the yield was found to be increased up to 1 unit when radiation and temperature were subtracted, and then it was found to decrease with more units subtracted. The highest benefits are obtained at the level of CO₂ concentration combined with 1 unit addition in temperature and radiation. A higher CO₂ concentration (600 ppm) may be able to reduce the negative effects of low radiation intensity (up to -2 units) and temperature increases on seed yield. Conversely, a low level of CO₂ (450 and 500 ppm) may be able to nullified the negative effects of low solar radiation (up to -1 units), but yield did not increase as much as it would with a higher CO₂ concentration. Our result was in good line with a finding of Pandey *et al.*, 2007.

Table 4: % change of yield due to interactive effect of solar radiation, temperature and CO₂ under different sowing date

Change in parameters	% Change yield from base seed yield		
	27/10/2021 (D ₁) (1376 kg ha ⁻¹ base yield)	11/11/2021 (D ₂) (1449 kg ha ⁻¹ base yield)	26/11/2021 (D ₃) (1461 kg ha ⁻¹ base yield)
450 ppm			
-3	-29.51	-28.85	-22.66
-2	-12.06	-9.25	-3.70
-1	2.11	5.24	5.68
+1	25.36	29.12	20.81
+2	27.11	32.23	32.92
+3	29.07	36.44	27.38
500 ppm			
-3	-23.04	-22.64	-15.74
-2	-3.85	-1.24	4.86
-1	12.06	14.63	15.13

+1	38.44	40.79	31.83
+2	39.61	44.24	44.56
+3	42.22	49.34	38.60
600 ppm			
-3	-12.50	-12.35	-4.11
-2	9.96	11.87	18.75
-1	28.20	29.88	30.60
+1	58.43	59.49	49.08
+2	59.81	63.77	62.90
+3	63.66	69.77	56.13

4. CONCLUSION

Calibrated and validated model of the DSSAT CROPGRO model is successfully used for simulating the seed yield of green gram under projected climate change scenarios. The model demonstrated that increased CO₂ concentration and ~~increased~~ solar radiation had a favourable influence on crop productivity while rising temperatures have a negative impact. The combined study found that increases in CO₂ concentration and maximum temperature have a detrimental impact on crop productivity whereas increases in CO₂ concentration along with solar radiation have a beneficial influence because ~~of the~~ nullified ~~the~~ negative impact of rises in maximum temperature. Overall, the beneficial effect of projected climate change on crop productivity can be harvested when solar radiation and temperature increase by 1 unit under all levels of CO₂ concentration.

REFERENCES

- 1) Butterfield RE and Morison JI. Modelling the impact of climatic warming on winter cereal development. *Agril. and Forest Meteorol.* 1992; 62(4): 241-261.
- 2) Dhage SS and Patil RH. Response of Greengram to Climate Change in Northern Transition Zone of Karnataka: DSSAT Model Based Assessment. *Legume Res. Int. J.* 2022; 1(5).
- 3) IPCC (2021). The physical science basis of climate change-A report of the Intergovernmental Panel on Climate Change. Fourth Assessment report.
- 4) Kumar S, Niwas R, Khichar ML, Kumar Y and Premdeep AS. Sensitivity analysis of DSSAT CROPGRO-Cotton model for cotton under different growing environments. *Ind. J. Ecology.* 2017; 44(4): 237-241.
- 5) Pandey V, Patel HR and Patel VJ. Impact assessment of climate change on wheat yield in Gujarat using CERES-wheat model. *J. Agrometeorol.* 2007; 9(2): 149-157.
- 6) Patil DD, Karande BI, Satpute SB and Patel HR. Sensitivity analysis to study the impact of climate change on chickpea using DSSAT (4.6) CROPGRO model over middle Gujarat Agroclimatic region. *A Peer-Review. Multi-Disciplinary Int. J.* 2018; 223.
- 7) Shamim M, Shekh AM, Pandey VY, Patel HR and Lunagaria MM. Sensitivity of CERES-Rice model to different environmental parameters on the productivity of aromatic rice in middle Gujarat. *J. Agrometeorol.* 2010; 12(2): 213-216.

Comment [d3]: Reference pattern

- 8) Yadav MK, Singh R. S, Singh KK, Mall RK, Patel C, Yadav SK and Singh MK. Assessment of climate change impact on pulse, oilseed and vegetable crops at Varanasi, India. *J. Agrometeorol.* 2016; 18(1): 13.