

# **A Study on Knowledge level of rice growers on Green Technologies in Rice based Ecosystem for Sustainable Agriculture in Madurai and Trichy district of Tamil Nadu**

## **Abstract:**

The present study was conducted in Trichy and Madurai district of Tamil Nadu. From each district two blocks were selected. Thus, the total sample size of 240 were selected for the present investigation. that higher percentage of the Trichy farmers had high level of knowledge towards green technologies (43.33%) followed by low (36.67%) and medium (20.00 %) level of knowledge towards green technologies. Meanwhile, half of the Madurai farmers had high level of knowledge (50.00 %) followed by low (36.67%) and medium (13.33%) level of knowledge towards green technologies. The  $R^2$  value indicated that all the variables contributed 59.60 % and 66.90 % variation in the knowledge level among the respondents of Trichy and Madurai district respectively.

**Key words :** Green technology, knowledge level, agricultural practices

## **2. Introduction**

An eco-friendly technology otherwise also known as green technology can be defined as the use of knowledge and resources in a systematic way to produce desired output without harming the environment. The world population is expected to increase over 10 billion by 2050. Such a growth in population has created and would create unprecedented pressures on the limited natural resources base to produce additional food, fiber, fuel and raw materials. The development and the transfer of eco-friendly technologies is interlinked with government extension agencies. Hence, it becomes necessary to study the efforts of the major promoters of agriculture in promoting eco-friendly technologies. Considerable attention is being paid to the eco-friendly technologies in different parts of the world. As the onus for promoting agricultural technologies is with the extension agencies, they must adapt themselves to the new environmental imperatives. In this point of view, agricultural extension is the educational

process of advising and assisting farm women in getting the best use of agricultural land and other natural resources in their care in the context of prevailing economical, technical, social and institutional conditions. The present study was undertaken to study the extent of knowledge level of recommended green technologies for paddy cultivation.

### **3. Materials and Methods**

The study was conducted in Trichy and Madurai districts of Tamil Nadu. From Madurai district two blocks namely Melur and Madurai east, from Trichy, Lalgudi and Thiruverambur blocks were selected and from each block three villages were identified for the study. A total of sample size of 240 farmers were selected from each village by using proportionate random sampling. For measuring knowledge, a Teacher made knowledge test was developed. A list of 25 knowledge items was constructed by using pertinent literature as well as extensive consultations with scientists and specialists. Judges opinion were obtained. The responses were binary in nature, such as 'Yes' and 'No'. Each accurate response received a score of 2 and each incorrect responses received a score of 1. By using cumulative frequency method, knowledge level of respondents were categorized into low, medium and high. Correlation and regression analysis was carried out to study the association and contribution of each variable on knowledge level for both the districts.

### **4. Results and discussion**

#### **4.1 Knowledge level of farmers utilizing green technologies in rice-based ecosystem**

The knowledge level of farmers utilizing green technologies in rice-based ecosystem in Madurai and Trichy district is studied and presented in table.1.

**Table.1. Knowledge level of Madurai and Trichy district farmers utilizing green technologies in rice-based ecosystem**

**(n = 240)**

S. No.	Statements	Response							
		Trichy (n=120)				Madurai (n=120)			
		Yes		No		Yes		No	
		No.	%	No.	%	No.	%	No.	%
1	The quantity of Azosprillum used per kg of seeds for seed treatment	108	90.00	12	10.00	100	83.33	20	16.67
2	Mention the ideal spacing of paddy cultivation	100	83.33	20	16.67	92	76.67	28	23.33
3	The recommended nursery area for one hectare of rice cultivation .	100	83.33	20	16.67	108	90.00	12	10.00
4	The recommended water level after panicle initiation to before maturation	4	3.33	116	96.67	64	53.33	56	46.67
5	How much quantity of FYM is required for paddy cultivation	100	83.33	20	16.67	108	90.00	12	10.00
6	Name any one crop that is used as green manure in paddy cultivation ?	12	10.00	108	90.00	100	83.33	20	16.67
7	What is the quantity of incorporation of green manure in paddy cultivation / ha?	16	13.33	104	86.67	76	63.33	44	36.67
8	Name one purpose of wood ash	28	23.33	92	76.67	28	23.33	92	76.67
9	Azolla is incorporated in the paddy field	12	10.00	108	90.00	40	33.33	80	66.67

	during which stage								
10	Mention the quantity of azospirillum that can be broadcast in paddy field	84	70.00	36	30.00	32	26.67	88	73.33
11	Mention one purpose of leaf colour chart	24	20.00	96	80.00	60	50.00	60	50.00
12	Mention one advantage of SRI technique	32	26.67	88	73.33	24	20.00	96	80.00
13	Name one method that is used to test the nutrients in the soil	40	33.33	80	66.67	28	23.33	92	76.67
14	Name one purpose of cono weeder	36	30.00	84	70.00	32	26.23	88	76.67
15	What is the proportion of raw materials in panchakavya and amirdhakaraisal?	60	50.00	60	50.00	84	70.00	36	30.00
16	How can be egg masses of stem borer be eliminated	32	26.67	88	73.33	28	23.33	92	76.67
17	Mention the stage at which clipping off rice seedling tips can be done	32	26.67	88	73.33	36	30.00	84	70.00
18	Trap effective against the infestation of yellow stem borer	28	23.33	92	76.67	32	26.67	88	73.33
19	During which stage paddy egg parasitoids is released	76	63.33	44	36.67	76	63.33	44	36.67

20	Mention the quantity of neem oil that can be sprayed /ha	80	66.67	40	33.33	88	73.33	32	26.67
21	The purpose of light trap installation	8	6.67	112	93.33	76	63.33	44	36.67
22	The number of bird perches that can be set up/ha	12	10.00	108	90.00	88	73.33	32	26.67
23	Name the purpose of bunding in paddy cultivation	80	66.67	40	33.33	80	66.67	40	33.33
24	The quantity of <i>Pseudomonas fluorescens</i> used as foliar spray	12	10.00	108	90.00	36	30.00	84	70.00
25	Mention the disadvantage of burning of crop residues in paddy field	32	26.67	88	73.33	32	26.67	88	73.33

From Table.1, it was observed that among the Trichy district farmers, most of them had knowledge on Azospirillum seed treatment in rice cultivation and its quantity to be used in seed treatment (90.00%). In the case of Madurai district farmers, 83.33 per cent of the paddy farmers had knowledge on azospirillum seed treatment. Majority of the farmers of Trichy district had knowledge on the recommended water level after panicle initiation to before maturation (3.33%) whereas more than half of the rice growers of Madurai district had knowledge (53.33%) on recommended water level in paddy cultivation.

Further examination of the table reveals that 83.33 and 90 per cent of Trichy district and Madurai district farmers were found to be knowledgeable on FYM required for paddy cultivation respectively. Meanwhile, it was found that 10.00 per cent of the farmers of Trichy district had knowledge on the crops which can be grown as a green manure in paddy cultivation. Among Madurai district farmers, 83.33 per cent knowledge towards usage of green manure in paddy cultivation. Around 13.33 per cent of the

farmers of Trichy district had knowledge on the quantity of green manure to be incorporated in paddy cultivation. Less than two-third of the Madurai rice growers(63.33%) were found to be knowledgeable on incorporation of green manure in paddy cultivation.

Similarly, majority of the Trichy farmers had no knowledge (90%) and knowledge (10%) of the azolla usage as a biofertilizer; whereas, among Madurai farmers, two-third of the farmers had no knowledge (66.67%) on azolla as biofertilizer. Higher percentage of the farmers of Trichy district had knowledge (70.00%) and no knowledge (30%) on azospirillum usage as biofertilizer respectively. Two-third of the Trichy paddy farmers (66.67%) and three-fourth of the paddy farmers in Madurai district had no knowledge (76.67%) on soil health card management.

Around thirty per cent of the farmers of Trichy district were found to be knowledgeable on useconoweeder for weeding in rice field. Among Madurai farmers, 26.23 per cent had knowledge on weeding in rice field. Exactly half of the Trichy district farmers had knowledge (50%) on the usage of Panchakavya and Amirdhakaraisal as plant growth regulators whereas, 70 per cent and 30 per cent of the Madurai district farmers had knowledge and no knowledge on usage of plant growth regulators. Similarly, 26.67 per cent of them had knowledge on the clipping of tip of the seedlings for elimination of stem borer egg mass remaining three-fourth of the farmers of Trichy district had no knowledge on this practice

Regarding the effective trap for yellow stem borer, 76.67 and 26.67 per cent of Trichy and Madurai rice growers were found to be knowledgeable. With reference to the farmer's knowledge level regarding the egg parasitoids, 63.33 per cent and 36.67 per cent of Trichy and Madurai district farmers had knowledge and no knowledge on it.

Eventually, 10.00 per cent of the Trichy farmers had knowledge and no knowledge on the installation of bird perches for destruction of leaf folder respectively; 73.33 per cent of the Madurai farmers had knowledge. Most of the farmers had no knowledge (90.00%) and knowledge (10.00%) on the usage of *Pseudomonas fluroscensas* foliar spray. Similarly, a little less than three-fourth of the Madurai farmers

had no knowledge (70.00%) and knowledge (30.00%) on usage of *Pseudomonas fluroscensas* foliar spray.

Based on the findings, it can be observed that farmers were found to have high level of knowledge on organic farming, increased social and environmental concern regarding good and healthy food motivate the farmers to adopt organic and eco-friendly practices. Simultaneously, increased level of information sources, higher level of extension participation and extension contact inspired the farmers to adopt eco-friendly practices; which in turn increased their knowledge level towards green technologies in paddy cultivation. Further, contrary findings were reported by Rathod et.al., (2017), Suji et.al., (2017), Rahaman et.al., (2018) and Balamurugan et.al., (2021).

#### 4.2. Overall knowledge level of farmers utilizing green technologies

The overall knowledge level of farmers utilizing green technologies in rice-based ecosystem in Madurai and Trichy district is studied and presented in Table.2

**Table.2. Overall knowledge level of Madurai and Trichy district farmers utilizing green technologies in rice-based ecosystem  
n=240**

S. No.	Overall knowledge level of green technologies	Respondents – Farmers					
		Trichy		Madurai		Total	
		F	%	F	F	%	F
1	Low	44	36.67	44	36.67	88	36.67
2	Medium	24	20.00	16	13.33	40	16.66
3	High	52	43.33	60	50.00	112	46.67
Total		120	100.00	120	100.00	240	100.00

From Table..2,it was revealed that higher percentage of the Trichy farmers had high level of knowledge towards green technologies (43.33%) followed by low (36.67%) and medium (20.00 %) level of knowledge towards green technologies. Meanwhile, half

of the Madurai farmers had high level of knowledge (50.00 %) followed by low (36.67%) and medium (13.33%) level of knowledge towards green technologies.

On contrary to findings of Rathod *et.al.*, (2017), Suji *et.al.*, (2017), Rahaman *et.al.*, (2018) and Balamurugan *et.al.*, (2021), from the present study, the farmers of Madurai and Trichy district were found to have high level of knowledge towards green technologies. Meanwhile, similar findings were reported by Suji *et.al.*, (2017) and Upadhyay *et.al.*, (2020a). Eventually, Singh *et.al.*, (2013) identified higher level of knowledge gap among the farmers and Masso and Man (2016) &Rahaman *et.al.*, (2018) suggested to intensify the awareness and knowledge level of farmers towards green technology practices.

#### **4.3.Relationship between the profile characteristics of Trichy and Madurai farmers and their knowledge level on green technologies**

To find out the relationship between the dependent variable knowledge and the selected independent variables, simple correlation and multiple regression analysis were worked out separately for Trichy and Madurai and the results are presented in Table 3.

**Table.3. Correlation analysis of profile characteristics of Trichy and Madurai farmers with their knowledge level on green technologies**

<b>S.No</b>	<b>Characteristics</b>	<b>'r' value Trichy farmers</b>	<b>'r' value Madurai farmers</b>
X <sub>1</sub>	Age	-0.350**	0.087
X <sub>2</sub>	Education	0.240**	0.097
X <sub>3</sub>	Occupation	-0.169	0.222*
X <sub>4</sub>	Farm size	0.022	-0.165
X <sub>5</sub>	Farming experience	-0.241**	-0.037
X <sub>6</sub>	Annual income	-0.128	-0.061
X <sub>7</sub>	Information seeking behavior	-0.227*	-0.224*
X <sub>8</sub>	Extension participation	-0.015	-0.161
X <sub>9</sub>	Social participation	-0.681**	-0.673**
X <sub>10</sub>	Innovativeness	-0.011	-0.171
X <sub>11</sub>	Economic motivation	-0.045	-0.149
X <sub>12</sub>	Perception on use of green	-0.015	-0.312**

	technology		
X <sub>13</sub>	Perception on environmental degradation	-0.212*	0.295**
X <sub>14</sub>	Perception on feasibility of green technology agricultural practices	-0.011	0.075
X <sub>15</sub>	Perception on use of synthetic fertilizers	0.008	0.224*

In the case of Trichy district, it could be observed from Table 3, that out of fifteen variables analysed, six variables have established significant association with knowledge on green technologies. The variable education(X<sub>2</sub>) was found to be significant and positively associated at one percent level of probability. The variables age (X<sub>1</sub>), farming experience (X<sub>5</sub>) and social participation(X<sub>9</sub>) were found to negatively significant and associate with knowledge level on one per cent level of probability. Information seeking behavior (X<sub>7</sub>) and perception on environmental degradation innovativeness (X<sub>13</sub>) have been found to be significant and negatively associated with knowledge level at five percent level of probability.

From Table.3, it was reported that education status of the Trichy farmers had positive and significant association with their knowledge level on green technologies at 1 per cent level of probability. Since education status of a farmer increases, they learn new things and their knowledge level increases which significantly increases their knowledge level on green technologies.

From Table.3, it revealed that information seeking behaviour of the Trichy farmers had negative and significant association with their knowledge level on green technologies at 5 per cent level of significance. Since information seeking behaviour of farmers on chemical fertilizers increases, their knowledge on green technologies decreases.

From Table.3, it was evident that social participation of the Trichy farmers had negative and significant association with their knowledge level on green technologies at 1 per cent level of significance. Increased social participation of farmers hinders their

participation in agricultural activities; thus, it limits the knowledge level of the farmers on green technologies.

From Table.3, it was observed that perception on environmental degradation of the Trichy farmers had negative and significant association with their knowledge level on green technologies at 5 per cent level of significance. Since the observability of the relative advantage of the green technology is relatively low than chemical fertilizers, farmers were not aware of the harmful effects of chemical fertilizers in deteriorating and polluting the environment.

In the case of Madurai district, it could be observed from Table 3, that out of fifteen variables analysed, six variables have established significant association with knowledge on green technologies. The variable perception on environmental degradation innovativeness ( $X_{13}$ ) was found to be significant and positively associated at one percent level of probability.

Variables namely occupation( $X_3$ ) and Perception on use of synthetic fertilizers( $X_{15}$ ) was found to be significant and positively associated at five percent level of probability whereas variables social participation( $X_9$ ) and Perception on use of green technology( $X_{12}$ ) were found to significant and associated negatively with knowledge level on one per cent level of probability. Information seeking behavior ( $X_7$ ) have been found to besignificant and negatively associated with knowledge level at five percent level of probability.

**Table. 4. Multiple regression analysis of profile characteristics of Trichy and Madurai farmers with their knowledge level on green technologies**

S.N o	Characteristics	Trichy			Madurai		
		Regressio n co- efficient	Standard error	t- value	Regressi on co- efficient	Standard error	t- value
X <sub>1</sub>	Age	-0.112	0.097	-1.153	-0.269	0.186	-1.446

X <sub>2</sub>	Education	0.201	0.034	5.890	0.070	0.069	1.018
X <sub>3</sub>	Occupation	-0.136	0.052	-2.607	0.189	0.065	2.900
X <sub>4</sub>	Farm size	0.750	0.074	10.19 6	-0.272	0.118	-2.310
X <sub>5</sub>	Farming experience	0.055	0.066	0.840	0.145	0.114	1.269
X <sub>6</sub>	Annual income	0.149	0.094	1.585	0.124	0.137	0.905
X <sub>7</sub>	Information seeking behaviour	-0.404	0.099	-4.100	-0.640	0.147	-4.362
X <sub>8</sub>	Extension participation	0.282	0.082	3.428	-0.211	0.122	-1.727
X <sub>9</sub>	Social participation	-1.234	0.075	- 16.48 7	-0.831	0.130	-6.410
X <sub>10</sub>	Innovativeness	0.133	0.064	2.061	-0.200	0.102	-1.955
X <sub>11</sub>	Economic motivation	-0.566	0.092	-6.163	-0.395	0.132	-3.000
X <sub>12</sub>	Perception on use of green technology	0.237	0.100	2.363	-0.360	0.123	-2.922
X <sub>13</sub>	Perception on environmental degradation	-0.170	0.081	-2.104	0.219	0.142	1.538
X <sub>14</sub>	Perception on feasibility of green technology agricultural practices	0.050	0.082	0.616	0.087	0.124	0.705
X <sub>15</sub>	Perception on use of synthetic fertilizers	0.295	0.089	3.300	0.231	0.130	1.778

\* = Significant at 5% \*\* = Significant at 1%

#### Trichy district

R<sup>2</sup> = 0.596  
1.838

F=32.89

a=

#### Madurai district

$R^2 = 0.669$	$F=14.005$	$a=$
5.155		

The contribution of each characteristic towards knowledge level of Trichy farmers on green technologies was examined using multiple regression analysis and the results are presented in table 4. It indicated that co-efficient of determination ( $R^2$ value) for the farmers was 0.596, which depicts that all the fifteen independent variables chosen for the study together explained for 59.60 per cent of variation in the knowledge level of farmers on green technologies. The prediction equation fitted for knowledge level of farmers on green technologies is given below.

$$Y1=1.838 - 0.350^{**} (X_1) + 0.240^{**} (X_2) - 0.169 (X_3) + 0.022 (X_4) - 0.241^{**} (X_5) - 0.128 (X_6) - 0.227^* (X_7) - 0.015 (X_8) - 0.681^{**} (X_9) - 0.011 (X_{10}) - 0.045 (X_{11}) - 0.015 (X_{12}) - 0.212^* (X_{13}) - 0.011 (X_{14}) + 0.008 (X_{15})$$

In the case of Madurai district, co-efficient of determination ( $R^2$ value) for the farmers was 0.669, which depicts that all the fifteen independent variables together explained for 66.90 per cent of variation in the knowledge level of the farmers on green technologies. The prediction equation fitted for knowledge level of the farmers on green technologies is given below.

$$Y2=5.155 + 0.087 (X_1) + 0.097 (X_2) + 0.222^* (X_3) - 0.165 (X_4) - 0.037 (X_5) - 0.061 (X_6) - 0.224^* (X_7) - 0.161 (X_8) - 0.673^{**} (X_9) - 0.171 (X_{10}) - 0.149 (X_{11}) - 0.312^{**} (X_{12}) + 0.295^{**} (X_{13}) + 0.075 (X_{14}) + 0.224^* (X_{15})$$

From Table.4, it was observed that occupation of Madurai farmers had positive and significant association with their knowledge level on green technologies at 5 per cent level of significance. Since, occupation of the majority of the farmers is farming alone, they tend to increase their knowledge on improved agricultural practices to keep them updated in their occupation.

From the above table, it was observed that information seeking behaviour of Madurai farmers had negative and significant association with their knowledge level on green technologies at 5 per cent level of significance. As information source of farmers might vary and may prefer the usage of chemical fertilizers, which affects the knowledge level on green technologies in a negative way.

From Table.4, it was reported that perception on environmental degradation of the Madurai farmers had positive and significant association with their knowledge level on green technologies at 1 per cent level of significance. As farmers were not aware of the harmful effects of synthetic fertilizers, as they concern for increased yield and income; their perception on environmental degradation might not improve their knowledge on green technology.

From the above Table.4, it was observed that perception on use of agricultural practices of the Madurai farmers had positive and significant association with their knowledge level on green technologies at 5 per cent level of significance. Though, farmers prefer chemical fertilizers for quicker results, their concern on field fertility and longterm productivity might influence their knowledge level on green technology

### **3. Conclusion**

The study revealed that majority of the respondents of both Trichy and Madurai district came under higher knowledge level on green technologies. The policy makers should arrange training programmes at farm level to increase the knowledge level of farmers on green technologies. Extension professionals may concentrate to impart skill demonstration on various aspect like bio input , use of various traps, encouraging minimal use of chemicals. The green technologies can ensure healthy, culturally compatible and sustainable life. Hence efforts need to be taken by the extension officials to popularize the importance of these practices.

### **4. References**

- Balamurugan V., V. Kalirajan, A. Thirumal. (2021). A Study on Practice - Wise Knowledge Level of the Paddy Farmers about the Recommended Biofertilizers

Practices in Paddy Cultivation in Vellore District. *Annals of the Romanian Society for Cell Biology*, 25(6),12376–12382.

Retrieved from <https://www.annalsofrscb.ro/index.php/journal/article/view/7865>

- Masso, W. Y. A., & Man, N. (2016). Identify the Knowledge Level of Rural Leaders towards Paddy Farming Technologies in Muda Agriculture Development Authority. *Indian Journal of Science and Technology*, 9(15).
- Rahaman, M. M., Islam, K. S., & Jahan, M. (2018). Rice farmers' knowledge of the risks of pesticide use in Bangladesh. *Journal of Health and Pollution*, 8(20).
- Rahman, M.S., Majumder, M.K. Drivers of adoption and impacts of an eco-friendly agricultural technology in Bangladesh. *SN Bus Econ* 1, 162 (2021). <https://doi.org/10.1007/s43546-021-00165-3>.
- Rathod, M.K., Kad, S.B., and Koshti, N.R. 2017. Adoption behaviour of farmers towards bio-fertilizers inn paddy. *International Journal of Extension Education*. 13:83-88.
- Suji, B.D., Kavaskar, M., Sathish kumar, A.M. 2017. Knowledge level of the farmers in eco-friendly agricultural technologies in paddy cultivation. *EPRA International Journal of Economic and Business Review*. 5(5):57-61.
- Suji, B.D., and Sathish Kumar, A.M. 2020. A study on adoption of ecofriendly technologies and constraints. *Plant Archives*. 20(1):42-46.
- Upadhyay, V., Khare, N. K., &Naberia, S. 2020a. Attributes of tribal paddy growers using eco-friendly practices in Mandla district, Madhya Pradesh. *Gomal University Journal of Research*. 29(2):
- Upadhyay, V., Khare, N. K., &Naberia, S. 2020b. Utilization of Eco-friendly practices of paddy crop by tribal farmers of Mandla district, Madhya Pradesh. *Journal of Pharmacognosy and Phytochemistry*. 9(2):183-185.