

# **Effect of Seaweed Extracts in Fruit Crops: A Review**

## **ABSTRACT**

Seaweed or Marine macro-algae are often regarded as an underutilized bioresource, many have been used as a source of food, raw materials for industry, and therapeutic and botanical purposes for decades. Moreover, several plant growth-stimulating compounds found in seaweed and products derived from seaweed have made them popular supplements to crop production systems. This article provides an in-depth analysis of how different seaweed species and seaweed products affect plant growth and development of fruit crops. They also increase the biochemical components of plants and have ability to alleviate abiotic stress.

*Keywords: Seaweed; plant growth-stimulating compounds; plant growth and development; biochemical components, abiotic stress*

## **1. INTRODUCTION**

Due to the rising need for food and fiber per unit of land area, chemical fertilizers are becoming increasingly popular to increase yields from small plots of land. Chemical fertilizers pose major health risks and harm the environment when they are used in excess. Because of this, research in this area is heavily focused on finding and evaluating the efficacy and efficiency of new products. One such approach is the use of biostimulants that can enhance the effectiveness of conventional mineral fertilizers [1]. Biostimulants are loosely defined as non-fertilizer products which have beneficial effects on plant growth and didn't contain any chemicals or synthetic plant growth regulators [2]. Nearly 10,000 different species of macroalgae (seaweeds) exist and account for 10% of marine productivity globally. Seaweeds are quintessential members of inshore, marine ecosystems as they provide shelter and food to numerous marine biota and can even contribute to the modification of physicochemical properties of seawater. Only a small part of all seaweed species is used extensively in agriculture as mulches, manures, and modified extracts, as well as food and supplements for humans and animals [3,4,5].

The earliest evidence of the use of seaweed by humans' dates back 15,000 years and was discovered in organic remains excavated from prehistoric sites in Monte Verde, Southern Chile, where an ancient migratory population of maritime origin had been established [6]. Seaweeds have been utilized by humans for a variety of purposes, including food, medicine, agriculture, cosmetic items, colouring pigments, and textiles, according to the available resources, which range from ancient scriptures and folklore to anthropological studies. Various seaweeds have found their places in diverse cultures throughout the globe [7]. Deity Ukimochi, the Goddess of food, is depicted to be associated with seaweed in Japanese mythology [8]. The most famous author of Roman agriculture, Columella, described how roots were to be wrapped in "seaweed" (genus not specified) to preserve the greenness of the seedlings, demonstrating that the ancient Romans were aware of the fertilizing properties of seaweeds [9]. In the British Isles, including Scotland and Ireland, different seaweeds have been used as manure more recently [10]. Seaweeds are macroscopic marine algae. They serve as human food, livestock feed, a source

of numerous fine chemicals, and a replacement for chemical fertilizer. In addition, it is employed to produce a variety of industrial goods, including agar and alginate [11]. Natural seaweeds have begun to take the place of synthetic fertilizers in recent years. Several commercial seaweed extract products are available for use in fruit crops [12] (Table 1). Because they contain numerous growth regulators, including cytokinins [13], auxins [14], gibberellins [15], and various macro and micronutrients required for plant growth and development, seaweed extracts are marketed as liquid fertilizers and bio-stimulants. Additionally, it facilitates the development of advantageous soil microbes [16], builds environmental stress tolerance [17], increases nutrient uptake from soil [18], and improves antioxidant characteristics [19]. Numerous studies have shown that applying seaweed extract to plants has a wide range of advantageous effects, including promoting early seed germination and establishment, increasing crop performance and yield, boosting resistance to biotic and abiotic stress, and extending the post-harvest shelf life of perishable goods [20,21,22] (Fig. 1). In this review we reported the effect of seaweed extracts as plant biostimulants in fruit crops and the possible modes of action including the chemical components that may be responsible for causing plants to respond physiologically to treatment.

Table 1. Commercial seaweed products used in fruit crops

<b>Product name</b>	<b>Seaweed name</b>
Acadian	<i>Ascophyllum nodosum</i>
Acid Buf	<i>Lithothamnium calcareum</i>
Agri-A-Mic	<i>Ascophyllum nodosum</i>
Bio-Genesis™ High Tide™	<i>Ascophyllum nodosum</i>
Biovita	<i>Ascophyllum nodosum</i>
Emerald RMA	Red marine algae
Fartumm	Unspecified
Guarantee	<i>Ascophyllum nodosum</i>
Goeman GA 14	<i>Ascophyllum nodosum</i>
Kelp Meal	<i>Ascophyllum nodosum</i>
Kelpak	<i>Ecklonia maxima</i>
Kelpro	<i>Ascophyllum nodosum</i>
Maxicrop	<i>Ascophyllum nodosum</i>
Profert, Seasol	<i>Durvillea potatorum</i>
Soluble Seaweed Extract	<i>Ascophyllum nodosum</i>
Simplex, Tasco	<i>Ascophyllum nodosum</i>

## **METHODS OF SEAWEED FARMING IN INDIA**

### **2.1 Floating bamboo raft method**

The principal structure for cultivation is an 8–10 cm wide by 3–3 m long bamboo raft. To keep it intact, the angular pieces are diagonally fastened with the aid of support bamboos (about 120 cm long). To fix the rafts and keep them buoyant, an anchor is attached around each group of rafts. To reduce grazing and prevent debris drift, bottom netting is offered. Total 20 plantings of about 100–150 g fresh weight each are planted on polypropylene ropes (3 mm) at regular intervals by raffia or braider. The seeded ropes are fastened to opposing bamboo at both ends, 15 cm apart and parallel to each other. Each raft with 20 such ropes thus will have an initial seeding weight of 60 kg fresh weight (Fig. 2a). The seeding is carried out onshore and seeded rafts are transplanted into the open sea subsequently. After 45 days, harvesting is often done. It is advised to use this technique in locations with calm water currents, such as the North Ramanathapuram district of Tamil Nadu [23].

## **2.2 Longline or monoline method**

There are a few similarities between this modified form of traditional off-bottom farming and the raft approach. Between anchors, seedlings weighing between 100 and 150 g are tied to a rope with raffia or braider. The main rope has a diameter of 6 to 8 mm and is roughly 20 meters long. The longline ropes are kept afloat in water with the help of floats tied at regular intervals and anchoring is done on both ends. The timber of casuarina, eucalyptus, bamboo, etc. is used for anchoring. In a square pattern, poles are erected at 3-m intervals, and 12-mm polypropylene rope is linked in any two parallel directions (depending upon tidal current). The 3 mm seeded ropes are then tied at intervals of 10 cm. It is important to take precautions to guarantee that the seaweed is always submerged (0.5 m below the surface) and gets enough sunshine (Fig. 2b). To prevent seedling loss that can occur if ropes are seeded on the coast and dragged into the sea, seeding in this situation is typically done in the water itself. This farming method is especially advised in regions with low grazer densities and moderate wave action. In Tamil Nadu's South Ramanathapuram, Pudukkottai, and Tuticorin districts, it is widely practiced [24].

## **2.3 Tube net method**

Higher wave action may result in crop loss at the preharvest stage, when seeding is at its heaviest, or drifting of germlings during the early growth phase. So, the tube net method has a lot of potential [25]. The tube nets (15-cm diameter; mesh size of 2.0 cm) of 25-m length are held floating in the water column below the surface with an appropriate number and size of floats at regular intervals. Anchor stones (about 30 kg) are used at each end to hold the tube nets steady in the water column; if required, additional anchors of appropriate size and weight can be fixed intermediately. The seed material of 20 kg fresh weight is loaded into the tubes with the aid of a 1.0- or 1.5 m long plastic pipe acting as a funnel or a hopper. For effective seeding, the pipe diameter should be a bit less than that of the tube network. Once the plastic pipe is installed, the entire tube is dragged downward until the plastic pipe's mouth protrudes from the tube. To load the seedling material into the tube sequentially and without gaps between the seedlings, the tube net is carefully pushed down from the bottom of the plastic pipe. This procedure is repeated until algal biomass has been planted throughout the entire tube net. The tubes are closed at both ends with rope to prevent material from being lost (Fig. 2c). Recently, a mechanical harvester was created and put through testing to harvest the biomass from the tubes [26].

## **3. GROWTH STIMULATORY FACTORS IN SEAWEED EXTRACTS AND THEIR METHODS OF ACTION**

The use of seaweed formulations as biostimulants in crop production is well recognized, and seaweed products demonstrate growth-stimulating properties. Biostimulants, also known as "metabolic enhancers," are substances that "boost plant development when administered in tiny

doses, rather than fertilizers" [27]. Seaweed components such as macro- and microelement nutrients, amino acids, vitamins, cytokinins, auxins, and abscisic acid (ABA)-like growth substances affect cellular metabolism in treated plants leading to enhanced growth and crop yield [28,29,30,31,32,33]. At low concentrations (diluted as 1:1000 or higher), seaweed extracts are bioactive [34]. It is conceivable that the numerous chemical components of seaweed extracts exhibit synergistic activity, even though many of these components' actions and modes of action are still unknown [35,36].

#### **4. EFFECTS ON ABIOTIC STRESS ALLEVIATION IN FRUIT CROPS**

Abiotic stressors like drought, salinity, and temperature extremes can lower the yield of important crops and restrict agricultural production globally [37]. For instance, drought and salinity are both spreading in many parts of the world, with 50% of all arable lands likely becoming salinized by 2050 [38]. Many abiotic variables, including temperature, salt, and drought, manifest as osmotic stress and have downstream consequences that include oxidative stress, which results in an accumulation of reactive oxygen species (ROS), including hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and the superoxide anion (O<sub>2</sub> • -) [39]. These are recognized to cause harm to proteins, lipids, carbohydrates, and DNA as well as to produce abnormal cell signalling [40]. Plants perform better under abiotic stressors due to bioactive chemicals found in seaweed extracts. Spraying extracts on plants has been found to increase their resistance to the stress caused by freezing temperatures [41]. Applying an *A. nodosum* extract formulation specifically improved the ability of grapes to withstand freezing because it reduced the leaves' osmotic potential, which is a crucial sign of osmotic tolerance. After nine days of seaweed extract treatment, the average osmotic potential of the treated plants was -1.57 MPa as opposed to -1.51 MPa in the untreated controls [42]. Spinelli et al. (2010) [43] examined the impact of another commercial seaweed extract, named Actiwave® on the vegetative and productive performance of strawberry plants grown on an iron-deficient substrate. They observed that vegetative growth, chlorophyll content, stomatal density, and photosynthetic rate were increased after biostimulant application. Fruit production and weight were also enhanced. Nutrient uptake might have been positively influenced by the more developed root system of treated plants. Treatment also contrasted the negative effects of iron chlorosis and this could be linked to betaine contained in this product. Two seaweed-based plant biostimulants containing *Ascophyllum nodosum* named Super Fifty® and Acadian were applied respectively on strawberry and was associated with a significant improvement in yield and root dry weight, despite the adverse salinity condition [44].

#### **5. EFFECT ON PLANT METABOLISM AND PHYSIOLOGY OF FRUIT CROPS**

One of the characteristic impacts of seaweed extract treatment is an increase in the chlorophyll content in the treated plants; this result has been seen in a variety of crops, including grapevine and strawberry [45,46,47,48].

Under nutrient-deprived conditions, the application of a commercial seaweed extract to apple trees of the variety Fuji reduced the problem of alternate bearing, but not under normal nutrient management conditions [49]. Application of plant growth regulators such as gibberellin, auxin, and cytokinin can decrease alternate bearing. Accordingly, it would seem that the influence of commercial seaweed extracts prepared from *A. nodosum* on alternate bearing under nutrient-deprived conditions may be caused, at least in part, by the presence of a hormone or a signalling molecule in the extract. However, it is challenging to explain why the commercial *A. nodosum* extract only reduces alternate bearing in conditions of inadequate nutrition [50,51].

#### **6. EFFECT ON GROWTH AND YIELD OF FRUIT CROPS**

In many crop plants, seaweed concentrate induces early flowering and fruit set [52,53,54]. For example, Roshdy (2014) [55] recommended the use of potassium silicate in combination with seaweed extract both applied four times at 0.05 % in the Grand Naine variety of bananas for higher bunch weight, average hand weight, finger weight, finger length, and finger diameter. Foliar application of seaweed extracts at 0.2% recorded maximum no. of fruits/tree, fruit weight, and yield/tree in valencia orange [56]. Seaweed extract had a higher amount of essential nutrients, amino acids, vitamins, antioxidants, and natural hormones [57]. The hormonal components found in the extracts, particularly cytokinins, are assumed to be responsible for the increased yield in plants treated with seaweed [58,59,60]. High amounts of cytokinins in reproductive organs may be related to nutrient mobilization, whereas cytokinins in vegetative plant organs are involved with nutrient partitioning. Fruit ripening generally causes an increase in the transport of nutrient resources within the developing plant [61,62]. and the fruits can serve as strong sinks for nutrients [63]. According to Norrie and Keathley (2006) [64]. over three years, *A. nodosum* extracts consistently increased the production of "Thompson seedless" grapes (*Vitis vinifera* L.). They noticed that the *A. nodosum*-treated plants always outperformed (in terms of berries per bunch, berry size, berry weight, rachis length, and the number of primary bunches per plant) the controls maintained under the regular crop management program. This led to improved fruit size (13% increase), weight (39% increase), and yields (60.4% increase over the control).

## **7. EFFECT ON QUALITY AND SHELF LIFE OF FRUIT CROPS**

*A. nodosum* extract increased the productivity, quality, and nutrient status of olives [65]. Foliar spray of *A. nodosum* extracts along with root application of nitrogen and boron improved K, Fe, and Cu concentrations in the leaves with a concomitant decrease of Mn. The seaweed extract treatment caused changes in the fatty acid profile of the olive oil; there was a significant improvement in the concentration of linolenic and oleic acid and a considerable decrease in palmitoleic, stearic, and linoleic acid. In addition to increasing productivity and quality, using seaweed extract prolonged the shelf life of pears and avocados [66,67]. Additionally, post-harvest treatments have been employed with seaweed extracts. When held at room temperature or in cold storage, post-harvest treatment of navel oranges with seaweed extract including a combination of Sargassum, laminaria, and *A. nodosum* extracts dramatically increased post-harvest shelf-life and quality [68].

## **8. CONCLUSION AND PROSPECTS**

Seaweeds and seaweed products are increasingly used in crop production. However, it is currently uncertain what mechanism(s) seaweed extract uses to cause its physiological effects. To better understand the mechanisms of action of the seaweed-induced growth response and stress reduction, it is possible to examine the effects of seaweed extracts and components on the entire genome/transcriptome of plants. This is possible because the genomes of many plants have now either been fully sequenced or are almost there. For instance, using model plants like *Arabidopsis thaliana* and *Medicago truncatula* might help elucidate the molecular mechanism(s) underlying the effects of seaweed extracts [69]. Climate change is likely to be to blame for the recent challenges to food production brought on by the rising incidence of biotic and abiotic stresses, which will further diminish yields and/or have an effect on crops in the twenty-first century (IPCC 2007) [70]. Therefore, it should be a top priority to research to create sustainable means of reducing these strains. Recent research has demonstrated that seaweed extracts made from different starting raw materials, and by different procedures are attributed to several beneficial effects such as increased nutrient uptake, abiotic stress tolerances, and improvement in the growth, yield, and quality of fruit crops. Seaweed extracts are also regarded as an organic agricultural input due to their safety for both human and animal health as well as the environment.

## REFERENCES

1. Frankenberger WT, Arshad, M. Phytohormones in soils. MarcelDekker, New York.1995.
2. Russo, RO, Berlyn, GP. The use of organic biostimulants to low input sustainable agriculture. J. of Sust. Agriculture. 1990;1:9–42.
3. Craigie JS. Seaweed extract stimuli in plant science and agriculture. J. Appl. Phycol. 2011;23:371–393.
4. Rönnbäck P, Kautsky N, Pihl L, Troell M, Söderqvist T, Wennhage H. Ecosystem goods and services from Swedish coastal habitats: identification, valuation, and implications of ecosystem shifts. *Ambio*. 2007;36:534–544.
5. Rayorath P, Benkel B, Hodges DM, Allan-Wojtas P, MacKinnon S, Critchley AT, Prithiviraj B. Lipophilic components of the brown seaweed, *Ascophyllum nodosum*, enhance freezing tolerance in *Arabidopsis thaliana*. *Planta*. 2009;230:135–147.
6. Dillehay TD, Ramirez C, Pino M, Collins MB, Rossen J, Pino-Navarro JD. Monte Verde: seaweed food, medicine, and the peopling of S. Am. *Sci*. 2008;320:784–789.
7. Guiry MD, Blunden G. Seaweed Resources in Europe: Uses and Potential. John Wiley and Sons Ltd. 1991.
8. Coulter CR, Turner P. Encyclopedia of Ancient Deities. Routledge. 2013
9. Henderson J. The Roman Book of Gardening. Psychology Press. 2004.
10. Wells E, Wilkinson M, Wood P, Scanla C. The use of macroalgal species richness and composition on intertidal rocky seashores in the assessment of ecological quality under the European water framework directive. *Mar. Pollut. Bull*. 2007;55:151–161.
11. Wang T, Jonsdottir R, Ólafsdóttir G. Total phenolic compounds, radical scavenging and metal chelation of extracts from Icelandic seaweeds. *Food Chem*. 2009;116:240–248.
12. Khan W, Rayirath UP, Subramanian S, Jithesh MN, Rayorath P, Hodges DM, Critchley AT, Raigie JS, Norrie J, Prithiviraj B. Seaweed extracts as biostimulants of plant growth and development. *J. Plant Growth Regul*. 2009;28:386-399.
13. Durand, Brian X, Meyer C. The effect of marine bioactive substances (N PRO) and endogenous cytokinins on nitrate reductase activity in *Arabidopsis thaliana*. *Physiol. Plant*. 2003;119:489-493.
14. Sahoo D. Farming the ocean. In: Seaweeds Cultivation and Utilization. Aravali Books International. New Delhi, India, 2000, 40.
15. Strik W, Staden VJ. Isolation and identification of cytokinins in a new commercial seaweed product made from *Fucus serratus* L. *J App Phycol*. 1997;9: 327-330.
16. Craigie JS. Seaweed extract stimuli in plant science and agriculture. *J. Appl. Phycol*. 2011;23:371–393.
17. Zhang X, Ervin EH, Schmidt, ER. Plant growth regulators can enhance the recovery of Kentucky bluegrass sod from heat injury. *Crop Sci*. 2003;43:952–956.
18. Turan M, Kose C. Seaweed extracts improve copper uptake of grapevine. *Acta. Agric. Scand*. 2004;54:213-220.
19. Verkleij FN. Seaweed extracts in agriculture and horticulture: a review. *Biol. Agricul. Horticul*. 1992;8:309–324.
20. Beckett RP, van Staden J. The effect of seaweed concentrate on the growth and yield of potassium stressed wheat. *Plant Soil*. 1989;116:29–36.

21. Hankins SD, Hockey HP. The effect of a liquid seaweed extract from *Ascophyllum nodosum* (*Fucales*, *Phaeophyta*) on the two spotted red spider mite *Tetranychus urticae*. *Hydrobiologia*. 1990;204:555–559.
22. Blunden G. Agricultural uses of seaweeds and seaweed extracts. In: Guiry MD, Blunden G (eds) *Seaweed resources in Europe: uses and potential*. Wiley, Chichester, 1991;65–81.
23. Mantri VA, Eswaran K, Shanmugam M, Ganesan M, Veeragurunathan V, Thiruppsthi S, Reddy CRK, Seth, A. An appraisal on commercial farming of *Kappaphycus alvarezii* in India: success in diversification of livelihood and prospects. *J Appl Phycol*. 2016;20:600–613.
24. Anderson G. Seaweed extract shows improved fruit quality at McLaren Vale vineyard trial. *Aust. N. Z. Grapegrow. Winemaker*. 2009;548;17–22.
25. De Góes HG, Reis RP. An initial comparison of tubular netting versus tie-tie methods of cultivation for *Kappaphycus alvarezii* (Rhodophyta, Solieriaceae) on the south coast of Rio de Janeiro State, Brazil. *J Appl Phycol*. 2012;23:607–613.
26. Ghosh A, Vijay Anand KG, Seth A (2015) Life cycle impact assessment of seaweed based bio-stimulant production from onshore cultivated *Kappaphycus alvarezii* (Doty) Doty ex Silva—is it environmentally sustainable? *Algal Res* 12:513–521.
27. Zhang X, Schmidt RE. The impact of growth regulators on the a-tocopherol status in water-stressed *Poa pratensis* L. *Int Turfgrass Res J*. 1997;8:1364–1373.
28. Crouch IJ, Smith MT, van Staden J, Lewis MJ, Hoad GV. Identification of auxins in a commercial seaweed concentrates. *J Plant Physiol*. 1992;139:590–594.
29. Crouch IJ, van Staden J. Evidence for the presence of plant growth regulators in commercial seaweed products. *Plant Growth Regul*. 1993;13:21–29.
30. Reitz SR, Trumble JT. Effects of cytokinin-containing seaweed extract on *Phaseolus lunatus* L.: influence of nutrient availability and apex removal. *Bot Mar*. 1996;39:33–38.
31. Durand N, Briand X, Meyer C. The effect of marine bioactive substances (NPRO) and exogenous cytokinins on nitrate reductase activity in *Arabidopsis thaliana*. *Physiol Plant*. 2003;119:489–49.
32. Stirk WA, Novak MS, van Staden J. Cytokinins in macroalgae. *Plant Growth Regul*. 2003;41:13–24.
33. Ordog V, Stirk WA, van Staden J, Novak O, Strnad M. Endogenous cytokinins in the three genera of microalgae from the Chlorophyta. *J Phycol*. 2004;40:88–95.
34. Crouch IJ, Van Staden J. Effect of seaweed concentrate from *Ecklonia maxima* (Osbeck) Papenfuss on *Meloidogyne incognita* infestation on tomato. *J Appl Phycol*. 1993;5:37–43.
35. Fornes F, Sañchez-Perales M, Guadiola JL. Effect of a seaweed extract on the productivity of 'de Nules' clementine mandarin and navelina orange. *Bot Mar*. 2002; 45:486–489.
36. Vernieri P, Borghesi E, Ferrante A, Magnani G. Application of biostimulants in floating system for improving rocket quality. *J Food Agric Environ*. 2005;3:86–88.
37. Wang Z, Pote J, Huang B. Responses of cytokinins, antioxidant enzymes, and lipid peroxidation in shoots of creeping bentgrass to high root-zone temperatures. *J Am Soc Hortic Sci*. 2003;128:648–655.

38. Flowers TJ, Yeo AR. Breeding for salinity resistance in crop plants—where next? *Aust J Plant Physiol.*1995;22:875–884.
39. Mittler R. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.* 2002;7:405–410.
40. Arora A, Sairam RK, Srivastava GC. Oxidative stress and antioxidative systems in plants. *Curr Sci.* 2002;82:1227–1238.
41. Mancuso S, Azzarello E, Mugnai S, Briand X. Marine bioactive substances (IPA extract) improve foliar ion uptake and water stress tolerance in potted *Vitis vinifera* plants. *Adv. Hortic. Sci.* 2006;20:156–161.
42. Wilson S. 2001. Frost Management in Cool Climate Vineyards. Final Report to Grape and Wine Research & Development Corporation.
43. Spinelli F, Fiori G, Noferini M, Sprocatti M, Costa GA. A novel type of seaweed extract as a natural alternative to the use of iron chelates in strawberry production. *Sci. Hortic.* 2010;125:263–269.
44. Ross R, Holden D. Commercial extracts of the brown seaweed *Ascophyllum nodosum* enhance growth and yield of strawberries. *HortScience.* 2010;45:141-148.
45. Blunden G, Jenkins T, Liu Y. Enhanced leaf chlorophyll levels in plants treated with seaweed extract. *J. Appl. Phycol.* 1997;8:535–543.
46. Fan D, Hodges, DM, Critchley AT, Prithiviraj, B. A commercial extract of brown macroalga (*Ascophyllum nodosum*) affects yield and the nutritional quality of spinach in vitro. *Commun. Soil Sci. Plant Anal.* 2013;44:1873–1884.
47. Jannin L, Arkoun M, Etienne P, Laîné P, Goux D, Garnica M, Fuentes M, Francisco SS, Baigorri R, Cruz F. Brassica napus growth is promoted by *Ascophyllum nodosum* (L.) Le Jol. seaweed extract: microarray analysis and physiological characterization of N, C, and S metabolisms. *J. Plant Growth Regul.* 2013;32:31–52.
48. Sivasankari S, Venkatesalu V, Anantharaj M, Chandrasekaran M, Effect of seaweed extracts on the growth and biochemical constituents of *Vigna sinensis*. *Biores. Technol.* 2006;97:1745–1751.
49. Spinelli F, Fiori G, Noferini M, Sprocatti M, Costa G. Perspectives on the use of a seaweed extract to moderate the negative effects of alternate bearing in apple trees. *J. Hortic. Sci. Biotechnol.* 2009;84:131–137.
50. Battacharyya D, Babgohari MH, Rathor P. Seaweed extracts as biostimulants in horticulture. *Sci Hortic.* 2015;96:39–48.
51. Kuwada K, Ishii T, Matsushita I, Matsumoto I, Kadoya K. Effect of seaweed extracts on hyphal growth of vesicular-arbuscular mycorrhizal fungi and their infectivity on trifoliolate orange roots. *J Jpn Soc Hortic Sci.* 1999;68:321–326.
52. Abetz P, Young CL. The effect of seaweed extract sprays derived from *Ascophyllum nodosum* on lettuce and cauliflower crops. *Bot Mar.* 1983;26:487–492.
53. Featonby-Smith BC, van Staden J. Effects of seaweed concentrate on grain yield in barley. *S Afr J Bot.* 1987;53:125–128.
54. Arthur GD, Stirk WA, van Staden J. Effect of a seaweed concentrate on the growth and yield of three varieties of *Capsicum annum*. *S Afr J Bot.* 2003;69:207–211.
55. Roshdy, K. A. Effect of spraying silicon and seaweed extract on growth and fruiting of Grand Naine banana. *Egypt J Agric Res.* 2014;92:979–991.

56. Ahmed FF, Mansour AE, Montasser MAA, Merwad MA, Mostafa EAM. Response of Valencia orange trees to foliar application of roselle, turmeric and seaweed extracts. *J Appl Sci Res.* 2013;9:960–964.
57. Ahmed FF, Ahmed MM, Ahmed AF. Partial replacement of inorganic nitrogen fertilizer by spraying some vitamins, yeast and seaweed extract in Ewaise mango orchard under upper Egypt conditions. *Stem Cell*, 2013;4:1–13.
58. Featonby-Smith BC, Van Staden J. The effect of seaweed concentrate on the growth of tomato plants in nematode-infested soil. *Sci Hortic.*1983;20:137–146.
59. Featonby-Smith BC, Van Staden J. The effect of seaweed concentrate and fertilizer on the growth of *Beta vulgaris*. *Z Pflanzenphysiol* 1983;112:155–162.
60. Featonby-Smith BC, van Staden J. The effect of seaweed concentrate and fertilizer on growth and the endogenous cytokinin content of *Phaseolus vulgaris*. *S Afr J Bot.* 1984 3:375–379.
61. Hutton MJ, van Staden J. Transport and metabolism of labeled zeatin applied to the stems of *Phaseolus vulgaris* at different stages of development. *Z Pflanzenphysiol.* 1984;114:331–339.
62. Adams-Phillips L, Barry C, Giovannoni J. Signal transduction systems regulating fruit ripening. *Trends Plant Sci.* 2004;9:331–338.
63. Varga A, Bruinsma J. The growth and ripening of tomato fruits at different levels of endogenous cytokinins. *J Hortic Sci.* 1974;49:135–142.
64. Norrie J, Keathley JP. Benefits of *Ascophyllum nodosum* marine-plant extract applications to ‘Thompson seedless’ grape production. (Proceedings of the Xth International Symposium on Plant Bioregulators in Fruit Production, 2005). *Acta Hortic.* 2006;727:243–247.
65. Chouliaras V, Tasioula M, Chatzissavvidis C, Therios I, Tsabolidou E. The effects of a seaweed extract in addition to nitrogen and boron fertilization on productivity, fruit maturation, leaf nutritional status and oil quality of the olive (*Olea europaea* L.) cultivar Koroneiki. *J. Sci. Food Agric.* 2009;89:984–988.
66. Blunden G, Jones EM, Passam JC. Effects of postharvest treatment of fruit and vegetables with cytokinin-active seaweed extract and kinetin solutions. *Bot. Mar.* 1978;21:237–240.
67. Kamel HM. Impact of garlic oil, seaweed extract and imazalil on keeping quality of valencia orange fruits during cold storage. *J. Hortic. Sci. Ornam. Plants.* 2014;6:116–125.
68. Fornes F, Sanchez-Perales M, Guardiola JL. Effect of a seaweed extract on the productivity of ‘de Nules’ Clementine mandarin and Navelina orange. *Bot. Mar.* 2002;45: 486–489.
69. Rayorath P, Narayanan JM, Farid A, Khan W, Palanisamy R, Hankins S, Critchley AT, Prithiviraj B. Rapid bioassays to evaluate the plant growth promoting activity of *Ascophyllum nodosum* (L.) Le Jol. using a model plant, *Arabidopsis thaliana* (L.) Heynh. *J Appl Phycol.* 2008;20:423–429.
70. IPCC (Intergovernmental Panel on Climate Change). Summary for policymakers. In: Parry ML, Canziani OF, Palutikot JP, van der Linden PJ, Hanson CE (eds) *Climate change 2007: impacts, adaptation and vulnerability. Contribution of Working Group II to*

the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.  
Cambridge University Press, Cambridge, UK; 2007.