

Screening and storage analysis of cider of bael (*Aegle marmelos* (L.) Correa) fruits

Abstract:- Bael is an underutilized, indigenous fruit with excellent nutritional content. During the summer, it is widely consumed as sharbat, and many of the beverages and goods derived from it are in great demand among customers. Cider is one of the fermented beverages made from bael that is in high demand on the worldwide market owing to its exceptional scent, flavour, and nutritional content. The cider market has enormous potential. Consequently, an experiment was conducted in the Post-Harvest Laboratory, Department of PHT, College of Horticulture and Forestry, Acharya Narendra Deva University of Agriculture and Technology Kumarganj Ayodhya (U.P) in the year 2020-21 in order to determine the stage and genotypes of bael for ideal cider, as well as the changes that occurred during the storage period of ideal cider. For screening bael cider in terms of organoleptic score and alcohol %, the study was conducted using a completely randomized design with three replications. After one month, another experiment will be conducted to assess the physicochemical changes that have occurred during storage. The results indicated that the cider made from NB-5 in May was superior than cider made from other stages and genotypes. Alcohol, hedonic (organoleptic) score, and titrable acidity exhibited a rising trend, whereas TSS, ascorbic acid, reducing sugars, non-reducing sugars, and total sugars exhibited a declining trend

.Key words – Bael, cider, physico-chemical, genotype, completely randomized design, alcohol, organoleptic score, fermentation

Introduction

The Bael fruit, or *Aegle marmelos* (L.) Correa, has been one of the most important fruits in India since the prehistoric time. It belongs to the Rutaceae family, aurantiodeae subfamily, and hard-shelled citroid fruit tree group and is thought to have its origins in the eastern ghats and central India (Rahman & Praven, 2014). The bael grown in tropical and subtropical regions. (Lambole *et al.* 2010) Bael juice and pulp make great summertime refreshers. The fruit is widely used in India as a liver and cardiac tonic and, when unripe or partially ripe, as an astringent, digestive, stomachic, useful in the treatment of diarrhoea and dysentery, as well as an efficient remedy for hi-cough, sore throat, and gum disease. The bael fruit possesses

laxative, cooling, and laxative effects as well as the capacity to treat constipation. Aegle marmelos has been utilised in Ayurvedic, Unani, and Siddha medicine as a herbal treatment for diabetes mellitus. High nutritional value can be found in marmelos' fruit. The pulp contains lignin, oil, inulin, water, sugar, protein, fibre, fat, calcium, phosphorus, potassium, iron, and vitamins (A, B, C, and riboflavin), as well as alkaloids, cardiac glycosides, terpenoids, flavonoids, and flavonoids. Fruit has a 61.5% water content, 31.8% carbohydrates, 1.8% protein, 1.75% minerals, 55mg carotene, 0.39mg fat, 0.13mg thiamine, 1.19mg riboflavin (the greatest amount among fruit crops), 1.1mg niacin, and 8mg vitamin C per 100g of edible portion. (Gopalan et al, 1971)

Bael is not widely used as a table fruit due of its hard shell, mucilaginous texture, and abundant seeds, which make it difficult to eat by hand. A processing opportunity exists in the excellent flavour, nutritional content, and therapeutic benefits of bael fruit (Roy and Singh,1979a). Refined beverages like Bael Cider, RTS, squash, etc. can be made from ripe bael fruits. The best alcoholic sweet beverage is Bael Cider. The diluted bael pulp was improved with sugar and acid and fermented at $30\pm 2^{\circ}$ C using the yeast *Saccharomyces cerevisiae*. (Garg et al. 2014). Patra et al. (2015) reported fermented bael juice contains titrable acidity and lactic acid with 1.7mg tartaric acid/L and 7.2mg/L, respectively. It is rich in antioxidants and can be utilised as a medicinal drink with significant nutritional potential. The Bael cider is healthier than any other fermented beverages. have PH (4.1), total phenolics (0.93g/100 ml), beta carotene (33g/100 ml), ascorbic acid (80 mg/100 ml), lactic acid (0.64 mg/100 ml), TSS (12.90B), total sugar (2.03g/100 ml), titrable acidity (0.15g tartaric acid/100 ml), and ethanol (7.87%). (Panda et al. 2013). This drink would be healthier than any other artificially-flavoured alcoholic beverages, which are widely available in our nation.

Materials and Methodology

The present investigation was carried out at Post-Harvest Laboratory, Department of PHT, College of Horticulture and Forestry, Acharya Narendra Deva University of Agriculture and Technology Kumarganj Ayodhya (U.P) in the year 2020-21 the healthy fruits free from disease, pest and bruises were randomly selected from the main experiment station, Horticulture, Acharya Narendra Deva University of Agriculture and Technology Kumarganj Ayodhya (U.P) India

Techniques for Preparation of bael cider:

Preparation of starter:

Bael fruits are picked and sliced. After retaining T.S.S. (20 percent), these bael slices were combined with water in a ratio of 1:1, brought to a boil, and chilled to room temperature. After cooling, *Saccharomyces cerevisiae* yeast culture was inoculated under laminar airflow, and aluminum foil was put on top and secured with a rubber band so that no pathogen could infect the culture. To create the starting culture, the whole combination was incubated for 24 hours at 30° C.

Methodology:

Firm ripe fruits free from any disease were selected, thoroughly washed and broken the shell, removed pulp with seed and fiber. The bael pulp was mixed with water in a ratio of one part pulp and one part water and heated up to 80⁰C. After cooling it was massed with hands and the acidity (0.5%) and T.S.S (20%) were adjusted, then transferred to the container (at least 1/4th part of the container should be vacant) thereafter Sulphur dioxide 100 ppm was added in form of potassium metabisulphite. After half an hour, it was inoculated with the starter of *Saccharomyces cerevisiae var. ellipsoids* yeast (4%) close the container airtight and the airlock was adjusted on the top of the container for the exhaustion of carbon-dioxide gas from the container to the outside and restrict air from outside to inside the container. At the peak level of fermentation pure pectic enzyme was added (0.5%) for getting an easy clearing. After that the must was filtered through the two layered muslin cloth discarding seeds and pomace. Filtered juice was kept again in the container for fermentation. Two- three weeks were required for primary fermentation phase.

After the completion of primary fermentation (when the bubble does not come out in the airlock), the clear fermented juice was siphoned off and filled into the containers up to the brim and 50ppm Sulphur dioxide in form of potassium metabisulphite was added, sealed, airtight and kept at room temperature for maturation. In order to carry out racking for a period of six months (at an interval of two months) storage

After the maturation of cider fermented juice filled in 200ml glass bottles (upto drum), crown corked, pasteurized at 65⁰C for 20 minutes and cooled in air and stored at ambient temperature. Ideal Cider was stored at ambient temperature for the investigation of storage stability.

Chemical estimation of bael cider

TSS was estimated by using a hand refractometer (Erma) of 0 to 32% and 28-62 % range at room temperature and the reading was corrected at 20⁰ C temperature. Sugars was estimated by using Fehling solution 'A' and 'B' as per method given by Lane and Eynon (1923) and Ranganna , 2010 The acidity and ascorbic acid was estimated as per method described by (A.O.A.C., 1970), the alcohol estimation was done by experts of RFRAC Lucknow through the procedure mention in

Statistical analysis

Statistical analyses of the data obtained in the experiments were calculated as suggested by Panse and Sukhatme (1985) and results were evaluated at 5% level of significance.

Result and Discussion

Screening of suitable genotype and stage of fruit for bael cider

The data arranged in table 1 indicate that cider produced in the month of May was noticeably superior to earlier stages. Cider made from the NB-5 genotype had a much higher organoleptic score than cider prepared from other genotypes of bael. The interaction impact between genotype and stage was found non-significant. Cider made from the NB-5 in May is of superior quality, followed by cider made from the NB-9 in May.

The alcohol produced in May has the greatest quantitative alcohol content compared to the other phases. The bael cider made from NB-5 was found to have the highest alcohol content among the genotypes. Bael's genotypes and the interaction effect were determined to be statistically non-significant. The bael cider made with NB-5 in May has the greatest alcohol content, followed by bael cider made with NB-9 in May.

Chemical changes during the storage analysis of bael cider

The data arranged in the Table 2 clearly indicate the consistent increasing trend of the organoleptic quality of bael cider during the storage period was recorded. The organoleptic score increasing from 7.68 at the time of cider preparation to 8.46 at the 7th month of observation. The increase in organoleptic quality may be due to increase in the percentage of alcohol through secondary fermentation and reduction in sulphur dioxide and yeast odour. The similar findings were also reported by Singh *et al.*, 2006. Apple cider's organoleptic score increased with storage Tripathi *et al.*, 1989

The alcohol content of bael cider and storage duration have a positive correlation. The trend of increasing alcohol content along with the elongation of bael cider storage. The alcohol

content increased from 7.19% at the time of cider production to 7.22% during the seventh month of monitoring. The rise in alcohol (percent) may be attributed to polysaccharide hydrolysis to fermentable sugar, secondary fermentation of fermentable sugar, and conversion to ethyl alcohol during bael cider storage. Ram (1996) found that bael cider's alcohol concentration increased for five months then stabilised. Similar results were also reported by Singh *et al.* (1976)

Total soluble solids have been steadily decreasing while the bael cider has been stored. Up to the third month of storage, the decreasing tendency was more pronounced than the subsequent shifting pattern. At the time of preparation, the maximum amount of cider that could be stored was noted. It then steadily decreased to a level of 6.81 °Brix at the end of the seventh observational month. The decrease in TSS may be due to hydrolysis of polysaccharide to simple sugars. Kiribhaga *et al.* (2020) also reported similar downward trend during storage of banana wine

Acidity percentage of the bael cider gradually decrease with the storage period afterwards. The acidity percentage changes from 0.764 % at the time of preparation to 0.391% at the end of the storage period. The decrease in acidity content might be due to breakdown of citric acid in the pulp to mobilize the sugar. The results were in close agreement with Jespersen *et al.* (2005); Nielsen *et al.* (2007) in cocoa beans

Ascorbic acid levels in bael cider have been declining steadily over time. When bael cider was being prepared, ascorbic acid levels were found to be 82.36 mg/100 g of pulp, but by the seventh month of storage, it had dropped to 35.23 mg/100 g of pulp. Oxidation of ascorbic acid with trapped oxygen in the bottling which ultimately leads to synthesis of dehydroascorbic acid. Kiribhaga *et al.* (2020) also reported similar downward trend during storage of banana wine.

the tendency of reducing sugars to decrease as bael cider was stored. The drop started off gradually during the first two months of storage, but it then became more steady up to the fifth month. After that, in the fifth month of storage, the reducing sugars almost exactly remained constant. The non-reducing sugar in bael cider was also reduced during storage. For the first two months of storage, the drop was moderate. After that, it continued to decline gradually until the fifth month of storage, at which point the non-reducing sugar levels were nearly constant. The negative relationship between total sugars and bael cider storage duration was observed. Total sugars were shown to trend downward similarly to reducing and non-reducing sugars. The downward trend in the sugars might be due to utilization of pectic

substances by the yeast during fermentation in presence of pectic enzyme added during must preparation. Similar findings were also reported by Aradhna and Fleet (2003); Garg (2014)

Conclusion

Based on the results, it can be concluded that Bael cider prepared from NB-5 in the month of May has highest alcohol (%) and hedonic score. The alcohol (%) and acceptability of cider were increasing whereas the TSS, Sugars and ascorbic acid, acidity decreased during the storage significantly improving the quality of bael cider during the storage

References

- Aradhna, M. M. and Fleet, G. H. (2003) The microbial ecology of cocoa bean fermentations in Indonesia. *Int. J. of Food Microbiology* **86**: 87-99
- Association of Official Analytical Chemists (AOAC)(1970.), "Official Methods of Analysis," *10th Edition, Washington D.C.*, pp. 154-170.
- Garg, N.; Kumar, S.; Yadav, K. and Yadav, P. (2014) Development of vitamin C fortified Bael Cider. *Lucknow Science Congress, LUSCON-2014* on March27,2014 organized by BBAU, Lucknow, Uttar Pradesh, India.
- Gopalan, C.B.V.; Shastri, R. and Subramanian, B. (1971) Nutritive value of Indian food, National institute of Nutrition, *ICMR*, Hyderabad, India.
- Jesperersen, L.; Nielsen, D.S. Honholt, S. and Jakobsen, M. (2005). Occurrence and diversity of yeasts involved in fermentation of west African cocoa beans. *FEMS yeast Research* **5**: 441-453

- Kiribhaga, Sarthak & Gomez, Saji & Joseph, Meagle & K., Surendra & Thomachan, Seeja. (2020). Biochemical changes of banana wine during storage *International Journal of Chemical Studies*; **8**(1): 119-124
- Lambole, V. B., Krishna M., Upendra K, Bhatt SKP, Vipul, G. (2010) Phytopharmacological properties of *Aegle marmelos* as a potential medicinal tree: an overview. *Int. J. Pharm. Sci. Rev. Res.* 5(2): 67-72
- Lane, J.H. and Eynon, L. (1923) Methods for Determination of Reducing and Non-Reducing Sugars. *Journal of Sciences*, 42, 32-37.
- Nielsen, D.S.; Teniola, O. D.; Ban, K. L.; Owusu, M.; Andersson, T. S. and Holzapfel, W. H. (2007) The microbiology of Ghanaian cocoa fermentations analyzed using culture-dependent and culture independent methods. *Int. J. of Food Microbiology*. **114**: 168-186
- Panda, S.K.; Sahu, U.C.; Roy, R.C. (2013) Bio-processing of Bael (*Aegle marmelos* (L.) Correa.) fruits into wine with antioxidants. *Food Bio-Science*. 5(10).10-16
- Panse, V.G. and Sukhatme, P.V. (1985) Statistical Methods for Agricultural Workers. *Indian Council of Agricultural Research Publication*, 87-89..
- Patra, J. K.; Sahoo, S.K.; Swain, M.R. (2015) Nutritional and antioxidant potential of *Aegle marmelos* fermented fruit juice. *Proc. Natl. Acad. Sci., India, Sect. B Biol. Sci.* DOI 10.1007/s40011-015-0644-4
- Rahman S., Parvin R (2014) Therapeutic potential of *aegle marmelos* (L.). An Overview. *Asian. Pac. J. Trop. Dis.* 4(1):71-77
- Ram. D. (1996) Studies on ripening storage and processing of bael (*Aegle marmelos* Correa) fruit. Ph.D. Thesis. *NDUAT Faizabad* pp-79
- Rangana, S. 2010. Analysis and quality control for fruit and vegetable products. *Tata McGraw Hill Ltd. New Delhi*.
- Singh Nagi, H. P. P. and Manjrekar, S. P. (1976). Studies in the preparation of cider from North Indian apple. II. Storage studies. *India food Packer*. **30** (1): 12
- Singh, S.; Godara, R. K.; Saini, R.S.; Patahk, D. V. and Sharma, J. R. (2006) Effect of pH on the quality of bael wine fermented under controlled condition. *Haryana J. Hort. Sci.*, **35**(3&4): 249-250
- Tripathi, U. K.; Ram, B. H.; Jain, S.P. and Singh, S. (1989). Changes in developing banana fruit. *Prog. Hort.* **13**(1):45-51

Table -1 Screening of suitable genotype and stage of fruit for bael cider

Treatment	Organoleptic score	Alcohol (%)
March (S ₁)	6.58	6.84
April (S ₂)	7.17	7.03
May (S ₃)	7.42	7.15
SEm±	0.067	0.043
C.D.(P=0.05)	0.197	0.127
NB-4 (V ₁)	6.83	6.98
NB-5 (V ₂)	7.3	7.12
NB-7 (V ₃)	6.97	6.89
NB-9 (V ₄)	7.12	7.04
SEm±	0.077	0.05
C.D.(P=0.05)	0.227	0.147
S₁V₁	6.32	6.78
S₁V₂	6.88	7.01
S₁V₃	6.46	6.64
S₁V₄	6.64	6.92
S₂V₁	6.94	7.02
S₂V₂	7.35	7.16
S₂V₃	7.12	6.91
S₂V₄	7.26	7.03
S₃V₁	7.22	7.14
S₃V₂	7.68	7.2
S₃V₃	7.34	7.11
S₃V₄	7.45	7.16
SEm±	0.134	0.087
C.D.(P=0.05)	NS	NS

Fig 1.1.: Organoleptic score of the bael cider

Fig 1.2. Alcohol (%) of the bael cider

Table-2 Chemical changes during the storage analysis of bael cider

Storage period	Organoleptic score	Alcohol content (%)	TSS⁰ (°Brix)	Acidity (%)	Ascorbic acid (mg/100g pulp)	Reducing sugars (%)	Non-reducing sugar (%)	Total sugars (%)
0	7.68	7.19	11.13	0.764	82.36	4.42	5.62	10.04
1	7.87	7.20	10.04	0.715	78.6	4.16	5.34	9.5
2	7.93	7.22	8.86	0.643	71.52	3.52	4.31	7.83
3	8.06	7.23	7.98	0.61	62.13	2.48	4.18	6.66
4	8.24	7.24	7.52	0.552	54.29	2.31	3.97	6.28
5	8.31	7.28	7.15	0.501	46.35	2.11	3.81	5.92
6	8.39	7.28	6.97	0.461	41.28	1.97	3.68	5.65
7	8.46	7.31	6.81	0.391	35.23	1.94	3.64	5.58
SEm±	0.11	0.03	0.14	0.009	0.88	0.03	0.05	0.11
C. D. at 5%	0.32	0.08	0.43	0.025	2.65	0.11	0.14	0.34

Fig 2.1: Changes in organoleptic score during storage of bael cider

Fig 2.2: Changes in alcohol (%) during storage of bael cider

Fig 2.3: Changes in TSS during storage of bael cider

Fig 2.4: Changes in acidity (%) during storage of bael cider

Fig 2.5: Changes in ascorbic acid (mg/100g pulp) during storage of bael cider

Fig 2.6: Changes in sugars (%) during storage of bael cider