

**Effect of Foliar Spray of Potassium and Micronutrients on Yield and Quality of Papaya (*Carica papaya* L.)  
cv. Red Lady**

**ABSTRACT**

The study was carried out in Farmer's field at Basur village, Sorab taluk, Shivamogga dist, during the year 2018-2019. The experiment was laid out in randomized complete block design with three replication and eleven treatments. Results of this study reveal that the combined application of T<sub>7</sub> - K<sub>2</sub>SO<sub>4</sub> at 1% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25% recorded significantly maximum yield and yield attributing parameters *i.e.* fruit length (27.20cm), fruit diameter (16.17cm), fruit weight (2438g and 2.43kg), number of fruits per plant (49.67), yield plant<sup>-1</sup> (90.07kg) and yield (224.96 t ha<sup>-1</sup>), pulp weight (2343.74g), peel weight (54.98g), seed weight (52.70g) and quality characters like total soluble solids (13.52 °brix), ascorbic acid (54.42mg 100g<sup>-1</sup>), total sugar (5.64%), reducing sugar (4.60%) and minimum acidity (0.13%) over control and which was on par with T<sub>8</sub> (K<sub>2</sub>SO<sub>4</sub> at 2% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25%) and non-reducing sugar was found to be statistically non-significant among the different treatments. The results revealed that the plants treated with T<sub>7</sub> - K<sub>2</sub>SO<sub>4</sub> at 1% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25% were found to be effective to improve the yield and quality of papaya cv. Red Lady.

Key words: Papaya, Foliar spray, Potash and Micro nutrients.

**Introduction**

Papaya (*Carica papaya* L.) is an important fruit of tropical and subtropical regions of the world, belonging to the family Caricaceae and also known as “wonder fruit of the tropics” (Parmaret *al.*, 2017). It is native of Tropical America was introduced to India in the

16th century from Malacca (Singh, 1990). India is the largest producer of papaya in the world producing 5.83 million MT of fruits from an area of 0.13 mha with productivity of 42.3 MT/ha (NHB, 2019). Karnataka is the third largest producer of papaya in the country. It is commercially cultivated in Andhra Pradesh, Gujarat, Karnataka, West Bengal, Madhya Pradesh, Maharashtra and Tamil Nadu. Papaya fruits are also good source of many minerals (potassium, phosphorus and magnesium) in human diet. The papaya fruit belongs to the group of low acid content fruit and the pH of pulp ranges from 5.5 to 5.9 (Chan *et al.*, 1971).

About 40-55% of Indian soils are moderately deficient in Zinc and 25-30% is deficient in Boron. Deficiency of other micronutrients occurs under 15% of soils. Deficiency of micronutrient has become a major constraint to the productivity, stability and sustainability of crops in many Indian soils and may further deteriorate due to global warming (Kumar *et al.*, 2011).

Adequate K nutrition has associated with increased yields, fruit size, increased soluble solids and ascorbic acid concentrations, improved fruit colour, increased shelf life, and shipping quality of many horticultural crops (Geraldson, 1985; Ganeshamurthy *et al.*, 2011). Transportation of photo assimilates from source tissues via the phloem to sink tissues, enzyme activation, turgor maintenance, and stress tolerance (Marschner, 1995).

Micronutrients can tremendously increase crop yield and get better quality and post-harvest life of produce. They play a significant role in disease resistance, since they function as enzyme activators and also play a function in lignin biosynthesis (Parmar *et al.*, 2017).

Foliar application of micronutrients has gained importance because foliar application is a well-established operation to complete and enrich plant nutrition. Foliar application can meet the expense of nutrients where absorption of nutrients from the soil is unavailable due to plant stress or in adverse soil conditions. Foliar sprays with fertilizers including macro element like potassium and microelements such as Zn and B have been shown to be convenient for field use, have a good effectiveness and very rapid plant

response (Fernandez *et al.*, 2013).Hence, foliar application of potassium, zinc and boron in specific plant developmental and critical stages improves the yield and quality of papaya.

## Materials and Methods

An investigation was carried out in Farmer's field at Basur village, Sorab taluk, Shivamogga dist with the support of Department of Fruit science at College of Horticulture, Bengaluru, University of Horticultural Sciences, Bagalkot during 2018-2019. The experiment was laid out in randomized block design with three replications and eleven treatments viz., T<sub>1</sub> - K<sub>2</sub>SO<sub>4</sub>at 1%, T<sub>2</sub> - K<sub>2</sub>SO<sub>4</sub>at 2%, T<sub>3</sub> - Borax at 0.25%, T<sub>4</sub> - Borax at 0.50%, T<sub>5</sub> - ZnSO<sub>4</sub>at 0.25%, T<sub>6</sub> - ZnSO<sub>4</sub>at 0.50%, T<sub>7</sub> - K<sub>2</sub>SO<sub>4</sub>at 1% + Borax at 0.25% + ZnSO<sub>4</sub>at 0.25%, T<sub>8</sub> - K<sub>2</sub>SO<sub>4</sub>at 2% + Borax at 0.25% + ZnSO<sub>4</sub>at 0.25%, T<sub>9</sub> - K<sub>2</sub>SO<sub>4</sub>at 1% + Borax at 0.50% + ZnSO<sub>4</sub>at 0.50%, T<sub>10</sub> - K<sub>2</sub>SO<sub>4</sub>at 2% + Borax at 0.50% + ZnSO<sub>4</sub>at 0.50%, T<sub>11</sub> - Control (water spray) with a spacing 2 x 2 m. Foliar application of potassium, zinc and boron were applied in six split doses with 45 days interval after transplanting. Three plants per replication were randomly selected and tagged in each treatment and in each replication. The fruits were harvested based on their maturity indices viz., change in colour of fruit from dark green to yellowish orange. Observations were recorded and statistically analyzed as per the methods given by Panse and Sukhatme(1967).

## Results and Discussion

In the present investigation, the data obtained on yield and yield attributing parameters *i.e.* fruit length (cm), fruit diameter (cm), fruit weight (g), number of fruits per plant, fruit yield plant per plant (kg), yield per ha (ton), pulp weight (g), peel weight (g), seed weight (g)and quality characters like total soluble solids, ascorbic acid, total sugar, reducing sugar and acidity were significantly influenced by spraying of different concentrations of potassium, zinc and boron are recorded and presented in Table1 and Table 2. The data obtained on non-reducing sugar was found to be statistically non-significant among the different treatments.

Among different treatments, the results observed that the fruit length, fruit diameter and fruit weight showed significantly maximum (27.20cm, 16.17cm and 2438.00g, respectively) in treatment T<sub>7</sub> (K<sub>2</sub>SO<sub>4</sub> at 1% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25%), which were on par with application of K<sub>2</sub>SO<sub>4</sub> at 2% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25% (26.97cm,15.84cm and 2397.64g, respectively) and were followed by ZnSO<sub>4</sub> at 0.25% (24.38cm,14.53cm and 2148.32g, respectively), Borax at 0.25% (24.05cm,14.29cm and 2113.55g, respectively). Whereas, minimum fruit length,fruit diameter and fruit weight (20.07cm, 11.98cm and 1802.43g, respectively) were observed in T<sub>11</sub> (control-water spray).This might be due to combined effect of zinc and boroninvolved in hormonal metabolism, starch formation, cell division, cell expansion and increased volume of intercellular spaces in the mesocarpic cells and higher mobilization of photosynthates from other parts of the plant towards the developing fruits that are extremely active metabolic sink (Singh *et al.*, 2001) and (El-Rhman and Shadia, 2012). The similar findings were reported by Singh *et al.* (2009), Waskela *et al.* (2013) and Bhalerao *et al.* (2014). Zinc helps in regulating the semi permeability of cell walls, thus mobilizing more water into fruits increased rate of cell division and cell enlargement leading to more accumulation of metabolites in the fruit resulting increase in fruit length, fruit diameter and fruit weight (Babu and Singh, 2001). Similar results were obtained by Chaitanya (1997), (Wali *et al.*, 2005), Singh *et al.* (2010), Yadav *et al.* (2010) and Rajkumar *et al.* (2014). Boron activates rapid mobilization of water and sugar in the fruit which intern increased in accumulation of dry matter within the fruit (Bhatt *etal.* 2012) and application of boron might have caused rapid synthesis of protein and translocation of carbohydrate which ultimately led to increase in fruit weight (Singh *et al.*, 2012).

The results obtained for number of fruits per plant, fruit yield plant per plant (kg) and yield per ha(ton)were significantly maximum (49.67, 90.07 and 224.96, respectively) with foliar application of K<sub>2</sub>SO<sub>4</sub> at 1% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25%, which were on par with the application of K<sub>2</sub>SO<sub>4</sub> at 2% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25% (47.22, 88.37 and 221.67, respectively) and were followed by ZnSO<sub>4</sub> at 0.25% (42.12, 80.79 and 201.95, respectively), Borax at 0.25% (42.07, 80.48 and 201.68, respectively). Whereas, T<sub>7</sub>, T<sub>8</sub>, T<sub>5</sub> and T<sub>3</sub> was registered significantly superior over all other treatments respectively. Whereas, in T<sub>11</sub> (control-water spray) was noticed minimum number of fruits per plant, fruit yield plant per plant (kg) and yield per ha(ton) (33.52,71.85 and 178.64, respectively). This might be due to combined effect offoliar application of potassium, zinc and borondirectly or indirectly, involved in

fruit setting, retention and their activity improved number of fruits per plant associated with photosynthesis, hormone metabolism which promotes synthesis of auxin, necessary for fruit set and fruit growth (Rajkumar *et al.* 2014). Zinc is helpful in chlorophyll synthesis which increases photosynthetic activities of leaves, which leading to development of primary flowers, production of viable flowers with improve pollination and fruit setting (Jat and Kacha, 2014). Similar results were also obtained by (Ali, 1991), (Ghanta, 1992), Babu and Yadav (2005) and (Kundu *et al.*, 1989; Singh *et al.*, 2010; Yadav and Singh, 2010; Modi *et al.*, 2012; Abhijith *et al.*, 2018). The application of boron might have caused rapid synthesis of protein and translocation of carbohydrate which ultimately led to increase in fruit weight which is directly correlated with total yield (Parr and Loughman, 1983) and (Singh *et al.*, 2012). Similar results have been reported by (Panwar *et al.*, 1995) and (Babu and Singh, 2001) and (Allah, 2006) who reported that increase in number of fruits probably due to influence of boron which increases pollen grain germination and pollen tube elongation, consequently leads to higher fruit set and finally more number of fruits per plant. The number of fruits produced per plant has significantly reflected on high yield. Higher the fruit number more will be the yield. Similar results were obtained by Kudada and Prasad (2002), Singh *et al.* (2007), Singh *et al.* (2010), Sajid *et al.* (2010) and Trivedi *et al.* (2012).

Significantly the maximum pulp weight, peel weight and seed weight (2343.74g, 54.98g and 52.70g, respectively) was obtained with foliar application T<sub>7</sub> (K<sub>2</sub>SO<sub>4</sub> at 1% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25%), which was on par (2268.64g, 52.63g and 50.65g, respectively) with T<sub>8</sub> (K<sub>2</sub>SO<sub>4</sub> at 2% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25%) and were followed by (2043.56g, 48.07g and 46.18g, respectively) with T<sub>5</sub> (ZnSO<sub>4</sub> at 0.25%), (2015.27g, 47.63g and 45.95g, respectively) with T<sub>3</sub> (Borax at 0.25%). Whereas, T<sub>7</sub>, T<sub>8</sub>, T<sub>5</sub> and T<sub>3</sub> was registered significantly superior over all other treatments respectively. The treatment T<sub>11</sub> (control-water spray) were observed minimum pulp weight, peel weight and seed weight (1692.28g, 41.34g and 40.05g, respectively). This increase in fruit pulp weight might be due to minimum fruit cavity index, increased fruit length, diameter, fruit weight and more accumulation of photosynthates in the matured fruits by beneficial effect of boron and zinc. Similar results were reported by Singh *et al.* (2010).

The present study indicates that significantly maximum total soluble solids (13.52 °brix), ascorbic acid (54.42mg 100g<sup>-1</sup>), total sugar (5.64%), reducing sugar (4.60%) and minimum acidity (0.13%) were observed in T<sub>7</sub> (K<sub>2</sub>SO<sub>4</sub> at 1% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25%). Whereas, T<sub>11</sub> (control-water spray) has registered significantly minimum total soluble solids (12.52%), ascorbic acid (50.45mg 100 g<sup>-1</sup>), total sugars (4.84%), reducing sugar (3.82%) and maximum acidity (0.16). This might be due to influence of potassium which regulates photosynthesis and it is also essential for starch synthesis and sugar accumulation by activating enzymes viz., starch synthetase, etc., (Lester *et al.*, 2010). Potassium is excessively required during fruit maturation stage for enhancing the fruit size, colour and taste of fruits (Ganeshamurthy *et al.*, 2011). Cell elongation accompanied with increase in sugar content (Syamal *et al.*, 2010). Similar results were obtained by (Kaur and Dhillon, 2006; Singh *et al.*, 2009; Manju and Kumar, 2015). TSS is directly associated with reducing and non-reducing sugars, the TSS will significantly reflect on more conversion of starch into reducing and non-reducing sugars during ripening process and this also might be due to influenced by zinc and boron application by regulating translocation of photosynthates to fruit pulp and hydrolysis of complex polysaccharides into simple sugars reported by Kavitha (2000).

## Conclusion

Present experiment reveals that foliar application is an instant effective way of application of nutrients. Plants sprayed with K<sub>2</sub>SO<sub>4</sub> at 1% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25% significantly increased yield and quality of papaya cv. Red Lady.

## References

- Abhijith, Y. C., Dinakara, A. J., Kishor, H. and Sindhu, C., 2018, Effect of micronutrients on yield and quality of aonla (*Emblica officinalis* Gaertn.) cv. NA-7. *Int. J. Curr. Microbiol. App. Sci.*, 7(3): 140-145.
- Ali, W., Pathak, R. K. and Yadav, A. L., 1991, Effect of foliar application of nutrients on guava cv. Allahabad Safeda. *Prog. Hort.*, 23(1-4): 18-31.

- Allah, A. S. E., 2006, Effect of spraying some macro and micro nutrients on fruit set, yield and fruit quality of Washington navel trees. *J. App. Sci.*, 22(1): 54-56.
- Anonymous. 2018-19. State wise area, production and productivity of papaya. National Horticulture Board, Indian Horticulture Database-2018-19.
- Babu, K. D. and Yadav, D. S., 2005, Foliar spray of micronutrients for yield and quality improvement in Khasi mandarin (*Citrus reticulata* Blanco.). *Ind. J. Hort.*, 62(3): 280-281.
- Babu, N. and Singh, A. R., 2001, Effect of foliar application of boron, zinc and copper on chemical characterization of litchi fruits. *Bioved.*, 12 (1/2): 45-48.
- Bhalerao, P. P., Patel, B. N., Patil, S. J. and Gaikwad, S. S., 2014, Effect of foliar application of Ca, Zn, Fe and B on growth, yield and quality of papaya (*Carica papaya*) cv. Taiwan Red Lady. *Current Hort.*, 2(2): 35-39.
- Cakmak, I., 2008, Enrichment of cereal grain with zinc: agronomic or genetic biofertilization. *Pl. Soil.*, 302: 1-17.
- Chaitanya, C. G., Kumar, G., Raina, B. L. and Muthoo, A. K., 1997, Effect of foliar application of zinc and boron on yield and quality of guava. *Haryana J. Hort.Sci.*, 261: 78-80.
- Chan, Jr. H. T., Chang, T. S. K., Stafford, A. E., Brekke, J. E., 1971, Non volatile acids in papaya. *J. Agri. Food. Chem.*, 19(2): 263–265.
- El-Rhman, A. I. E. and Shadia, A. A., 2012, Effect of foliar sprays of urea and zinc on yield and physio-chemical composition on jujube (*Ziziphus mauritiana*). *M. E. J. of Agri. Res.*, 1(1): 52-57.
- Fernandez, V., Sotiropoulos, T., Brown, P. H., 2013, Foliar fertilization. In: Scientific Principles and Field Practices. International Fertilizer Industry Association, Paris.
- Ganeshamurthy, N., Satisha, G. C. and Patil, P., 2011, Potassium nutrition on yield and quality of fruit crops with special emphasis on banana and grapes. *Karnataka J. Agric. Sci.*, 24(1): 29-38.

- Geraldson, C. M., 1985, Potassium nutrition of vegetable crops. *In Munson, R.D. (Ed) Potassium in Agriculture*, Madison, W. I., 94(1): 915- 927.
- Ghanta, P. K., Dhus, R. S. and Mitra, S. K., 1992, Response of papaya to foliar spray of Boron, Manganese and Copper. *Hort. J.*, 5(1): 43-48.
- Jat, G. and Kacha, H. L., 2014, Response of guava to foliar application of urea and zinc on fruit set, yield and quality. *J. Agri. Search.*, 1(2): 86-91.
- Kaur, G., Dhillon, W. S., 2006, Effect of foliar application of chemicals on physicochemical characters of variety Allahabad Safeda of guava during winter. *Res. Punjab Agri. Univ.*, 43 (2): 114-116.
- Kavitha, M., Kumar, N. and Jeyakumar, P., 2000, Effect of zinc and boron on biochemical and quality characters of papaya cv. CO.5. *South Indian Horti.*, 48 (1/6): 1-5.
- Kudada, N. and Prasad, S. M., 2002, Effect of manuring on incidence of papaya ring spot virus and yield attributes of pot grown papaya cv. Rajdoot. *J. Res.*, 52(3): 224-227.
- Kumar, M., Swarup, A., Patra, A. K., Chandrakala, J. U., 2011, Micronutrient fertilization under rising atmospheric CO<sub>2</sub> for micronutrient security in India. *Ind. J. Ferti.*, 7(7): 52-60.
- Kundu, A., Mitra, S. K., Ghosh, S. K. and Bose, T. K., 1989, Effect of micronutrient on foliar application in papaya. *Mysore j. Agric. Sci.*, 23: 65-70.
- Lester, G. E., Jifon, J. L. and Makus, D. J., 2010, Impact of potassium nutrition on postharvest fruit quality: Melon (*Cucumis melo* L) case study. *Plant Soil*, 335:117–131.
- Manju, V. and Kumar, S., 2015, A dynamic response of potassium and micro nutrients combined with brassinosteroids - a steroidal plant hormone, on accumulation of sugars in papaya cv. TNAU Papaya CO-8. *Int. J. Agri. Sci. Res.*, 5(6):277-282.
- Marschner, H., 1995, Functions of mineral nutrients: macronutrients, In: H. Marschner (Ed.). *Mineral nutrition of higher plants* 2nd Edition. Academic Press, N. Y., 94(1): 299-312.

- Marschner, H., 2012, Mineral Nutrition of Higher Plants. Academic Press Limited Harcourt Brace and Company, Publishers, London, pp. 347-364.
- Modi, P. K., Varma, L. R., Bhalerao, P. P., Verma, P. and Khade, A., 2012, Micronutrient sprays effects on growth, yield and quality of papaya (*Caricapapaya* L.) cv. Madhu Bindu. *Madrass Agric. J.*, 99(7/9): 500-502.
- Panse, V. G. and Sukhatme, P. V., 1967, Statistical method of Agricultural workers, ICAR Publication. New Delhi. pp.381.
- Panwar, R. D., Saini, R. S., Kaushik, R. A. and Yamdagni, R., 1995, Effect of micronutrients on fruit retention, yield and quality of aonlacv. Banarasi under rainfed condition. *Haryana J. Hort. Sci.*, 22 (1-2): 250-251.
- Parmar, P., Patil, S. J., Kumar, S., Asha, C. M. and Tandel, B. M., 2017, Response of fertilizer application on growth of papaya var. Red Lady. *Int. J. Curr. Microbiol. App. Sci.*, 6(12): 2375-2379.
- Parr, A. J. and Laughman, B. C., 1983, Boron and membrane functions in plants. In: Metals and micronutrients: Uptake and utilization by plants. Robb, D. A. and Pirsiont, W. S. Eds. *Annu. Proc. Phytochem. Soc. Eur.*, 21:87.
- Rajkumar, Tiwari, J. P. and Shantlal, 2014, Effect of foliar application of zinc and boron on fruit yield and quality of winter season guava (*Psidium guajava*) cv. Pant Prabhat. *Annals of Agri Bio Res.*, 19(1): 105-108.
- Sajid, M., Abdur-Rab, Ali, N. and Arif, M., 2010, Effect of foliar application of Zn and B on fruit production and physiological disorders in sweet orange cv. Blood Orange. *Sarhad J. Agric.*, 26(3): 355-360.
- Singh, D. K., Ghosh, S. K., Paul, P. K. and Suresh, C. P., 2010, Effect of different micronutrients on growth, yield and quality of papaya (*Carica papaya* L.) cv. Ranchi. *Acta Hort.*, 851:351-356.
- Singh, D. M., Singh, H. K., Vishwanath and Pratap, B., 2009, Effect of foliar feeding of nutrients on yield and quality of aonla (*Emblica officinalis*Gaertn.) fruits. *Annals of Hort.*, 2(1): 95-97.
- Singh, I. D., 1990, Papaya, New Delhi, India: Oxford and IBH Publishing company private Ltd.
- Singh, J. K., Prasad, J. and Singh, H. K., 2007, Effect of micro-nutrients and plant growth regulators on yield and physic-chemical characteristics of aonla fruits in cv. Narendra Aonla-10. *Indian J. Hort.*, 64 (2): 216-218.

- Singh, P. C., Gangwar, R. S. and Singh, V. K., 2012, Effect of micronutrients spray on fruit drop, fruit quality and yield of aonla cv. Banarasi. *HortFlora Res. Spect.*, 1(1): 73-76.
- Singh, P. C., Gangwar, R. S. and Singh, V. K., 2012, Effect of micronutrients spray on fruit drop, fruit quality and yield of aonla cv. Banarasi. *HortFlora Res. Spect.*, 1(1): 73-76.
- Singh, R., Godara, N. R., Singh, R. and Dahiya, S. S. 2001. Response of foliar application of growth regulators and nutrients in ber (*Zizyphus mauritiana* L.) cv. Umran. *Haryana J. Hort. Sci.*, 30: 161-164.
- Syamal, M. M., Bordoloi, B. and Pakkianathan, K., 2010, Influence of plant growth substances on vegetative growth, flowering, fruiting and fruit quality of papaya. *Indian J. Hort.*, 67(2): 173-176.
- Trivedi, N., Singh, D., Bahadur, V., Prasad, V. M. and Collis, J. P., 2012, Effect of foliar application of zinc and boron on yield and fruit quality of guava (*Psidium guajava* L.). *HortFlora Res. Spect.*, 1(3): 281-283.
- Wali, V. K., Kaul, R. and Kher, R., 2005, Effect of foliar sprays of nitrogen, potassium and zinc on yield and physico-chemical composition of phalsa (*Grewia subinqualis*) cv. Purple Round. *Haryana J. Hort. Sci.*, 34: 56-57.
- Waskela, R. S., Kanpure, R. N., Kumawat, B. R. and Kachouli, B. K., 2013, Effect of foliar spray of micronutrients on growth, yield and quality of guava (*Psidium guajava* L.) cv. Dharidar. *Int. J. agric. Sci.*, 9(2): 551-556.
- Yadav, A. L., Singh, H. K. and Singh, M. K., 2010, Influence of micronutrients of plant growth, yield and quality of papaya fruit (*Carica papaya* L.) cv. Washington. *Asian J. Hort.*, 5(2): 326-329.
- Yadav, A. L., Singh, H. K. and Singh, M. K., 2010, Influence of micronutrients of plant growth, yield and quality of papaya fruit (*Carica papaya* L.) cv. Washington. *Asian J. Hort.*, 5(2): 326-329.

**Table 1. Effect of foliar spray of potassium and micronutrients on yield and yield attributing characters of Papaya *cv.* Red Lady.**

Treatments	Treatment Details	Yield and yield attributing characters								
		Fruit length (cm)	Fruit diameter (cm)	Fruit weight (g)	Number of fruits per plant	Fruit yield per plant (kg)	Fruit yield per ha (ton)	Pulp weight (g)	Peel weight (g)	Seed weight (g)
T <sub>1</sub>	K <sub>2</sub> SO <sub>4</sub> at 1%	21.20	12.95	1865.37	36.88	72.92	181.75	1805.65	43.27	41.23
T <sub>2</sub>	K <sub>2</sub> SO <sub>4</sub> at 2%	20.33	12.05	1812.13	35.84	71.99	179.50	1703.87	41.64	40.28
T <sub>3</sub>	Borax at 0.25%	24.05	14.29	2113.55	42.07	80.48	201.68	2015.27	47.83	45.95
T <sub>4</sub>	Borax at 0.50%	20.65	12.37	1834.29	36.31	72.54	181.08	1764.13	42.82	40.79
T <sub>5</sub>	ZnSO <sub>4</sub> at 0.25%	24.38	14.53	2148.32	42.12	80.79	201.95	2043.56	48.07	46.18

<b>T<sub>6</sub></b>	<b>ZnSO<sub>4</sub> at 0.50%</b>	20.88	12.68	1850.93	36.57	72.68	181.34	1793.21	43.04	41.00
<b>T<sub>7</sub></b>	<b>K<sub>2</sub>SO<sub>4</sub> at 1% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25%</b>	27.20	16.17	2438.00	49.67	90.07	224.96	2343.74	54.98	52.70
<b>T<sub>8</sub></b>	<b>K<sub>2</sub>SO<sub>4</sub> at 2% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25%</b>	26.97	15.84	2397.64	47.22	88.37	221.67	2268.64	52.63	50.65
<b>T<sub>9</sub></b>	<b>K<sub>2</sub>SO<sub>4</sub> at 1% + Borax at 0.50% + ZnSO<sub>4</sub> at 0.50%</b>	20.59	12.21	1822.16	36.18	72.14	180.58	1730.37	42.50	40.63
<b>T<sub>10</sub></b>	<b>K<sub>2</sub>SO<sub>4</sub> at 2% + Borax at 0.50% + ZnSO<sub>4</sub> at 0.50%</b>	20.52	12.13	1819.92	36.01	72.05	179.67	1715.14	41.92	40.42
<b>T<sub>11</sub></b>	<b>Control (water spray).</b>	20.07	11.98	1802.43	33.52	67.65	169.65	1692.28	41.34	40.05
<b>S.Em±</b>		<b>0.62</b>	<b>0.40</b>	<b>69.75</b>	<b>1.66</b>	<b>2.48</b>	<b>6.41</b>	<b>68.08</b>	<b>1.32</b>	<b>1.37</b>
<b>C.D @ 5 %</b>		<b>2.10</b>	<b>1.20</b>	<b>205.76</b>	<b>4.89</b>	<b>7.33</b>	<b>19.30</b>	<b>200.83</b>	<b>4.23</b>	<b>4.05</b>

**Table 2. Effect of foliar spray of potassium and micronutrients on quality characteristics of Papaya cv. Red Lady.**

<b>Treatments</b>	<b>Treatment Details</b>	<b>Quality characteristics</b>					
		<b>TSS (° brix)</b>	<b>Acidity (%)</b>	<b>Ascorbic Acid (mg 100 g<sup>-1</sup>)</b>	<b>Total sugar (%)</b>	<b>Reducing sugar (%)</b>	<b>Non-Reducing sugar (%)</b>
<b>T<sub>1</sub></b>	<b>K<sub>2</sub>SO<sub>4</sub> at 1%</b>	13.44	0.14	53.84	5.48	4.43	1.05
<b>T<sub>2</sub></b>	<b>K<sub>2</sub>SO<sub>4</sub> at 2%</b>	13.20	0.14	52.95	5.28	4.28	1.09
<b>T<sub>3</sub></b>	<b>Borax at 0.25%</b>	13.28	0.14	53.57	5.34	4.38	1.06
<b>T<sub>4</sub></b>	<b>Borax at 0.50%</b>	13.15	0.15	52.80	5.15	4.15	1.10

<b>T<sub>5</sub></b>	<b>ZnSO<sub>4</sub> at 0.25%</b>	12.90	0.15	51.87	5.06	3.90	1.08
<b>T<sub>6</sub></b>	<b>ZnSO<sub>4</sub> at 0.50%</b>	12.84	0.16	51.54	4.98	3.86	1.06
<b>T<sub>7</sub></b>	<b>K<sub>2</sub>SO<sub>4</sub> at 1% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25%</b>	13.52	0.13	54.42	5.64	4.60	1.04
<b>T<sub>8</sub></b>	<b>K<sub>2</sub>SO<sub>4</sub> at 2% + Borax at 0.25% + ZnSO<sub>4</sub> at 0.25%</b>	13.50	0.13	54.07	5.56	4.55	1.02
<b>T<sub>9</sub></b>	<b>K<sub>2</sub>SO<sub>4</sub> at 1% + Borax at 0.50% + ZnSO<sub>4</sub> at 0.50%</b>	13.02	0.15	52.61	5.13	4.01	1.02
<b>T<sub>10</sub></b>	<b>K<sub>2</sub>SO<sub>4</sub> at 2% + Borax at 0.50% + ZnSO<sub>4</sub> at 0.50%</b>	12.98	0.15	52.23	5.10	3.97	1.01
<b>T<sub>11</sub></b>	<b>Control (water spray).</b>	12.52	0.16	50.45	4.84	3.82	1.04
<b>S.Em±</b>		<b>0.42</b>	<b>0.01</b>	<b>0.27</b>	<b>0.23</b>	<b>0.14</b>	<b>NS</b>
<b>C.D @ 5 %</b>		<b>1.22</b>	<b>0.03</b>	<b>0.80</b>	<b>0.56</b>	<b>0.42</b>	<b>NS</b>