

**“ Comparative efficacy and economics of certain chemicals and Biopesticides against podborer, *Helicoverpa armigera* (Hubner) in chickpea at Naini,Prayagraj”**

**ABSTRACT**

---

The experiment was conducted at the research plot of the Department of Agricultural Entomology at Central Research Farm, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during the *rabi* season of 2021. The field was laid in randomised block design (RBD) with seven treatments and one controlled plot. The Mean larval population plant was taken day before and 3, 7 and 14 days after each spray. All the insecticides tested significantly reduced the pest infestation compared to control. The results of the efficacy of treatments of first spray showed that T<sub>4</sub> Spinosad 45% SC is most effective treatment against gram pod borer of Mean larval population producing maximum yield and recorded highest Cost-Benefit ratio compared to other treatments. While T<sub>1</sub> Chlorantraniliprole 18.5 % sc, T<sub>2</sub> Spinetoram 11.7EC, T<sub>3</sub> Flubendamide 39.35% SC , T<sub>4</sub> Spinosad 45% SC, T<sub>5</sub> *Bacillus thuringiensis* has shown averageresults has proved to be least effective chemicals. T<sub>6</sub> *Beauveria bassiana* and T<sub>7</sub> *Metarhizium anisopilae* found to be least effective in managing *Helicoverpa armigera*. When cost benefit ratio was worked out, interesting result was achieved. Among the treatment studied, the best and most economical treatment was Spinosad 45% SC (1:3.98), Spinetoram 11.7EC (1:3.95), followed by Chlorantraniliprole 18.5 % sc (1:3.82), Flubendamide 39.35% SC (1:3.73), *Bacillus thuringiensis* (1:3.21), and *Beauveria bassiana* (1:2.96), The lowest cost benefit ratio was recorded in *Metarhiziun anisopilae* (1:2.80) when compared to Control(1:1.95).

**Key words:** Biopesticides, chemicals, chickpea, effect, *Helicoverpa armigera*, pod borer.

**INTRODUCTION**

---

Chickpea, *Cicer arietinum*, is a member of the legume, pea, or pulse, family "Fabaceae". Chickpea is the common name for an annual plant, *Cicer arietinum*, of the Fabaceae (or Leguminosae) family that is widely cultivated for its typically yellow-brown, pea like seeds. Legumes are multipurpose crops and are consumed either directly as food or in various processed forms or as feed in many farming systems (**Rashid *et al.*, 2014**)<sup>[8]</sup>

Madhya Pradesh ranked first contributing an area of 30.76 lakh ha, production 33.98 lakh tonnes and productivity 1105 kg/ha (34.46% and 40.62% of total area and production of country). Maharashtra is one of the second rank for area 15.41 lakh ha (17.26%) and third for production 11.98 lakh tones (14.32%). Where as, Rajasthan stood second in production (14.47%) and third in area (15.37%). The highest yield was recorded in the state of Telangana (1459 kg/ha) followed by Gujarat (1201 kg/ha) and West Bengal (1163 kg/ha). The lowest yield was recorded in Karnataka (578 kg/ha). (**Annual Report DPD 2017**).

The crop has multiple uses in rural as well as urban India. Chickpea is a good source of protein (20 mg/100 g), carbohydrate and minerals, also its posses a high nutritional value. 100 g of gram seed provides 358 calories which is more than that of any other legume, except groundnut and lupine seeds (Kanwar, 1979). It is good source of amino acid. The amino acid content per gram of chickpea is 0.44 mg lysine, 0.30 mg thiamine, 0.51 mg riboflavin and 2.1 mg niacin. Chickpea seeds are good source of Vitamin A [I.U.]- (316 mg), vitamin C-(3 mg), vitamin K-(0.29 mg) and minerals (12 mg) along with the Folic-acid (125g/100g). Germinated seeds are recommended against scurvy disease. Chickpea also contains 56.5 per cent carbohydrate, besides ash, calcium and iron etc.

Among biotic factors chickpea is infested by nearly 60 insect's species in which cutworm, *Agrotis ipsilon* (Ratt.), gram pod borer, *Helicoverpa armigera* (Hubner), semilooper, *Autographa nigrisigna* (Walk.), and aphid, *Aphis craccivora* (Koch.) are the pests of major importance (**Acharjee and Sharma, 2013**)<sup>[1]</sup>. Among these, the major damage is caused by gram pod borer which is polyphagous in nature; *Helicoverpa armigera* is one of the serious pests of chickpea, which feeds more than 150 crops throughout the world (**Vinutha *et al.*, 2013**)<sup>[11]</sup>. Gram pod borer is widely distributed and a serious pest of chickpea causing heavy crop losses (20- 60%) throughout the India.

## MATERIALS AND METHODS

---

The experiment was conducted during *rabi* season 2021 at the Central Research Farm (CRF) of Sam Higginbottom University of Agriculture, Technology and Sciences, Naini, Prayagraj, Uttar Pradesh, India, in a randomized block design with eight treatments replicated three times using Ankur-chirag (local variety) in a plot size of 2m×2m at a spacing of 30×10cm with a recommended package of practices excluding plant protection. Seven treatments of chemicals and biopesticides were evaluated against, *Helicoverpa armigera* i.e., T<sub>1</sub> Chlorantraniliprole 18.5 % sc, T<sub>2</sub> Spinetoram 11.7EC, T<sub>3</sub> Flubendamide 39.35% SC , T<sub>4</sub> Spinosad 45% SC, T<sub>5</sub> *Bacillus thuringiensis*, T<sub>6</sub> *Beauveria bassiana* , T<sub>7</sub> *Metarhizium anisopilae* and T<sub>0</sub> control plot. The population of chickpea pod borer was recorded before 1-day spraying and on 3rd day, 7th day and 14th day after insecticidal application. The populations of chickpea pod borer was recorded on 5 randomly selected and tagged plants from each plot for investigating larval population and cost benefit ratio.

### Cost benefit ratio:

Based on the yield data, the gross returns and net returns were calculated for each treatment. The benefit cost ratio (BCR) was determined by dividing the additional returns with the additional cost of imposing the respective treatment on hectare basis.

$$B.C.R = \frac{\text{Gross Returns}}{\text{Total Cost of Protection}}$$

## RESULTS AND DISCUSSION

---

The results of the experiment Evaluation of different insecticides chickpea pod borer *Helicoverpa armigera* (Hubner) to study cost benefit ratio during *rabi* season of 2021-2022. The data so obtained through observation on various aspects were subjected to statistical analysis wherever necessary and the compiled mean data are tabulated in the following pages. Results obtained are presented aspect wise here under.

The data revealed on population of *Helicoverpa armigera* over control on mean (3,7 and 14 DAS) 1<sup>st</sup> spray revealed that all the treatments were significantly superior over control (3.28). Among all the treatments minimum larval population was recorded in T4 Spinosad 45% SC (1.44), followed by T1 Chlorantraniliprole 18.5% SC (1.59) T2 Spinetoram 11.7EC (1.79), T3 Flubendamide 39.35% SC ( 1.99), T5 *Bacillus thuringiensis* (2.23), T6 *Beauveria bassiana* (2.33). In this the maximum larval population was recorded in T7 *Metarhizium anisopliae* (2.46).

Table.1. Efficacy of certain chemicals and biopesticides on the population of pod borer, *H. armigera* on chickpea during *rabi* season of 2021-22  
( 1<sup>st</sup> Spray)

	Treatment	Average number of Larvae/5Plants(1 <sup>st</sup> spray)				
		1DBS	3DAS	7DAS	14DAS	Mean
T <sub>4</sub>	Spinosad 45% SC	2.8	1.66	1.13	1.53	1.44
T <sub>1</sub>	Chlorantraniliprole 18.5% SC	2.86	1.86	1.26	1.66	1.59
T <sub>2</sub>	Spinoterum 11.7 EC	2.8	2.06	1.46	1.86	1.79
T <sub>3</sub>	Flubendamide 39.35% SC	3.00	2.26	1.66	2.06	1.99
T <sub>5</sub>	<i>Bacillus thuringiensis</i>	2.93	2.46	1.86	2.26	2.19
T <sub>6</sub>	<i>Beauveria bassiana</i>	3.06	2.6	2.00	2.4	2.33
T <sub>7</sub>	<i>Metarhizium anisopilae</i>	2.93	2.73	2.13	2.53	2.46
T <sub>0</sub>	Control	3.00	3.13	3.26	3.46	3.28
	F-test	NS	S	S	S	S
	C.D. at 0.5%	---	0.111	0.130	0.173	0.134
	S.EdA (±)	0.22	0.044	0.054	0.077	0.063

The data revealed on population of *Helicoverpa armigera* over control on mean (3,7 and 14 DAS) 2<sup>nd</sup> spray revealed that all the treatments were significantly superior over control

(3.64 ). Among all the treatments minimum larval population was recorded in T<sub>4</sub> Spinosad 45% SC (0.94), followed by T<sub>1</sub> Chlorantraniliprole 18.5 % sc 18.5 SC (1.26), T<sub>2</sub> Spinetoram 11.7EC (1.24),T<sub>3</sub> Flubendamide 39.35% SC (1.44), T<sub>5</sub> *Bacillus thuringiensis* (1.64), T<sub>6</sub> *Beauveria bassiana* (1.82).In this the maximum larval population was recorded in T<sub>7</sub> *Metarhizium anisopliae* (1.91).

Table.2. Efficacy of certain chemicals and biopesticides on the population of pod borer, *H. armigera* on chickpea during *rabi* season of 2021-22 - ( 2<sup>nd</sup> Spray)

	Treatment	Average number of Larvae/5 Plants(2 <sup>nd</sup> spray)				
		1DBS	3DAS	7DAS	14DAS	Mean
T <sub>4</sub>	Spinosad 45% SC	1.53	1.13	0.66	1.06	0.95
T <sub>1</sub>	Chlorantraniliprole 18.5% SC	1.66	1.53	1.06	1.2	1.26
T <sub>2</sub>	Spinetoram 11.7 EC	1.86	1.46	0.93	1.33	1.24
T <sub>3</sub>	Flubendamide 39.35% SC	2.06	1.66	1.13	1.53	1.44
T <sub>5</sub>	<i>Bacillus thuringiensis</i>	2.26	1.86	1.33	1.73	1.64
T <sub>6</sub>	<i>Beauveria bassiana</i>	2.4	2.00	1.53	1.93	1.82
T <sub>7</sub>	<i>Metarhizium anisopliae</i>	2.53	2.13	1.6	2.00	1.91
T <sub>0</sub>	Control	3.46	3.6	3.6	3.73	3.64
	F-test	S	S	S	S	S
	C.D. at 0.5%	0.173	0.353	0.302	0.333	0.355
	S.EdA (±)	0.812	0.522	0.141	0.154	0.165

The data revealed on population of *Helicoverpa armigera* over control on Overall mean revealed that all the treatments were significantly superior over control (3.46 ). Among all the treatments minimum larval population was recorded in T<sub>4</sub> Spinosad 45% SC (1.19), followed by T<sub>1</sub> Chlorantraniliprole 18.5 % SC (1.43) T<sub>2</sub> Spinetoram 11.7EC (1.52), T<sub>3</sub> Flubendamide 39.35% SC ( 1.72), T<sub>5</sub> *Bacillus thuringiensis* (1.92), T<sub>6</sub> *Beauveria bassiana* (2.07). In this the maximum larval population was recorded in T<sub>7</sub> *Metarhizium anisopliae* (2.18).


Table.3 Efficacy of certain chemicals and biopesticides on the population of pod borer, *H.armigera* on chickpea during *rabi* season of 2021-22 - (1<sup>st</sup> and 2<sup>nd</sup> Spray).

S.NO	Treatments	Over all mean population/5 plants		1 <sup>st</sup> & 2 <sup>nd</sup> spray
		1 <sup>st</sup> Spray	2 <sup>nd</sup> Spray	
T <sub>4</sub>	Spinosad 45% SC	1.44	0.955	1.995
T <sub>1</sub>	Chlorantraniliprole 18.5% SC	1.599	1.266	1.432
T <sub>2</sub>	Spinetoram 11.7 EC	1.799	1.242	1.520
T <sub>3</sub>	Flubendamide 39.35% SC	1.999	1.443	1.721
T <sub>5</sub>	<i>Bacillus thuringiensis</i>	2.199	1.643	1.921
T <sub>6</sub>	<i>Beauveria bassiana</i>	2.333	1.822	2.077
T <sub>7</sub>	<i>Metarhizium anisopliae</i>	2.466	1.911	2.188
T <sub>0</sub>	Control	3.287	3.644	3.465

Table .4 Economics of Cultivation:


Treatments		Yield q/ha	Cost of yield q/₹	Total cost of yield in ₹	Common cost ₹	Treatment cost ₹	Total treatment cost ₹	C:B Ratio
T4	Spinosad 45% SC	23.52	5500	129630	25365	7195	32560	1:3.98
T1	Chlorantraniliprole 18.5 % sc	22.75	5500	125125	25365	7390	32755	1:3.82
T2	Spinetoram 11.7EC	20.50	5500	112750	25365	3135	28500	1:3.95
T3	Flubendamide 39.35% SC	18.00	5500	99000	25365	1144	26509	1:3.73
T5	<i>Bacillus thuringiensis</i>	16.24	5500	89320	25365	2440	27805	1:3.21
T6	<i>Beauveria bassiana</i>	14.42	5500	79310	25365	1420	26785	1:2.96
T7	<i>Metarhizium anisopilae</i>	14	5500	77000	25365	2050	27415	1:2.80
T0	Control	9.02	5500	49610	25365	----	25365	1:1.95

**Figure.1 Graphical representation of Yield**



The yields among the treatment were significant. The highest yield was recorded in T<sub>4</sub> Spinosad 45% SC (23.52q/ha), followed by T<sub>1</sub> Chlorantraniliprole 18.5 % SC (22.75q/ha), T<sub>2</sub> Spinetoram 11.7EC (20.50q/ha), T<sub>3</sub> Flubendamide 39.35% SC (18.00q/ha), T<sub>5</sub> *Bacillus thuringiensis* (16.24q/ha), T<sub>6</sub> *Beauveria bassiana* (16.24q/ha) and T<sub>7</sub> *Metarhizium annisopliae* (14.00q/ha) and T<sub>0</sub> control (9.02q/ha).

**Figure.2 Graphical representation of Cost Benefit Ratio.**



When cost benefit ratio worked out, interesting result was achieved, among the treatment studied, the best and most economical treatment T<sub>4</sub> Spinosad 45% SC 45% SC (1:3.98), followed by T<sub>2</sub> Spinetoram 11.7EC (1:3.95), T<sub>1</sub> Chlorantraniliprole 18.5 % SC (1:3.82), T<sub>3</sub> Flubendamide 39.35% SC (1:3.73), T<sub>5</sub> *Bacillus thuringiensis* (1:3.21), T<sub>6</sub> *Beauveria bassiana* (1:2.96) and T<sub>7</sub> *Metarhizium anisopliae* (1:2.80) and T<sub>0</sub> control (1:1.95).

## DISCUSSION

Overall mean revealed that all the treatments were significantly superior over control (3.67). Among all the treatments minimum larval population was recorded in T<sub>4</sub> Spinosad 45% SC (1.99) these results are similar to the findings of **Suryavanshi et al., (2008)**, followed by T<sub>1</sub> Chlorantraniliprole 18.5 % SC (1.43), T<sub>2</sub> Spinetoram 11.7EC (1.52), T<sub>3</sub> Flubendamide 39.35% SC (1.72), T<sub>5</sub> *Bacillus thuringiensis* (1.92), T<sub>6</sub> *Beauveria bassiana* (2.07). In this the maximum larval population was recorded in T<sub>7</sub> *Metarhizium anisopliae* (2.18) and control plot (3.46).

All the insecticides were found very effective and significantly superior over control. The minimum larval population was recorded in T<sub>4</sub> Spinosad 45% SC yield (23.52q/ha) these results are similar to the findings of **Suryavanshi et al., (2008)**, T<sub>1</sub> Chlorantraniliprole 18.5 % SC and maximum yield (22.75q/ha) these results are similar to the findings of followed by **Kapulai**

**Santhosh and Ashwani Kumar (2022)** T<sub>2</sub> Spinetoram 11.7EC yield (20.50q/ha) these results are similar to the findings of **Akbar et al., (2017)**<sup>[2]</sup> followed by T<sub>3</sub> Flubendamide 39.35% SC yield (18.00 q/ha) these results are similar to the findings of **Kapulai Santhosh and Ashwani Kumar (2022)**, T<sub>5</sub> *Bacillus thuringiensis* yield ( 16.24 q/ha) these results are similar to the findings of **S. Bhushan et al., (2011)**<sup>[3]</sup>. T<sub>6</sub> *Beauveria bassiana* yield (14.42q/ha), in this the maximum larval population was recorded in T<sub>7</sub> *Metarhizium anisopliae* with minimum yield (14.00q/ha) and control(9.02q/ha).

When cost benefit ratio worked out, interesting result was achieved, among the treatment studied, the best and most economical treatment T<sub>4</sub> Spinosad 45% SC 45% SC (1:3.98), followed by T<sub>2</sub> Spinetoram 11.7EC (1:3.95), T<sub>1</sub> Chlorantraniliprole 18.5 % SC (1:3.82), T<sub>3</sub> Flubendamide 39.35% SC (1:3.73), T<sub>5</sub> *Bacillus thuringiensis* (1:3.21), T<sub>6</sub> *Beauveria bassiana* (1:2.96) and T<sub>7</sub> *Metarhizium annisopliae* (1:2.80) and T<sub>0</sub> control (1:1.95).

## CONCLUSION

From the present study, the results it showed that T<sub>4</sub> Spinosad 45% SC most effective treatment against gram pod borer of Mean larval population and producing maximum yield and recorded highest Cost-Benefit ratio compared to other treatments. While T<sub>1</sub> Chlorantraniliprole 18.5 % SC, T<sub>2</sub> Spinetoram 11.7EC, T<sub>3</sub> Flubendamide, has shown average results has proved to be least effective chemicals. *Bacillus thuringiensis*, *Beauveria bassiana*, and *Metarhizium annisopliae* found to be least effective in managing *Helicoverpa armigera*. Botanicals are the part of integrated pest management in order to avoid indiscriminate use of pesticides causing pollution in the environment and not much harmful to beneficial insects.

## Reference

1. **Acharjee, S. and Sharma, B.K.(2013).** Transgenic *Bacillus thuringiensis* (Bt) chickpea: India's most wanted genetically modified (GM) pulse crop. *African Journal of Biotechnology*, **12** (39): 5709-5713.
2. **Akbar, W., Usman, M. A., Memon, R. A., Moula, B. and Sohail, M.(2017).** Validation of some new chemistry and conventional insecticides against gram pod borer (*Helicoverpa armigera*) in chickpea. *Pakistan Entomologist*, **40**(1):45-49.
3. **Bhushan, S., Singh, R. P., and Shanker, R. (2011).** Bioefficacy of neem and Bt against pod borer, *Helicoverpa armigera* in chickpea. *Journal of Biopesticides*, **4**(1), 87.
4. **Deshmukh, S. G., Sureja, B. V., Jethva, D. M., & Chatar, V. P.(2010).** Field efficacy of different insecticides against *Helicoverpa armigera* (Hubner) infesting chickpea. *Legume Research*, **33**(4):269-273.
5. **Gayathri, L. and Kumar, A.(2021).** Field efficacy of certain insecticides against pod borer, *Helicoverpa armigera* (Hubner) on chick pea in Prayagraj. *Journal of Entomology and Zoology Studies*, **9**(3): 280-283.
6. **Ghugal, S.G., Shrivastava, S.K., Bhowmick, A.K. and Saxena, A.K.(2013).** Management of *Helicoverpa armigera* (Hubner) in chickpea with biopesticides. *Jawaharlal Nehru Krishi Vishwa Vidyalaya Research Journal* , **47**(1): 84-87.
7. **Kumar P., Kumar S., Rana R., and Sachin S.K.,(2014).** Comparative efficacy of novel insecticides and bio-pesticides on larval population density of gram pod borer (*Helicoverpa armigera* hubner) on chickpea. *Journal of Plant Development Sciences*, **6**(2): 335-338.
8. **Rashid, A., Hossain, S., Deb, U., Kumara Charyulu, D., Shyam, D. M. and Bantilan, C.(2014).** Targeting and introduction of Chickpea improved cultivars in Barind region of Bangladesh, (Tropical Legumes II Phase 2 Project).
9. **Srikanth M., Lakshmi M.S.M. and Dr. Y. Koteswar Rao.(2014).** Bio-efficacy and economics of certain new insecticides against gram pod borer, *Helicoverpa armigera* (Hubner). *International Journal of Plant, Animal and Environmental Sciences*, **4** (1):1-5.

10. **Upadhyay, R. R., Singh, P. S. and Singh, S. K.(2020).** Comparative efficacy and economics of certain insecticides against gram pod borer, *Helicoverpa armigera* (Hübner) in chickpea. *Indian Journal of Plant Protection*, **48**(4): 403-410.
11. **Vinutha, J.S., Bhagat, D. and Bakthavatsalam, N.(2013).** Nano technology in the management of polyphagous pest *Helicoverpa armigera* *Journal of Academic. Research*, **1** (10): 606-608.
12. **Yogeeswarudu, B. and Venkata, K.K.(2014).** Field studies on efficacy of novel insecticides against *Helicoverpa armigera* (Hubner) infesting on chickpea. *Journal of Entomology and Zoology studies*, **2**(4): 286-289.

UNDER PEER REVIEW