

Pre- Sowing Seed Treatments of Selected Organics on Growth, Yield and Yield Attributing Characters of Cowpea (*Vigna unguiculata* L.)

Abstract

During Kharif, 2021, present trail was carried out at the field experimentation center of the Genetics and Plant Breeding, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Allahabad. An experiment entitled **Pre- Sowing Seed Treatments of Selected Organics on Growth, Yield and Yield Attributing Characters of Cowpea (*Vigna unguiculata* L.)**. For this purpose ,13 priming treatments including Control on Cowpea seeds variety were used to study under field conditions. Field experiment was laid out in Randomized Block Design (RBD) with three replications respectively. T₀- Control, T₁ Moringa Leaves Extract 2% (8 hrs), T₂ Moringa Leaves Extract 4% (8 hrs), T₃ Moringa Leaves Extract 6% (8 hrs), T₄ Neem Leaves Extract 2% (8 hrs), T₅ Neem Leaves Extract 4% (8 hrs), T₆ Neem Leaves Extract 6% (8 hrs), T₇ Panchagavya 2% (8 hrs), T₈ Panchagavya 4% (8 hrs), T₉ Panchagavya 6% (8 hrs), T₁₀ Beejamruth 2% (8 hrs), T₁₁ Beejamruth 4% (8 hrs), T₁₂ Beejamruth (8 hrs). To find out Influence of different seed treatment on growth, yield and seed quality parameters of cowpea showed that significant treatment on Rate of Field emergence, Days to 50% Flowering, Plant Height, Number of primary branches, Days to Maturity, Number of pods per plant, Number of seeds per pod, 100 seed weight, Seed yield per plant, Seed yield per plot, biological yield, Harvest index. Pre-sowing treatment which is concluded from the present study that the seeds of Cowpea (Variety – Bali 265) were treated with T₉ Panchagavya 6 % (8 hrs) enhanced the above-mentioned characters.

Key words: *Cowpea, Priming, Seed treatment, Panchagavya, Beejamruth.*

INTRODUCTION

Cowpea has unique place in Indian agriculture particularly from the nutritional and soil fertility point of view. It is well adapted to stress condition and possesses excellent nutritional quality. It is a multipurpose crop. The immature pods and leaves which are used as vegetable, the stem or haulm serves as fodder for livestock. The mature beans used as pulses. It is likely that the crop was first introduced to India during the Neolithic period, and therefore India appears to be a secondary centre of genetic diversity (**Pant et al. 1982**). "Yardlong beans," a unique cultivar group (*Sesquipedialis*) of cowpea that produces very long pods widely consumed in Asia as a fresh green or "snap" bean, apparently evolved in Asia and is rare in African landrace germplasm.

The cowpea plant is an annual, erect or semi-erect, reaching a height of 2 m. It is slightly hairy with a well-developed root system. The flower colour varies through different shades of purple, pink, yellow and white. The pods contain 6 to 13 small, kidney shaped seeds. Cowpea (*Vigna sinensis* L.) is believed to have originated in Africa, Cowpea belongs to family leguminaceae. Recently, It is the important pulse crop cultivated throughout India. In India, cowpea is consumed in several ways in southern India as whole grain. It is an important dietary protein food to humans. In addition to protein (23.86 g), it supplies fibre (16.3 g), fat (1.15 g), vitamins like A, C, B, niacin, minerals like calcium, magnesium, potassium,

phosphorus, sodium, sugars and carbohydrates (60.3 g) per 100 g of grain. (Rosaiah et al., 1993) and known by many vernacular names like Lobia (Hindi), Alasande (Kannada) and Karamani (Tamil and Telugu) (Javeeda, 2001).

Panchagavya means “Mixture of five products (dung, urine, milk, ghee, and curd) of the cow”. Of these the direct constituents are cow dung, urine, milk and the two derived products are ghee and curd. It is used as fertilizer and pesticides in agriculture operation. Panchagavya plays an important role in the quality of fruits and vegetables (Krishna Kumar, 2020).

Beejamruth is used as a treatment for seeds, seedlings or any planting plant material. It is very effective in protecting young roots from fungus and other microbes as well as from soil borne and seed-borne diseases that commonly affect plants after the monsoon season. Beejamruth is composed of similar ingredients as Jeevamrutha. Maximum microbial population was observed on the day of preparation in beejamruth. observed on the day of preparation in beejamrutha and on 10th day after preparation in jeevamrutha. The presence of beneficial microorganisms in this liquid formulation might be mainly due to their constituents such as: cow dung, cow urine, legume flour and jaggery containing both macro and essential micro nutrients, many vitamins, essential amino acids, growth promoting substances like indole acetic acid (IAA), gibberellic acid (GA) and beneficial microorganisms (Neelima and Sreenivasa, 2011).

Moringa (*Moringa oleifera*) is well known as a miracle tree. Zeatin stimulates cell division and cell elongation. Disease-free leaves with no more than 40 days old are suitable for leaf extraction because it contains higher number of bioactive materials than older once (Michel, 2008). The most common method of preparing leaf extract from fresh moringa leaves is collecting 200g of fresh leaves then clean, wash and store them overnight at freezing temperature. After 24 hours, crash the stored leaves through mortar and pestle or blender and sieve through cheesecloth to collect the extract and dilute based on the plant requirement (Nouman, 2012).

Neem leaves are also used as a good priming agents contain flavonoids, steroids, carbohydrates, glycoside, antiquinone, terpenoids and alkaloids. It was shown that neem leaf extract has antifungal effects and can be used as fungicidal seed treatments for the control of seedborne diseases and for increasing seed germination and seedling emergence. Neem leaf extract was prepared using following method, the fresh neem leaves were collected separately and dried under shade. The shade dried leaves are powdered using mortar and pestle or electric grinder. Then exactly weighed one gram of leaf powder using weighing balance and dissolved in 100 ml of distilled water which was measured already in the beaker to make 1% neem leaf extract. The prepared neem leaf extract was filtered by using muslin cloth to remove unwanted material and leaf debris.

Research was carried out with following **objectives**:

1 To determine the effect of different doses of seed treatment of Neem leaves Extract, Moringa Leaves Extract ,Panchagavya and Beejamruth on growth, yield and yield attributing traitss of Cowpea CV. BALI265

2 To standardize the suitable doses of pre-sowing seed treatment of Neem leaves extract, Moringa leaves extract, Panchagavya and Beejamruth seed treatment for cowpea crop

MATERIAL AND METHODS

The present investigation was carried out to study the effect of Pre-sowing seed treatments of selected Organics on growth, yield and yield attributing traits of Cowpea (*Vigna unguiculata* L.) at the Field experimentation centre of Seed Science and Technology in the Department of Genetics and Plant Breeding, Sam Higginbottom Institute of Agriculture, Technology and Science, Naini Agriculture Institute, Prayagraj (U.P). Field experiment was laid out in Randomized Block Design (RBD) with treatment material consists of 12 treatments and untreated (control) seed of Cowpea and three replications respectively during Kharif 2021-22. viz., T₀- Control, T₁ – Moringa leaves extract priming @ 2%, 8 hrs, T₂- Moringa leaves extract @ 4%, 8hrs, T₃-Moringa leaves extract @ 6%, 8hrs, T₄- Neem leaves extract @ 2%, 8hrs, T₅- Neem leaves extract @ 4%, 8hrs, T₆- Neem leaves extract @ 6%, 8hrs, T₇- Panchagavya @ 2% 8 hrs, T₈- Panchagavya @ 4% 8 hrs, T₉ – Panchagavya @ 6%, 8hrs, T₁₀- Beejamruth – 2% 8hrs, T₁₁- Beejamruth priming – 4% 8hrs, T₁₂- Beejamruth @ 6% 8hrs, with the soaking durations of 8hrs with water. For the preparation of Panchagavya first mix thoroughly the fresh cow dung and ghee then incubate it for 2 days, after incubation add cow urine and 5 liters of water and mix it properly in morning and evening for one week, then add gram powder (500gm) along with cow milk and curd and mix it properly in the morning and evening for two weeks, after two weeks Panchagavya is ready and can be used directly. The unfortified seed served as control. The soaked seeds were surface dried for one day and were evaluated for the growth, yield attributing parameters viz Maximum Rate of Field emergence, Days to 50% Flowering, Plant Height(cm) (30,60,90 DAS), Number of primary branches, Days to Maturity, Number of pods per plant, Number of seeds per pod, 100 seed weight, Seed yield per plant, Seed yield per plot, biological yield, Harvest index (%) to find the best treatment results was observed in Cowpea variety (Bali 265).

RESULTS AND DISCUSSION

PRE-HARVEST PARAMETERS

Significantly, maximum Rate of field emergence (680.8) was recorded in the treatment T₉ Panchagavya 6% 8hrs and it was followed by T₁₁ Beejamruth 4% 8hrs (653). Minimum field emergence was recorded by T₀ – Control (436.3). Significantly, maximum height of plant (117.46 cm) was recorded by T₉ Panchagavya 6% 8hrs and it was followed by T₁₁ Beejamruth 4% 8hrs (103.13). Minimum plant height was recorded by T₀ – Control (67.50 cm). Significantly, maximum number of branches (5.00) was recorded by T₉ Panchagavya 6% 8hrs and it was followed by T₁₁ Beejamruth 4% 8hrs (4.86). Minimum number of branches was recorded by T₀– Control (3.00). Significantly, Maximum days to 50% flowering was recorded by T₀-Control (62.66) and Minimum was recorded by T₉

Panchagavya 6% 8hrs (32.56) followed by T₁₁ Beejamruth 4% 8hrs (32.76). Significantly, Maximum days to Maturity was recorded by T₀ Control (84.66) and Minimum was recorded

Notation	Treatments	Rate of Field Emergence	Days to 50% flowering	Plant height (cm) at 90 DAS	Number of Primary branches	Days To maturity
----------	------------	-------------------------	-----------------------	-----------------------------	----------------------------	------------------

by T₉-Panchagavya 6% 8hrs (72.00) followed by T₁₁ Beejamruth 4% 8hrs (72.33). Similar findings were recorded in cowpea by **Chandra et. al., (2021)** and **Khan et. al., (2021)**. Statistically analysed data was mentioned in table no. 1.

Table No. 1 Pre harvest observation mean performance of Cowpea for growth and yield parameters

T0	Control	436.3	62.66	67.50	3.00	84.66
T1	Moringa leaves extract 2% (8hrs)	575.3	35.00	88.40	3.66	76.00
T2	Moringa leaves extract 4% (8hrs)	570	35.33	78.93	3.83	74.66
T3	Moringa leaves extract 6% (8hrs)	565.2	33.33	83.70	4.00	75.33
T4	Neem leaves extract 2% (8hrs)	605.7	35.66	94.83	4.50	77.55
T5	Neem leaves Extract 4% (8hrs)	559.8	35.33	77.40	4.03	73.66
T6	Neem leaves extract 6% (8hrs)	550.4	36.23	89.26	4.40	76.33
T7	Panchagavya 2% (8hrs)	552.3	34.00	80.50	4.66	83.33
T8	Panchagavya 4% (8hrs)	583.4	36.43	95.40	4.16	77.66
T9	Panchagavya 6% (8hrs)	680.8	32.56	117.46	5.00	72.00
T10	Beejamruth 2%(8hrs)	586.9	33.00	79.60	3.50	74.00
T11	Beejamruth 4% (8hrs)	653	32.76	103.13	4.86	72.33
T12	Beejamruth 6% (8hrs)	582.7	61.66	74.80	4.26	76.33
	Grand Mean	577.06	38.77	87.00	4.14	76.46
	F test	S	S	S	S	S
	CD_{0.05}	1.20	1.30	8.64	0.92	1.82
	SE(m)	0.41	0.44	2.96	0.31	0.62
	SE(d)	0.59	0.63	4.19	0.45	0.88
	C.V	4.79	1.98	5.89	13.30	1.41

POST-HARVEST PARAMETERS

Significantly, Maximum Number of Pods per plant was recorded by T₉ Panchagavya 6% 8hrs (13.46) and it was followed by T₁₁ Beejamruth 4% 8hrs (13.33), Minimum number of Pods per plant was recorded by T₀– Control (5.80). Significantly, Maximum Number of seeds per pod was recorded by T₉ Panchagavya 6% 8hrs (16.13) and it was followed by, T₁₁Beejamruth 4% 8hrs (14.60) and Minimum Number of seeds per pod was recorded by T₀– Control (9.85). Significantly, Maximum 100 seed weight was recorded by T₉ Panchagavya 6% 8hrs (12.40g) followed by T₁₁ Beejamruth 4% 8hrs (12.16 g) and. Minimum 100 seed weight was recorded by T₀– Control (9.10g). Significantly, Maximum seed yield per plant was recorded by T₉ Panchagavya 6% 8hrs (18.55 g) followed by T₁₁ Beejamruth 4% 8hrs (16.29 g). Minimum seed yield per plant was recorded by T₀– Control (9.09 g). Significantly, maximum seed yield per plot was recorded by T₉ Panchagavya 6% 8hrs (1613.90g) and it was followed by T₁₁ Beejamruth 4% 8hrs (1295.80 g) and Minimum seed yield per plot was recorded by T₀– Control (615.80g). Significantly, maximum biological yield was recorded by T₉ Panchagavya 6% 8hrs (4636.94g) and it was followed by T₁₁ Beejamruth 4% 8hrs (3800.01g). Minimum biological yield was recorded by T₀ (2689.21 g). Significantly, maximum harvest index was recorded by T₉ Panchagavya 6% 8hrs (34.80 %) and it was

followed by T₁₁ Beejamruth 4% 8hrs (34.10%). Minimum harvest index was recorded by T₀– Control (22.90%). This was might be due to better water imbibition due to hydro priming and priming with Panchagavya, because seeds have a thick outer coat and they might take more time to start germination if sown unprimed because water imbibition is the first step of germination and insufficient moisture level hampers germination process. Similar findings were recorded in cowpea by **Farooq *et. al.*, (2019)** and **Moosavi *et. al.*, (2011)**. Statistically analysed data was mentioned in table no. 2.

Table No. 2 Post- harvest observation mean performance of Cowpea for growth and yield parameters

Notation	Treatments	Number of pods per plant	Number of seeds per pod	100 Seed Weight (g)	Seed yield per plant (g)	Seed yield per plot (g)	Biological Yield (g)	Harvest index (%)
T ₀	Control	5.80	9.85	9.10	9.09	615.80	2689.21	22.90
T ₁	Moringa leaves extract 2% (8hrs)	11.23	11.73	11.12	10.94	904.20	3391.59	26.66
T ₂	Moringa leaves extract 4% (8hrs)	10.83	10.46	8.17	10.19	797.80	3264.63	24.44
T ₃	Moringa leaves extract 6% (8hrs)	6.13	10.90	9.81	11.89	1050.40	3453.24	30.42
T ₄	Neem leaves extract 2%(8hrs)	9.66	12.75	10.53	11.29	997.06	3461.11	28.81
T ₅	Neem leaves Extract 4% (8hrs)	7.86	11.36	9.61	12.74	1120.70	3444.08	32.54
T ₆	Neem leaves extract 6% (8hrs)	10.46	13.93	11.70	11.62	1010.20	3384.25	29.85
T ₇	Panchagavya 2% (8hrs)	11.66	12.01	10.21	12.18	1080.86	3404.45	31.75
T ₈	Panchagavya 4% (8hrs)	12.16	14.21	12.03	13.49	1165.00	3467.29	33.60
T ₉	Panchagavya 6% (8hrs)	13.46	16.13	12.40	18.55	1613.90	4636.94	34.80
T ₁₀	Beejamruth 2% (8hrs)	8.33	13.53	11.31	9.86	802.23	2948.69	27.21
T ₁₁	Beejamruth 4% (8hrs)	13.33	14.60	12.16	16.29	1295.80	3800.01	34.10
T ₁₂	Beejamruth 6% (8hrs)	9.00	11.62	10.80	11.32	960.51	3427.99	28.02
	Grand Mean	10.00	12.55	10.69	12.26	1031.89	3444.12	29.62
	F test	S	S	S	S	S	S	S
	CD_{0.05}	1.76	1.44	0.40	1.55	46.63	70.87	0.57
	SE(m)	0.60	0.49	0.14	0.53	15.98	24.28	0.19
	SE(d)	0.85	0.70	0.20	0.75	22.59	34.34	0.27
	C.V	10.46	6.83	2.67	7.51	2.68	1.22	1.14

Conclusion:

It is concluded from the present study that the seeds of Cowpea (Variety – Bali 265) treated with Panchagavya @ 6% gave better performance than other treatments viz, Rate of field emergence (680.8), Days to 50% flowering (32.56), Plant height (117.46 cm), Number of primary branches per plant (5.00), Number of pods per plant (13.46), Number of seeds per pod (16.13), Days to Maturity (72.00), 100 Seed weight (12.40 g), Seed yield per plant (18.55 g), Seed yield per plot (1613.90 g), Biological yield (4636.94 g), Harvest index (34.80 %) followed by Beejamruth 4% 8hrs were recorded significantly higher compared to Control (Untreated). These conclusions are based on the results of crop duration investigation and therefore further investigation is needed to arrive at valid recommendation.

REFERENCES

- Ali, Q. R., Perveen, M. A., El-Esawi, S., Ali, S. M., Hussain, M., Amber, N., Iqbal, M., Rizwan, M. N., Alyemeni, H. A., El-Serehy, F. A., Al-Misned and Ahmad P. 2020.** "Low Doses of *Cuscutareflexa* Extract Act as Natural Biostimulants to Improve the Germination Vigor, Growth, and Grain Yield of Wheat Grown under Water Stress: Photosynthetic Pigments, Antioxidative Defense Mechanisms, and Nutrient Acquisition." *Biomolecules* **10** (9).
- Farooq, Muhammad Usman, Faisal Nadeem, Hafeez ur Rehman, Abdul Wahid, Shahzad MA Basra, and Kadambot HM Siddique. 2019.** "Seed priming in field crops: potential benefits, adoption and challenges." *Crop and Pasture Science*, **70** (9):731-771.
- Gella, Dessalegne, Habtamu Ashagre, and Takele Negewo. 2013.** "Allelopathic effect of aqueous extracts of major weed species plant parts on germination and growth of wheat." *Journal of Agricultural and Crop Research* **1** (3):30-35.
- Gunasekar, J, A Kamaraj, and S Padmavathi. 2017.** "EFFECT OF BOTANICAL SEED PRIMING ON SEED QUALITY CHARACTERS IN BLACKGRAM [VIGNAMUNGO (L.) HEPPER] cv. CO6." *Plant Archives* **17** (2):1383-1387.
- Kale, Dhananjay B, Prashant Kumar Rai, HIRAKANT V KALPANDE, and AK Rai. 2019.** "Studies on Organic Priming on Germination and Seedling Vigour in Sorghum [Sorghum bicolor (L.).]" *Int. J. Pure App. Biosci* **7** (1):511-519.
- Khan, Naeem, Asghari MD Bano, and Ali Babar. 2020.** "Impacts of plant growth promoters and plant growth regulators on rainfed agriculture." *PloS one* **15** (4):e0231426.

- Krishnasamy, V. 2003.** "Effect of Seed Soaking in Leaf Extract and Seed Coating with Leaf Powder on Seed Germination and Seedling Vigor in Sunflower." *Madras Agric. J* **91** (7-12):460-462.
- Kumaravelu, G, and D Kadamban. 2009.** "Panchagavya and its effect on the growth of the greengram cultivar K-851." *International Journal of Plant Sciences (Muzaffarnagar)* **4** (2):409-414.
- ANO HAR, K, Neelam Khare, and Hemant Kumar. 2017.** "Effects of leaf extract of *Lantana camara* on germination and growth behavior of selected tree species." *International Journal of Current Microbiology and Applied Sciences* **6** (7):2519-2526.
- MANOHAR, K, Neelam Khare, and Hemant Kumar. 2017.** "Effects of leaf extract of *Lantana camara* on germination and growth behavior of selected tree species." *International Journal of Current Microbiology and Applied Sciences* **6** (7):2519-2526.
- Moosavi, Amir, Reza Tavakkol AFSHARI, Abouzar Asadi, and Mohammad Hossain Gharineh. 2011.** "Allelopathic Effects of Aqueous Extract of Leaf Stem and Root of *Sorghum bicolor* on Seed Germination and Seedling Growth of *Vigna radiata L.*" *Notulae Scientia Biologicae* **3** (2):114-118.
- Moosavi, Amir, Reza Tavakkol AFSHARI, Abouzar Asadi, and Mohammad Hossain Gharineh. 2011.** "Allelopathic Effects of Aqueous Extract of Leaf Stem and Root of *Sorghum bicolor* on Seed Germination and Seedling Growth of *Vigna radiata L.*" *Notulae Scientia Biologicae* **3** (2):114-118.
- Muzzo, BI, AJ Mwilawa, DD Maleko, and EJ Mtengeti. 2018.** "Allelopathic effect of *Chromolaena odorata* aqueous leaf extracts on seed germination and seedling growth of selected crop and pasture species in Tanzania." *International Journal of Botany Studies* **3** (5):41-48.
- Perveen, Shagufta, Muhammad Naeem Mushtaq, Muhammad Yusu and Nighat Sarwar.**
- 2021.** "Allelopathic hormesis and potent allelochemicals from multipurpose tree *Moringa oleifera* leaf extract." *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology* **155** (1):154-158.
- Prakash, M, S Pallavamallan, G Sathiyarayanan, and S Rameshkumar. 2021.** "Effect of seed pelleting with botanicals on germination and seedling growth of cluster bean under induced saline condition." *Legume Research: An International Journal* **44** (1).
- Sasthri, G, and P Srimathi. 2010.** "Effect of organic and inorganic seed priming treatment on production of quality seed in cowpea." *Green Farming* **1** (4):366-368.

Sutar, R., Sujith, G. M., & Devakumar, N. (2019). Growth and yield of Cowpea [*Vigna unguiculata* (L.) Walp] as influenced by jeevamrutha and panchagavya application. *Legume Research-An International Journal*, **42**(6), 824-828.

Yashwanth Chandra, K., & Chaurasia, A. K. (2021). Effect of Seed Priming Treatments with Panchagavya, Jeevamrutha, Beejamrutha and Leaf Extracts, Sodium Molybdate on Seed Quality Parameters of Field Pea (*Pisum sativum*). *International Journal of Plant & Soil Science*, **33**(21), 24-33.