

**PHYTODIVERSITY STUDIES OF ANGIOSPERMIC AND GYMNOSPERMIC
TREES IN UMPHYRNAI PRIVATE FOREST OF EAST KHASI HILLS DISTRICT,
MEGHALAYA.**

Abstract

The study was carried out to identify and assess the variety of angiosperm and gymnosperm tree species in the study region in the East Khasi Hills District of Meghalaya during 2021-2022, located in Umphyrnai village at an elevation of 1578 m. A total of 187 unique trees from 17 different species have been identified. 4 gymnosperm species and 13 angiosperm species were discovered among them. The trees were discovered to belong to 9 different families. The most dominant gymnosperm species was discovered to be *Pinus kesiya*, and the most dominant angiosperm species was discovered to be *Alnus nepalensis*. *Pinus kesiya* has the highest IVI.

Key words: Angiosperms, gymnosperms, diversity, Umphyrnai, private forests.

Introduction

Positioned in the North Eastern part of the country, Meghalaya covers an area of 1,429 sq. km, which is 0.68 of the geographical area of the country. The State lies between 24 ° 58'N and 26 ° 07'N latitude and 89 ° 48' E to 92 ° 51' E longitude and is framed by Assam in the north and east and shares a transnational boundary with Bangladesh in the south and west. The State has three distinct regions, videlicet, Garo Hills, Khasi Hills, and Jaintia Hills.

The state, owing to the different ecological conditions similar to wide variation in downfall, temperature, altitude as well as soil conditions, supports luxuriant growth of different types of foliage, viz., tropical evergreen, tropical semi-evergreen, tropical wetish and dry deciduous, tropical broad-leaved hill timbers, tropical pine timbers, temperate timbers, and champignons (**Champion and Seth 1968; Rao and Hajra 1986**).

According to ISFR 2021, Meghalaya has 8389 square kilometers of unclassified wood. These sorts of forests are often tiny in size and are spread within the village boundaries. They are

Comment [OTU1]: Don't begin a sentence with numerics

handled and used in accordance with the owner's needs and desires. These forests are generally kept in order to produce wood. According to Tiwari, *B-K et al.* (2010), private forests in Meghalaya are the primary source of 76,870 m³ of timber valued at INR 284.5 million (USD 5.7 million) taken from Meghalaya's forests per year. Owners of poorly supplied private forests frequently transfer them to other land uses (for example, agriculture or charcoal burning).

Formatted: Font: Italic

Many private forests are secondary forests or pine (*Pinus kesiya*) plantations. In some cases, the owners have converted these forests into agricultural land, agroforests or home gardens. While collection of forest products by people other than the owners' family members is strictly prohibited, in few cases the owners allow fellow villagers to extract dead and fallen wood, and NTFPs for their personal use.

Methods and Materials:

Study Area

The research was conducted in private forest lands in Umphyrnai village, East Khasi Hills District, at a height of 1578m, during the years 2021-2022. The study area's geographic coordinates are 25.5359°N and 91.9590°E.

Fig 1: Location map of the Study Area.

Climate

The South-West monsoon, which originates in the Bay of Bengal, has a direct influence on the climate of Meghalaya. Depending on altitude and elevation, the climate differs from

eastern to western sections of the state. The area's climate is classified into four seasons: winter (December-February), spring (March-May), rainy (June-August), and autumn (September-November). The average maximum and lowest temperatures in June and July are 22°C and 8°C, respectively (January-February).

Methodology:

Standardized belt transects survey:

The forest was divided into four square sample plots, each measuring 20×20m. All of the species found in these plots were recorded. Each angiosperm or gymnosperm tree's botanical description was documented independently in each of these plots. All tree species were identified and recorded with the assistance of **informed individuals**. The phytodiversity of the study region was calculated by adding all of the species found in all of the plots. The diameter at breast height of trees in each plot was measured at 1.37 m.

Comment [OTU2]: The services of taxonomists should be employed always to ensure proper and accurate identification of the plant species

In the forest, vegetation data were quantitatively examined for density, frequency, and abundance (Curtis and McIntosh, 1950). The total of the relative values utilized for the Importance Value Index, namely frequency, density, and dominance (Curtis, 1959), The diameter at breast height (Dbh) of the tree was used to calculate the basal area, which was given in meters squared (m²). The Shannon-Weiner information index was used to calculate the diversity index (Shannon and Weiner, 1963). Simpson's Index was used to calculate dominance concentration (Simpson, 1949). Pielou's evenness index was used to calculate species' evenness (Pielou, 1969). Margalef's Index of Species Richness was used to calculate species richness (1958).

Quantitative analysis

In each forest community, field data was analysed for abundance, density, and frequency (Curtis and McIntosh, 1950). The total of the relative values utilised for the important value index, namely frequency, density, and dominance (Curtis, 1959), These are used to express a community's traits. Quantitative traits are analytical in nature and are typically expressed on a 5-point scale. These include characteristics such as frequency, density, quantity, cover, basal area, and so on.

Formatted: Indent: Left: 0"

1. Frequency

Formatted: Indent: Left: 0"

This term refers to the degree of dispersal of each species in a given area, which is usually stated as a percentage of occurrence. It will be examined by randomly sampling the research region and recording the names of the species that occur in each sampling unit. It is calculated by the following equation:

Formatted: Indent: Left: 0"

$$\text{Frequency (\%)} = \frac{\text{Number of quadrates in which the species occurred}}{\text{Total number of quadrates}} \times 100$$

1. Density

The numerical strength of a species within the community is represented by its density. The diversity of a species is defined by the number of individuals in any given unit area. The degree of competitiveness is indicated by density. It is called by the following formula

$$\text{Density} = \frac{\text{Total no. of individuals of a species in all quadrates}}{\text{Total no. of quadrates sampled}} \times 100$$

2. Abundance

This is the number of individuals per quadrate of occurrence of any species. It is calculated as follows:

$$\text{Abundance} = \frac{\text{Total no. of individuals of a species in all quadrates}}{\text{No. of quadrates in which the species occurred}} \times 100$$

3. Relative frequency

The proportion of individual species in an area in relation to the total number of species observed.

$$\text{Relative frequency} = \frac{\text{Number of quadrates in which the species occurs}}{\text{Total number of quadrates in which all the species occurred}} \times 100$$

4. Relative density

Relative density is the study of numerical strength of a species of a species in relation to the total number of individuals of all the species and can be calculated as:

$$\text{Relative density} = \frac{\text{Total no. of individual of a particular species in all quadrates}}{\text{Total no. of individuals of all species in all quadrates}} \times 100$$

5. Relative Dominance

The value of a species' entire basal cover determines its dominance. The coverage value of a species in relation to the total coverage of the other species in the area is known as relative dominance.

$$\text{Relative Dominance} = \frac{\text{Total basal area of a particular species}}{\text{Total basal area of all species}} \times 100$$

6. Basal Area

Basal Area is one of the primary determinants of community dominance. The girth of the tree stems at breast height (DBH) at 1.37m above ground level will be used to calculate the basal area.

$$\text{Basal Area (m}^2\text{)} = \frac{\pi \times (\text{DBH})^2}{4 \times 10000}$$

7. Importance Value Index (IVI)

This index is used to determine each species' overall relevance in the community structure. In order to calculate this index, the percentage values of relative frequency, relative density, and relative dominance are added together, and the resulting value is known as the Importance Value Index (IVI) of the species (Curtis, 1959).

$$\text{IVI} = \text{Relative Frequency} + \text{Relative Density} + \text{Relative Dominance}$$

8. Species richness

'Margalef's index of richness (Dmg) (Magurran, 1988)

$$\text{Dmg} = (S-1 / \ln N)$$

Where, S = Total number of species, N = Number of individuals.

9. Species diversity

Shannon and Weiner (1963)

$$H' = - \sum (P_i \ln P_i)$$

Where, $P_i = n/N$ (proportion of each species in the sample)

n = Number of individual species

N = Total number of individuals

10. Evenness Index

(Pielou, 1975)

$$E = H' / \ln S$$

Where, H' = Shannon Index Value

\ln = Bits per individual

11. Index of dominance (D)

Simpson (1949)

$$D = \sum (n/N)^2$$

Where, D = Simpson index of dominance

n = Number of individual species

N = Total number of individuals

Results and Discussion:

The vegetation composition observed in 4 quadrates of 20×20 m size at random locations revealed a total of 187 unique trees representing 17 species. Four species of gymnosperms and thirteen species of angiosperms were discovered. The trees were discovered to be members of nine separate families (Table 1). Table 1 lists the species, along with their scientific names, local/common names, and families.

Table 1: Taxonomic status of trees at Umphyrnai Private Forest of East Khasi Hills District.

Sl. No.	Scientific name	Local/Common Name	Family	No. of Individuals
1	<i>Pinus kesiya</i>	Dieng Kseh Khasi	Pinaceae	58
2	<i>Pinus roxburghii</i>	Dieng Kseh Bilat	Pinaceae	19

Comment [OTU3]: Write the Author Citation for all the species

3	<i>Cryptomeria japonica</i>	Dieng Cedar	Cupressaceae	8
4	<i>Juniperous japonica</i>	Juniper	Cupressaceae	5
5	<i>Alnus nepalensis</i>	Dieng iong	Betulaceae	23
6	<i>Alnus glutinosa</i>	Dieng lieh	Betulaceae	10
7	<i>Myrica esculenta</i>	Dieng Sohphie Heh	Myricaceae	4
8	<i>Rhus chinensis</i>	Dieng Sohma	Anacardiaceae	15
9	<i>Schima khasiana</i>	Dieng ngan		5
10	<i>Pyrus pashia</i>	Dieng Sohjhur	Rosaceae	6
11	<i>Prunus cerasoides</i>	Dieng Cherry (Jew)	Rosaceae	7
12	<i>Prunus nepalensis</i>	Dieng Cherry (Thiang)	Rosaceae	11
13	<i>Cinnamomum cecicodahne</i>	Dieng Pingwait	Lauraceae	7
14	<i>Quercus serrata</i>	Jolcham Oak	Fagaceae	1
15	<i>Acacia daelbata</i>	Dieng Baibl	Fabaceae	3
16	<i>Celtis tetrandia</i>	Nilgiri Elm	Ulmaceae	3
17	<i>Pyrus calleryana</i>	Dieng Sohphoh	Rosaceae	1
18	<i>Pourthia arguta</i>	Sohryngkham	Rosaceae	1

Gymnosperms

Table 1 indicates that from the occurrence of 90 gymnospermic trees, 4 species belonging to *Pinaceae* and *Cupressaceae* family were found. The individual gymnosperms with the maximum number of individuals includes *Pinus kesiya* (58), followed by *Pinus roxburghii* (19). The gymnosperms with the least number of occurrences include *Cryptomeria japonica* (8) and *Juniperous phoenica* (5).

Angiosperms

Table 1 shows that from the occurrence of 97 angiospermic trees, 14 species belonging to 7 families' viz., *Betulaceae*, *Myricaceae*, *Rosaceae*, *Fabaceae*, *Fagaceae*, *Anacardiaceae*, and *Ulmaceae* were recorded. The species with the highest number of occurrence belongs to *Alnus nepalensis* (23), and followed by *Rhus chinensis* (15). Whereas the individuals with the least number of occurrence includes *Quercus serrata*, *Pyrus calleryana* and *Pourthia arguta* each with one individual only.

Sl no.	Name of species	No. of species in each quadrat				Total no. of individuals	Total no. of species occurred	Total no. of quadrats studied	Basal Area (m ²)	Frequency	Density	Abundance	Relative frequency	Relative density	Relative dominance	IVI
		I	II	III	IV											
1.	<i>Pinus kesiya</i>	11	14	11	22	58	4	4	4.045	100	14.5	14.5	7.547	31.01	36.08	74.643
2.	<i>Pinus roxburghii</i>	4	5	4	6	19	4	4	1.219	100	4.75	4.75	7.547	10.16	10.87	28.577
3.	<i>Cryptomeria japonica</i>	2	2	2	2	8	4	4	0.498	100	2	2	7.547	4.27	4.45	16.275
4.	<i>Juniperous phoenica</i>	1	0	4	0	5	2	4	0.318	50	1.25	2.5	3.773	2.67	2.84	9.287
5.	<i>Alnus nepalensis</i>	5	7	5	6	23	4	4	1.51	100	5.75	5.75	7.547	12.29	13.5	33.346
6.	<i>Alnus glutinosa</i>	2	5	2	1	10	4	4	0.644	100	2.5	2.5	7.547	5.34	5.76	18.654
7.	<i>Myrica esculenta</i>	1	0	1	2	4	3	4	0.193	75	1	1.4	5.66	2.13	1.72	9.519
8.	<i>Rhus chinensis</i>	4	3	4	4	15	4	4	0.439	100	3.75	3.75	7.547	8.021	3.92	19.488
9.	<i>Schima khasiana</i>	1	3	1	0	5	3	4	0.419	75	1.25	1.7	5.66	2.67	3.74	12.074
10.	<i>Pyrus pashia</i>	1	2	0	3	6	3	4	0.294	75	1.5	2	5.66	3.208	2.63	11.498
11.	<i>Prunus cerasoides</i>	1	3	2	1	7	4	4	0.262	100	1.75	1.75	7.547	3.74	2.34	13.630
12.	<i>Prunus nepalensis</i>	3	5	2	1	11	4	4	0.678	100	2.75	2.75	7.547	5.882	6.06	19.489
13.	<i>Cinnamomum cecicodahne</i>	1	2	4	0	7	3	4	0.329	75	1.75	2.4	5.66	3.743	2.94	12.343
14.	<i>Quercus serrata</i>	1	0	0	0	1	1	4	0.056	25	0.25	1	1.886	0.534	0.5	2.921
15.	<i>Acacia daelbata</i>	0	1	0	2	3	2	4	0.0143	25	0.75	1.5	3.773	1.604	1.28	6.657
16.	<i>Celtis tetrandia</i>	0	0	2	1	3	2	4	0.099	50	0.75	1.5	3.773	1.604	0.89	6.267
17.	<i>Pyrus calleryana</i>	0	0	0	1	1	1	4	0.035	25	0.25	1	1.886	0.534	0.32	2.741
18.	<i>Pourthia arqueta</i>	0	0	1	0	1	1	4	0.022	25	0.25	1	1.886	0.534	0.2	2.621
						187	53		11.21							300.03

Table 2: Quantitative analysis of tree species at Umphyrnai Private Forest.




Fig 2: Importance Value Index at Umphyrnai Private Forest.

Diversity aspects

Various aspects were calculated on the basis of Shannon and Weiner index of species diversity (H'), Margalef's index of richness (D_{mg}), Pielou index of evenness (E), Simpson index of dominance (D).

Diversity Parameters in Umphyrnai Forest Stand

Table 3 shows that Shannon Weiner diversity index in the study site is greater in Angiosperms (2.30) than that of gymnosperms (0.99) which indicate that the angiosperms are more diverse.

Based on diversity indices values from table 3, it was concluded that tree species diversity is greater in Angiosperms (2.30) than that of gymnosperms (0.99). Simpson index has been shown to be higher in gymnosperms (0.459) and lower in angiosperms (0.114). Margalef's Species richness is found to be higher in Angiosperms (2.841) and lower in Gymnosperms (0.666). As per dominance index, it was found that Angiosperms (0.885) are found to be higher than that of gymnosperms (0.540).

Table 3: Gymnospermic and Angiospermic tree diversity in Umphyrnai Forest Stand

Sl. No.	Attributes/Parameters	Gymnosperms	Angiosperms
1.	Shannon-Weiner Diversity	0.990	2.30
2.	Simpson index	0.459	0.114
3.	Evenness/Equitability Index	0.712	0.872
4.	Margalef's Richness Index	0.666	2.841
5	Dominance index	0.540	0.885
6.	Total no. of individuals	90	97

Conclusion

The results of present study could be concluded as an analysis of angiospermic and gymnospermic tree diversity in the private forest of Umphyrnai village. It was observed that in Umphyrnai Forest Stand 4 species of gymnosperms and 14 species of angiosperms were recorded. The maximum frequency, density, abundance, and IVI of gymnosperms were recorded for *Pinus kesiya* and that of angiosperms was recorded for *Alnus nepalensis*. Shannon Weiner diversity index in the study site is greater in Angiosperms (2.30) than that of gymnosperms (0.99) which indicate that the angiosperms are more diverse.

References:

- **Ahmed, J., and Sharma, S. (2014).** "Spatial Pattern, Diversity, and Phytosociological Analysis of Woody Plant Species in Ponda Watershed, Rajouri, J&K, India", *International Journal of Current Research*,- ~~Vol-6, Issue-(06), pp-: 7022-7027.~~
- **Antony, I., Jose, S.K., and Madhu, G. (2015).** "Phytodiversity Assessment in Neyyar Wildlife Sanctuary Western Ghats, India", *International Journal of Current Research*, ~~Vol-7, Issue (6), pp- 16730-16732.~~

- **Ayyappan, N., and Parthasarathy, N. (1999).** Biodiversity inventory of trees in a large scale permanent plot of tropical evergreen forest at Varagalaiar, Anamalais, Western Ghats, India. *Biodiversity and Conservation*, 8: 1533-1554.
- **Bahugunal, A., Phondani, P.C., Vikram, S., Negi, ~~et.al.~~(2010).** Floristic Diversity and Indigenous Uses of Forest Vegetation of Dabka Watershed in Indian Central Himalaya, *Ethnobotanical Leaflets*, 14: 491-510.
- **Balakrishna, N.P. (1981-1983).** Flora of Jowai and Vicinity. 2 Vols., Botanical Survey of India, Howrah. 666 pages.
- **Bhuyan, P.,Khan, M.L. ~~and,~~ Tripathi, R.S. (2003).** Tree diversity and population structure in undisturbed and human-impacted stands of tropical wet evergreen forest in Arunachal Pradesh, Eastern Himalayas, India. *Biodiversity and Conservation*, 12 (8): 1753-1773.
- **Chaturvedi, S.K., and Jamir, M. (2015).** A checklist of angiospermic flora of Zunheboto district of Nagaland, India. *Pleione*, Vol. 9(1): 82 - 94.
- **Chiarucci, A., and Bonini, I. (2005).** Quantitative floristic as a tool for the assessment of diversity in Tuscan forest. *Forest Ecology and Management*. 242: 160-170.
- **Deb, P. and Sundriyal, R.C. (2011).** Vegetation dynamics of an old growth lowland tropical rainforest in Northeast India: Species composition and stand heterogeneity. *International Journal of Biodiversity and Conservation*. 12 (2): 104-112.
- **Dutta, G., and Devi, A. (2013).** Plant Diversity, Population and regeneration status in undisturbed tropical forests in Assam, Northeast, India. *Journal of Forestry Research*, 24: 715-720.
- **Dhyani, S., Mir, M.A., and Kumar, N. (2014).** Angiospermic diversity of Karwapani fresh water swamp forest in Doon valley, Uttaranchal. Environmental Science: *An Indian Journal Current Research Paper*. ~~Vol. 9, Issue (4); pp. 126-131.~~
- **Jamir, S.A., and Pandey, H.n. (2003).** Vascular plant diversity in the sacred grooves of Jaintia Hills in Northeast, India. *Biodiversity and Conservation*, 12, ~~pp.~~ 1457-1510.

Comment [OTU4]: Remove and write names of all authors

Formatted: Font: Italic

- **Kharkongor, B.M., and Tiwari, B.K. (2017).** Sacred Groves of Meghalaya: A Review. *International Journal of Science and Research (IJSR)*. ~~Vol. 6, Issue (3): pp.~~ 346-349.
- **Ovung, E.Y., Wapongnungsang, and Tripathi, S.K. (2021).** Assessment of tree diversity in tropical moist deciduous forest of Mizoram, University, Northeast, India. *Journal of Applied and Natural Sciences*, Vol. 13, Issue 1: 2436.
- **Sarkar, M., and Devi, A. (2014).** Assessment of diversity, population structure, and regeneration status of tree species in Hollongapar Gibbon Wildlife Sanctuary, Assam, Northeast, India. *Tropical Plant Research: An International Journal*, ~~Vol.~~ 1(2): 26-36.
- **Shannon, C.E., and Weiner, W. (1963).** The mathematical theory of communities. University of Illinois Press, Urbana Illinois.
- **Simpson, E.H. (1949).** Measurement of diversity: Nature ~ London) 163-688.

Comment [OTU5]: No initials

Comment [OTU6]: Incomplete reference