

**Effect of different organic manures on growth, yield and
Quality of cucumber (*Cucumis sativus* L.) c.v. (Harsh)
under
Prayagraj agro climate condition**

ABSTRACT

A field experiment ~~on entitled “the E~~ffect of different organic manures on growth, yield and quality of cucumber (*Cucumis sativus* L.) c.v. (Harsh) under Prayagraj agro climate condition² was conducted in a Randomized Block Design with three replications during the *Summer* season, 2021 at Horticultural Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj with a view to determine the influence effect of organic manures application on cucumber ~~variety Harsh for its~~ growth, quality and yield, ~~and to work out the economics of various treatments. Under this experiment, overall 10Ten~~ treatment combinations ~~were taken~~ viz. T₀ (Control), T₁ (FYM 75% + Poultry manure 25%), T₂ (FYM 75% + Vermicompost 25%), T₃ (FYM 25% + Poultry manure 75%), T₄ (Poultry manure 75% + Vermicompost 25%), T₅ (Vermicompost 75% + FYM 25%), T₆ (Vermicompost 25% + Poultry manure 25%), T₇ (FYM 50% + Poultry manure 50%), T₈ (Vermicompost 50% + Poultry manure 50%) and T₉ (FYM 50% + Vermicompost 50%). From the present investigation it was concluded that T₈ (Poultry manure 50% + Vermicompost 50%) was found to be best among all treatment in terms of growth, yield and quality of cucumber ~~i.e. T.S.S.~~ The highest net return was found from T₈ (~~PoultryPoultry~~ manure 50% + vermicompost 50%) with ₹ 1,56,775 and the B:C ratio ~~wasthat is~~ 1.92.

Keywords: FYM, Poultry manure, Vermicompost, ~~T.S.S.~~

INTRODUCTION

Cucumber (~~*Cucumis sativus*~~) is a widely-cultivated creeping vine plant ~~in the~~ *Cucurbitaceae* family that bears usually cylindrical fruits ~~that,~~ which are used as vegetables. Considered an annual plant, there are three main varieties of cucumber — slicing, pickling, and burpless/seedless, — within which several cultivars have been ~~evolved~~ created. The cucumber originated ~~in s from~~ South Asia, but now grown ~~in many countries s on most~~ continents and also, as many different types of cucumber are traded on the global market.

Botanically ~~C~~cucumber is known as *Cucumis Sativus* L. belongs to family *Cucurbitaceae*. It is a diploid self-pollinated species with chromosome number $2n=2x=14$ (Karpechenko, 1925). Cucumber probably originated from India (De Candolle, 1967). It is mainly cultivated in China, India, Turkey, Iran and other parts of south-east Asia. Progenitor of cucumber is "~~Cucumis hardwickii~~". Economic sex ratio is 15:1.

The area under ~~c~~Cucumber production in India ~~was accounts to~~ 107 million ha ~~withwith a~~ production of 1658 million tonnes ~~during in year~~ 2018-19. (Source: NHB, Ministry of Agriculture

& Farmers Welfare, Government of India, 2019-20). Haryana ranks first ~~in~~ area and production of ~~C~~cucumber in year 2018-19 followed by Madhya Pradesh and Karnataka. The production of ~~c~~Cucumber in Uttar Pradesh is 81.47 million tonnes for year 2018-19. ~~It The cucumber~~ is used as salad, pickles and also as ~~a~~ cooked vegetable. It has many uses in ayurvedic medicines. According to 'Unani' medicines, the ~~seed~~ oil ~~from its seed~~ is ~~used god~~ for the brain and the body ailments. Cucumber ~~contains~~ has 96.3 g water, magnesium 11 mg, sodium 10.2 mg, Vitamin C 7 mg, 2.5g Carbohydrates, Oxalic acid 15 mg, Calcium 10 mg, Sulphur 17 mg, Potassium 50 mg and many other nutrients out of 100 g of edible portion. (Choudhary and Basu, 1967).

Cucumber is well adapted crop for warm-season crop and it does not withstand even light frost. The crop performs well in temperature range between 18°C ~~and~~ - 24°C and soil having pH ranging between 5.5-6.7 ~~irrespective~~ of its kind from sandy to heavy clay soil. It is grown as sole crop in India in ~~Zaid and~~ Zaid season. It is well suited to hot and warm climate with annual rainfall of 60-75 cm. However, cucumber cannot withstand water lodging. It is a short duration cucurbit crop, ~~grown in all parts of India,~~ well adapted ~~to crop~~ for ~~many~~ diversified cropping systems. It is considered as quality dietary food due to its excellent digestibility and rich water content (96.3 g/100 g). Cucumber is a dependable laxative for those who suffer constipation. The juice of cucumber is a valuable food in the treatment of hyper acidity, gastric and duodenal ulcers.

Formatted: Indent: First line: 0.5", Right: 0.05", Space Before: 0 pt, Line spacing: Multiple 1.5 li

Formatted: Font: Italic

Farmyard manure (~~or~~ FYM) refers to the decomposed mixture of dung and urine of farm animals along with litter and left over material from roughages or fodder fed to the cattle. On an average well decomposed ~~FYM farmyard manure~~ contains 0.5 per cent N, 0.2 per cent P₂O₅ and .0.5 per cent K₂O. ~~Urine, which is wasted, contains one per cent nitrogen and 1.35 per cent potassium. The excreta of birds ferment very quickly.~~ If left exposed, 50 percent of its nitrogen is lost within

30 days. Poultry manure contains higher nitrogen and phosphorus compared to other bulky organic manures. The average nutrient content is 3.03 per cent N; 2.63 per cent P₂O₅ and 1.4 per cent K₂O. Vermicompost is the product of the decomposition ~~of farm and other wastes process~~ using various species of worms, usually ~~red wigglers, white worms, and other~~ earthworms, ~~to create a mixture of decomposing vegetable or food waste, bedding materials, and vermicast. This process is called vermicomposting, while the rearing of worms for this purpose is called vermiculture.~~

Intensive use of only chemical fertilizers to achieve high production has created various problems. Continuous application of heavy doses of chemical fertilizers without organic manures has led to deterioration of soil health in terms of physical and chemical properties of soil, decrease in soil microbial activities, and also reduction in soil humus (Anjanappa *et al.*, 2011). Intensive cultivation and improper fertilizer use leads to deficiency of nutrients in soil, thus resulting in lower yield of crops. Therefore, the use of farm input in the form of organic manure has become necessary. Number of investigations were undertaken to study the ~~effect~~ effect of different chemical fertilizers on cucumber in different soils; on the other hand, the use of organic nutrient sources such as FYM, poultry manures, vermicompost *etc.* remains as alternative choice for the production of residue free wholesome produce, and to maintain soil health. ~~However, The~~ information on holistic approach with suitable combination of organic manure and inorganic fertilizers on growth and yield and quality of cucumber in soil of eastern region of Uttar Pradesh is very meager.

Keeping in view the above facts, the present investigation was with an objective of studying the undertaken entitled, 'Effect of different organic manures on growth, yield and quality of cucumber (*Cucumis sativus* L.) c.v. (Harsh) under Prayagraj agro climate condition².

Formatted: Right: 0.05", Space Before: 1.45 pt

Formatted: Justified, Indent: Left: 1", Right: 0.05", Space Before: 1.45 pt, Line spacing: Multiple 1.49 li

MATERIAL AND METHODS

The experiment was conducted in Randomized Block Design comprising 10 treatments with three replications during *Summer 2021-22* at Horticultural Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj. The unit plot size was 2.25 m². The plants were planted with a spacing of 1.5 m between the rows and 90 cm between the plants. There were ten plants in each plot. The observations were recorded on randomly selected and tagged five plants for on growth characters. In general, soil properties of experimental site showed a typical alluvial soil of eastern region of Uttar Pradesh. ~~Soil was sandy loam in texture, slightly acidic in reaction and having low electrical conductivity, very high in organic carbon, low in available nitrogen, low in available phosphorus and moderately high in available potassium.~~

Include the Soil chemical properties – Available N, P,K, Organic carbon values. Soil pH and EC

IT IS A MUST.

Weeding and plant protection measure were followed as and when needed. Observations were recorded at different stages of growth periods. The data were statistically analyzed by the method suggested by **Fisher and Yates,**

1936.

The treatments included. T₀(Control), T₁ (FYM 75% + Poultry manure 25%), T₂ (FYM 75% + Vermicompost 25%), T₃ (FYM 25% + Poultry manure 75%), T₄ (Poultry manure 75% + Vermicompost 25%), T₅ (Vermicompost 75% + FYM 25%), T₆ (Vermicompost 25% + Poultry manure 25%), T₇ (FYM 50% + Poultry manure 50%), T₈ (Vermicompost 50% + Poultry manure 50%) and T₉ (FYM 50% + Vermicompost 50%). From the present investigation it was concluded that T₈ (Poultry manure 50% + Vermicompost 50%)

RESULTS AND DISCUSSION

The findings of the In the present investigation ~~an attempt has been made~~ to study the effect of different organic treatment combinations on growth and yield of cucumber (*Cucumis sativus* L.) in Prayagraj climatic zone are. ~~The results obtained are~~ presented herein relevant Tables as follows

A) Growth Parameters

Growth parameters like comprised of percentage of seed germination, vine length (30, 45,

Formatted: Indent: Left: 1.5", Right: 0.05", Space Before: 1.45 pt, Line spacing: Multiple 1.49 li

Formatted: Font: 12 pt

Formatted: Justified, Indent: Left: 1.5", Right: 0.06", Space Before: 0 pt, Line spacing: single

60 DAS), number of branches per plant, days to first emergence of male and female flower and node at first male and female flowers. ~~The data for these characters observed~~ have been listed out in table 1.1, 1.2 and 1.3.

~~1.~~ *Percentage of seed germination*

The maximum ~~value of percent of~~ seed germination was observed ~~in for~~ T₄ (Poultry manure 75% + Vermicompost 25%) with value ~~of~~ 91.67% followed by T₈ (Poultry manure 50% + vermicompost 50%) with value of 95.83%. ~~The M~~minimum value ~~for of percent of~~ seed germination was observed ~~in for~~ T₆ (Vermicompost 25% + Poultry manure 25%) with value of 66.67%.

Formatted: Font: Italic

~~2.~~ *Vine length (cm) [30, 45 and 60 DAS]*

Formatted: Font: Italic

The maximum ~~V~~yine length (29.28 cm) at 30 DAS was observed with treatment T₈ (Poultry manure 50% + Vermicompost 50%) followed by T₆ (Vermicompost 25% + Poultry manure 25%) with 29.02 cm. ~~A~~~~M~~minimum ~~V~~yine length (21.09 cm) was observed in T₀ (control), while the remaining treatments were moderate in their growth habit. ~~At 45 DAS,~~ ~~T~~the maximum ~~V~~yine length (101.96 cm) at 45 DAS was observed with treatment T₃ (FYM 25% + Poultry manure 75%) followed by T₁ (Vermicompost 75% + FYM 25%) with 101.96 cm. Minimum ~~y~~~~V~~yine length (94.99 cm) was observed in T₅ (Vermicompost 75% + FYM 25%), while the remaining treatments were moderate in their growth habit. The maximum ~~V~~yine length (162.99 cm) at 60 DAS was observed in treatment T₈ (Poultry manure 50% + Vermicompost 50%) followed by T₆ (Vermicompost 25% + Poultry manure 25%) with 162.19 cm. Minimum ~~y~~~~V~~yine length (155.17 cm) was observed in T₀ (control), while the remaining treatments were moderate in their growth habit. The application of organic manures might have improved the soil physical and chemical properties and leading to the adequate supply of nutrients to the plants which might have promoted the maximum vegetative growth while the minimum plant growth was due to non-availability of nutrients. Similar findings were reported by **Ikeh (2012); Hamdi et al. (2017); Ghayal et al. (2018) and Tahir et al. (2019)** in cucumber.

3.—Days to first male and female flower emergence

Among the application of organic manure, the minimum days to first male flower emergence was seen in T₇ (FYM 50% + Poultry manure 50%) with in 31.67 days, followed by T₈ (Poultry manure 50% + Vermicompost 50%) with 33.33 days, whereas maximum days to first male flower emergence took 35.67 days ~~was~~ recorded in T₅ (Vermicompost 75% + FYM 25%). Among the application of organic manure, the minimum days to first female flower emergence was seen in T₇ (FYM 50% + Poultry manure 50%) with 40.00 days, followed by T₈ (Poultry manure 50% + Vermicompost 50%) with 41.67 days whereas maximum days to first female flower emergence was 44.00 days, ~~was~~ recorded in T₅ (Vermicompost 75% + FYM 25%). Integration of FYM and poultry manure favoured vigorous growth and synthesized more hormones in plants, which might have helped in the translocation as well as more quantity of available phosphorus through the xylem vessels and their accumulation in the axillary buds that would have favoured the plant to enter into reproductive phase. Similar results have also been reported by **Kaur and Kaur (2018); Tahir et al. (2019); Jagraj et al. (2020) and Singh et al. (2021)** in cucumber.

Formatted: Font: Italic

4. Node at first male and female flower emergence

Formatted: Font: Italic

Among the application of organic manure, the maximum node at first male flower emergence was seen in T₅ (Vermicompost 75% + FYM 25%) at 4.11 node, followed by T₄ (Poultry manure 75% + Vermicompost 25%) ~~at with~~ 3.89 node, whereas minimum node at first male flower emergence ~~at~~ 3.44 node was recorded in T₄ (FYM 75% + Vermicompost

25%). Among the application of organic manure, the maximum node at first female flower emergence was seen in T₅ (Vermicompost 75% + FYM 25%) at 6.78 node, followed by T₈ (Poultry manure 50% + Vermicompost 50%) ~~with-at~~ 6.67 node, whereas minimum node at first female flower emergence ~~at~~ 6.00 node was recorded in T₄ (FYM 75% + Vermicompost 25%).

5. Number of branches per plant

Formatted: Font: Italic

It was also found that T₈ (Poultry manure 50% + ~~Vermiempost~~Vermicompost 50%) ~~had with~~ maximum ~~value i.e.~~ (5.56) branches per plant followed by T₆ (Vermicompost 25% + Poultry manure 25%) with 4.89 branches per plant, whereas the minimum score was observed in treatment T₂ (FYM 75% + Vermicompost 25%) with 4.22 ~~branches~~. It was noticed that number of branches per plant ~~increased~~ with increasing vine length successively with the increasing levels of nutrient. Combination of organic manures also recorded maximum vine length and number of branches also which helped the plants in better photosynthesis to attain vigour. The findings of the present investigation are in conformity with the reports of Ikeh *et al.* (2012); Hamdi *et al.* (2017); Ghayal *et al.* (2018); Tahir *et al.* (2019) and Singh *et al.* (2021) in cucumber.

Table 1.1 Effect of organic treatment combination on seed germination, days to first male and female flower emergence

Treatment Notation	Treatment Combination	Seed Germination (%)	Days to first male flower emergence	Days to first female flower emergence
T0	Control	79.17	34.00	42.33
T1	FYM (75%) + Poultry manure (25%)	70.83	34.67	43.00
T2	FYM (75%) + Vermicompost (25%)	79.17	34.33	42.67
T3	Poultry manure (75%) + FYM (25%)	72.92	35.00	43.33
T4	Poultry manure (75%) Vermicompost (25%)	91.67	33.33	41.67
T5	Vermicompost (75%) + FYM (25%)	75.00	35.67	44.00
T6	Vermicompost (25%) + Poultry manure (25%)	66.67	33.67	42.00
T7	FYM (50%) + Poultry manure (50%)	75.00	31.67	40.00
T8	Poultry manure (50%) + vermicompost (50%)	95.83	33.33	41.67
T9	Vermicompost (50%) + FYM (50%)	79.17	34.33	42.67
	Mean	78.54	34.00	42.33
	'F' Test	S	S	S
	C.V. <u>Percent or not check it</u>	0.29	0.21	0.48
	S.E. (m)	0.18	0.05	0.17
	C.D. at 5%	0.39	0.12	0.37

Formatted: Font color: Red

Table 1.2 Effect of organic manure treatment combinations on vine length (cm) at 30, 45 and 60 DAS.

Treatment Notation	Treatment Combination	30 DAS	45 DAS	60 DAS
T0	Control	21.99	97.09	155.17
T1	FYM (75%) + Poultry manure (25%)	26.48	101.96	160.41
T2	FYM (75%) + Vermicompost (25%)	23.02	98.16	156.34
T3	Poultry manure (75%) + FYM (25%)	27.88	103.07	161.44
T4	Poultry manure (75%) Vermicompost (25%)	26.22	101.77	159.53
T5	Vermicompost (75%) + FYM (25%)	25.09	94.99	158.22
T6	Vermicompost (25%) + Poultry manure (25%)	29.02	98.80	162.19

T7	FYM (50%) + Poultry manure (50%)	28.76	98.53	161.77
T8	Poultry manure (50%) + vermicompost (50%)	29.28	99.54	162.99
T9	Vermicompost (50%) + FYM (50%)	25.00	95.13	158.40
	Mean	26.27	98.90	159.64
	'F' Test	S	S	S
	C.V. Percent or not check it	0.24	0.11	0.08
	S.E. (d)	0.05	0.09	0.10
	THE CD VALUES ARE TOTALLY WRONG. KINDLY REDO ANALYSIS			
	C.D. at 5%	2.43	2.46	2.48

Formatted Table

Formatted: Font color: Red

Table 1.3 Effect of organic treatment combinations on node at first male and female flower and number of branches per plant.

Treatment Notation	Treatment Combination	Node at first male flower	Node at first female flower	Number of branches per plant
T0	Control	3.78	6.67	4.22
T1	FYM (75%) + Poultry manure (25%)	3.89	6.33	4.56
T2	FYM (75%) + Vermicompost (25%)	3.44	6.00	4.22
T3	Poultry manure (75%) + FYM (25%)	3.78	6.33	4.33
T4	Poultry manure (75%) Vermicompost (25%)	3.89	6.22	4.44
T5	Vermicompost (75%) + FYM (25%)	4.11	6.78	5.33
T6	Vermicompost (25%) + Poultry manure (25%)	3.56	6.22	4.89
T7	FYM (50%) + Poultry manure (50%)	3.67	6.56	4.44
T8	Poultry manure (50%) + vermicompost (50%)	3.44	6.67	5.56
T9	Vermicompost (50%) + FYM (50%)	3.56	6.11	4.56
	Mean	3.71	6.38	4.65
	'F' Test	S	S	S
	C.V. Percent or not check it	1.86	0.39	0.17
	S.E. (m)	0.05	0.02	0.03
	C.D. at 5%	0.11	0.04	0.01

B) Yield Parameters

Yield parameters comprised of number of fruits per plant, fruit length, fruit diameter, fruit yield per plant, fruit yield per plot, fruit yield per hectare and T.S.S. The data for these characters observed have been listed out in table 1.4.,

1. Number of fruits per plant

The maximum number of fruits per plants (26.56) was recorded in treatment T₈ (Poultry manure 50% + Vermicompost 50%) followed by T₉ (Vermicompost 50% + FYM 50%) i.e., 23.33 and the lowest fruits per plant (15.78) was observed in T₁ (FYM 75% + Poultry manure 25%). Maximum number of fruits per plant increase in T₈ (Poultry manure 50% + Vermicompost 50%) might be due to increased number of flowers which might have formed into fruits due to adequate availability of major and minor nutrients during its growth and development. Similar conclusions have also been reported by Ikeh *et al.* (2012); Hamdi *et al.* (2017); Singh *et al.* (2017); Ghayal *et al.* (2018); Tahir *et al.* (2019) and Singh *et al.* (2021) in cucumber.

2. Average fruit yield per plant, per plot and per hectare.

The maximum fruit yield per plant (6.12 kg/plant) was recorded in treatment T₈ (Poultry manure 50% + Vermicompost 50%) followed by T₇ (FYM 50% + Poultry manure 50%) i.e., 5.51 kg/plant and the lowest fruit yield per plant (3.69 kg/plant) was observed in T₅ (Vermicompost 75% + FYM 25%). The maximum fruit yield per plot 39.77 kg/plot were recorded in treatment T₈ (Poultry manure 50% + Vermicompost 50%) followed by T₇ (FYM 50% + Poultry manure 50%) i.e., 35.65 kg/plot and the lowest fruit yield per plot (23.77 kg/plot) was observed in T₅ (Vermicompost 75% + FYM 25%). The maximum fruit yield per hectare (15.90 t/ha) was recorded in treatment T₈ (Poultry manure 50% + Vermicompost 50%) followed by T₇ (FYM 50% + Poultry manure 50%) i.e., 14.31 t/ha and the lowest fruit yield per hectare (10.11 t/ha) was observed in T₅ (Vermicompost 75% + FYM 25%). Organic manures play an important role in improving productivity and quality of fruit. Doses of Poultry manure, FYM and vermicompost increased the vigour of plants, assimilating area, size of fruit, thereby resulting into higher weight of fruit. These results were in close conformity with the findings of Ikeh *et al.* (2012); Hamdi *et al.* (2017); Singh *et al.* (2017); Ghayal *et al.* (2018); Tahir *et al.* (2019) and Singh *et al.* (2021) in cucumber.

Formatted: Font: Italic

Formatted: Font: Italic

3. Average fruit length and diameter

Formatted: Font: Italic

The maximum fruit length of 18.34 cm was recorded in treatment T₈ (Poultry manure 50% + Vermicompost 50%) followed by T₉ (Vermicompost 50% + FYM 50%) i.e., 17.94 cm and the lowest fruit length (16.42 cm) were observed in T₅ (Vermicompost 75% + FYM 25%). The maximum fruit diameter of 7.29 cm was recorded in treatment T₀ (control) followed by T₈ (Poultry manure 50% + Vermicompost 50%) i.e., 6.80 cm and the lowest fruit diameter (5.67 cm) was observed in T₇ (FYM 50% + Poultry manure 50%). These results are in close conformity with the findings of Ikeh *et al.* (2012); Hamdi *et al.* (2017); Singh *et al.* (2017); Ghayal *et al.* (2018); Kaur and Kaur (2018); Tahir *et al.* (2019) and Singh *et al.* (2021) in cucumber.

4. T.S.S. EXPAND

Formatted: Font: Italic

The maximum T.S.S. (5.77 °Brix) was observed in treatment T₈ (Poultry manure 50% + vermicompost 50%) followed by T₅ (Vermicompost 75% + FYM 25%) with 5.70 °Brix. The minimum T.S.S. (5.15 °Brix) was noticed in treatment T₄ (Poultry manure 75% + Vermicompost 25%). These results are in close conformity with the findings of Ikeh *et al.* (2012); Hamdi *et al.* (2017); Singh *et al.* (2017); Ghayal *et al.* (2018); Kaur and Kaur (2018); Tahir *et al.* (2019); Jagraj *et al.* (2020); Naseer *et al.* (2020) and Singh *et al.* (2021) in cucumber.

Table 1.4 Effect of organic treatment combinations on number of fruits per plant, fruit length, fruit diameter and T.S.S.

Treatment Notation	Treatment Combination	Number of fruits per plant	Fruit diameter (cm)	Fruit Length (cm)	T.S.S.
T0	Control	18.11	7.29	17.67	5.42
T1	FYM (75%) + Poultry manure (25%)	15.78	6.35	17.73	5.35
T2	FYM (75%) + Vermicompost (25%)	17.67	6.79	16.78	5.42
T3	Poultry manure (75%) + FYM (25%)	21.00	6.11	17.00	5.30
T4	Poultry manure (75%) Vermicompost (25%)	20.89	6.63	16.75	5.15
T5	Vermicompost (75%) + FYM (25%)	23.33	5.69	16.42	5.70

T6	Vermicompost (25%) + Poultry manure (25%)	20.78	6.35	17.56	5.47
T7	FYM (50%) + Poultry manure (50%)	22.67	5.67	17.15	5.27
T8	Poultry manure (50%) + vermicompost (50%)	26.56	6.80	18.34	5.77
T9	Vermicompost (50%) + FYM (50%)	23.33	6.32	17.94	5.35
	Mean	23.33	6.40	17.33	5.42
	'F' Test	S	S	S	S
	C.V. Percent or not check it	0.18	0.14	0.06	6.32
	S.E. (m)	0.03	0.08	0.01	0.28
	C.D. at 5%	0.06	0.02	0.02	0.58

Table 1.5 Effect of organic treatment combinations on Fruit yield per plant, fruit yield per plot and fruit yield per hectare

Treatment Notation	Treatment Combination	Fruit yield per plant (Kg/plant)	Fruit yield per hectare (t/ha)	Fruit yield per plot (Kg)
T0	Control	3.88	10.32	24.31
T1	FYM (75%) + Poultry manure (25%)	3.60	9.62	22.46
T2	FYM (75%) + Vermicompost (25%)	3.84	10.22	24.03
T3	Poultry manure (75%) + FYM (25%)	4.51	11.97	28.60
T4	Poultry manure (75%) Vermicompost (25%)	4.79	12.68	30.44
T5	Vermicompost (75%) + FYM (25%)	3.69	10.11	23.77
T6	Vermicompost (25%) + Poultry manure (25%)	5.10	13.26	32.92
T7	FYM (50%) + Poultry manure (50%)	5.51	14.31	35.65
T8	Poultry manure (50%) + vermicompost (50%)	6.12	15.90	39.77
T9	Vermicompost (50%) + FYM (50%)	5.45	14.16	35.28
	Mean	4.65	12.25	29.72
	'F' Test	S	S	S
	C.V. Percent or not check it	1.98	1.87	0.66
	S.E. (m)	0.07	0.18	0.16
	C.D. at 5%	0.16	0.39	0.33

C) Economic Parameter

Maximum gross returns, net returns and benefit cost ratio was observed in the treatment T₈ (Poultry manure 50% + Vermicompost 50%) and recorded the best treatment among

all the other treatments. As the economics is the need of the farmers while taking decision regarding the adoption of the techniques and scientific knowledge hence, T₈ (Poultry manure 50% + Vermicompost 50%) gave the highest gross return ([????????????????](#)), net return([????????????????](#)), and cost benefit ratio ([????????????????](#)) is due to higher productivity and higher quality of fruits, which increase the market value of the fruits.

CONCLUSION

From the present investigation, it ~~could be was~~ concluded that T₈ (Poultry manure 50% + Vermicompost 50%) was found to be best among all treatment in terms of growth, yield and quality of cucumber i.e. T.S.S. The highest net return was found from T₈ (~~Poultry~~[Poultry](#) manure 50% + vermicompost 50%) with ₹1,56,775 and the B:C ratio that is 1.92.

REFERENCES

Anjanappa, M., Kumara, B. S. and Inderesh, K. M. (2011). Growth yield and quality attributes of cucumber (cv. Hassan local) as influenced by integrated nutrient management grown under protected conditions. *Mysore Journal of Agriculture Science*. **46**(1): 32-37.

Choudhary, K. D. and Basu, N. K. (1967). Phytochemical and hypoglycemic investigation of cucumber. *Journal of Pharmaceutical Sciences*. **56**(11): 1405-1409.


De Candolle A. (1967). Origin of cultivated plants. *Hafner Publication Co.*, New York, (Reprint of 2nd edition. 1959).

National Horticultural Board, Ministry of Agriculture & Farmers Welfare (DAC & FW), Government of India, 2020-21.

Fisher, R. A and Yates, F. (1936). Statistical Tables for Biological, Agricultural and Medical Research. *Oliver and Boyd, London* : 143.

FAOSTAT (2020). www.fao.org/faostat/en/data/CC. Food Supply- Crops Primary

Equivalent. Visited on 15/12/2021.



Ghayal, R. G., Vaidya, K. P. and Dademal, A. A. (2018). Effect of different organic and inorganic fertilizers on growth and yield of cucumber (*Cucumis sativus* L.) in lateritic soil of Konkan (M.S.). *International Journal of Chemical Studies*. **6**(2): 3452-3454.

Hamdi, E. Kh. H., Mosa A. A., Shazly, M. M. E. and Noha R. H. (2017). Response of Cucumber (*Cucumis sativus* L.) to Various Organic and Bio-Fertilization Treatments under an [Organic](#) Farming System. *Journal of Soil Sciences and Agricultural Engineering*. **8**(5): 189-194.

Ikeh, A. O., Udoh, E. I., Uduak, G. I., Udounang, P. I. and Etokeren U. E. (2012). Response of cucumber (*Cucumis sativus* L.) to different rates of goat and poultry manure on an ultisol. *Journal of Agriculture and Social Research*. **12**(2): 132-139.

Jagraj, S., Singh, M. K., Mukesh K., Gupta, A. and Singh, K. P. (2020). Growth, Yield and Quality Parameters of Cucumber (*Cucumis sativus* L.) as Influenced by Integrated Nutrient Management Application. *International Journal of Current Microbiology and Applied Sciences*. **9**(10): 1455-1462.

Kaur, K. and Kaur, A. (2018). Influence of integrated nutrient management practices in cucumber (*Cucumis sativus* L.) c.v. Punjab Naveen. *International Journal of Current Advanced Research*. **7**(12): 16680-16683.

Karpechenko GD. (1925) Chromosomes of Phaseoline. *Bulletin Application of Botany*, **14**: 143-148.

National Horticultural Board, Ministry of Agriculture & Farmers Welfare (DAC & FW), Government of India, 2019-20.

Nasser, J. R. A., Amer, N. K. and Zina, M. A. (2020). Effect of Organic and Chemical Fertilizers on Growth and Yield of Cucumber (*Cucumis sativus* L.). *Indian Journal of Ecology*. **47**(8): 158-164.

Singh, V., Prasad, V. M., Kasera, S. Singh, B. P. and Mishra, S. (2017). Influence of different organic and inorganic fertilizer combinations on growth, yield and quality of cucumber (*Cucumis sativus* L.) under protected cultivation. *Journal of Pharmacognosy and Phytochemistry*. **6**(4): 1079-1082.

Singh, A., Prasad, V. M., Vijay, B. and Samir, E. T. (2021). Effect of organic and inorganic fertilizer on growth, yield and quality traits of cucumber (*Cucumis sativus*) under Prayagraj Agro-climatic condition. *The Pharma Innovation Journal*. **10**(7): 1293-1296.

Tahir, S. M. Kabir, A. M., Ibrahim, H. and Sufiyanu, S. (2019). Studies on the performance of organic and inorganic fertilizer on the growth and yield of cucumber (*Cucumis sativus* L.) *Science World Journal*. **14**(1): 156-163.

MAY BE REDUCED
DELETE IRRELEVANT ONES IN TEXT AND REFERENCES

Formatted: Indent: Hanging: 2.11", Right: 0.11"