

Original Research Article

Field evaluation of selected insecticides and botanical against mustard aphid, *Lipaphis erysimi* (Kalt.) on mustard, *Brassica juncea* L.

ABSTRACT

A field trial was conducted at Central Research Farm [CRF], SHUATS, Naini, Prayagraj during the Rabi season. Six treatments were evaluated against *Lipaphis erysimi*. i.e. Fipronil 5% SC @ 0.5ml/lit, Difenthurion 50% WP @ 0.20g/lit, Profenophos 50% EC @ 1ml/lit, Thiamethoxam 25% WG @ 0.25g/lit, Imidacloprid 17.8% SL @ 0.5ml/lit and NSKE 5% @ 5ml/lit were evaluated against mustard aphid *Lipaphis erysimi*. Results revealed that, among the different treatments, the lowest population of mustard aphid was recorded in Imidacloprid 17.8% SL (22.82). Thiamethoxam 25% WG (33.63) was found to be the next best treatment followed by Difenthurion 50% WP (47.18), Fipronil 5% SC (54.15), and Profenophos 50% EC (60.12), whereas NSKE 5% (80.23) was found to be the least effective against this pest. The plot treated with the highest yield Imidacloprid 17.8% SL (24.6 q/ha) followed by Thiamethoxam 25% WG (21.8 q/ha), Difenthurion 50% WP (18.2 q/ha), Fipronil 5% SC (16.4 q/ha), Profenophos 50% EC (14.3 q/ha) and NSKE 5% (13.5 q/ha) as compared to the control plot (9.2 q/ha). Among the treatments, the best and most economical treatment was Imidacloprid 17.8% SL (1:7.24) followed by Thiamethoxam 25% WG (1:6.43), Difenthurion 50% WP (1:5.24), Fipronil 5% SC (1:4.64), Profenophos 50% EC (1:3.94) and NSKE 5% (1:3.90) as compared to control plot (1:2.83).

Keywords: Cost-Benefit ratio , Evaluation , Imidacloprid , Insecticides , *Lipaphis erysimi*.

INTRODUCTION

“Mustard, *Brassica juncea* (L.) Czern and Coss is an important oilseed crop belonging to the family cruciferaceae (Syn.Brassicaceae). Indian mustard or brown mustard is natural amphidiploids having chromosome no (2n=36). It is self-pollinated but certain amount (2-

15%) pollination occur due to insects and other factors. The origin place of mustard is China, northeastern India from where it has extended up to Afghanistan via Punjab” (Kalasariya, 2016). “Mustard (*Brassica spp.*) is one of the first domesticated crops which has wide dispersal, and has been grown as herb in Asia, North Africa, and Europe for thousands of years” (Mandal *et al.*, 2018).

“It contributes about 28.6% of the total oilseed production in India, whereas it is the second most important edible oilseed after groundnut, sharing 27.8% of India’s oilseed economy” (Kumar *et al.*, 2018). “Mustard plays an important role in the oil seed economy of the country. The oil contents of mustard seeds ranges from 32 to 40 %, and protein content ranges from 15 to 17 %” (Ghule and Bagde 2016).

“Mustard is also rich in minerals like Calcium, Manganese, Copper, Iron, Selenium, Zinc, Vitamin (A, B and C) and Proteins. 1000 g mustard seed contains 508 k. cal. energy, 28.09 g Carbohydrates, 26.08 g Proteins, 26.08 g Total fat and 12.2 g Dietary fiber, 31 I.U. Vitamin A, 4.733 mg Niacin, 7.1 mg Vitamin C, 266 mg Calcium, 9.21 mg iron, 370 mg Magnesium, 13 mg Sodium and 738 mg Potassium” (Daravath *et al.*, 2016).

“Mustard production in India stands in between 8-9 million tons with a significant increase in between 2000 and 2019” (Kalia *et al.*, 2021). “It is cultivated over an area of 5.75 million ha with production and productivity of 6.80 million tonnes and 1183 kg/ha respectively in India (Kumar *et al.*, 2020). In India, the major production regions of mustard are Rajasthan, Haryana, Uttar Pradesh, and Madhya Pradesh while the major consumption (as raw pellets) regions are Rajasthan, Uttar Pradesh, Haryana, and Punjab” (Kalia *et al.*, 2021). “Uttar Pradesh accounts for 1198.5 ha. 10.85% and 11.19% of area and production, respectively in the country with an average yield of 11.49 q/ha which is equivalent to the national average (11.17 q/ha)”. (Singh *et al.*, 2007).

Mustard plant is attacked by a number of insect pests. Bakhetia, 1983 found “more than three dozen insect pests, associated with various phonological stages of these crops. Among them, *Lipaphis erysimi*, commonly known as the mustard aphid is most destructive. It belongs to the family Aphididae of the order Homoptera”. “The insect is distributed to many other countries of the world. The attack is severe in those regions where the number of cloudy days is greater during the pest activity period” (Akter *et al.*, 2021).

“Aphids are small, soft-bodied, pearl-shaped insects that have a pair of cornicles (honey tubes) projecting out from the fifth or sixth abdominal segment. The aphid attacks generally during December and continue till March. About 45 generations are completed in a year” (Gautam *et al.*, 2019).

“The damage is caused by both nymph and adult stages as they suck the cell sap from leaves and turn yellow from green and growth of plants is stunted. It found on all parts of plant; leaves, stem, blooming flowers and silique forming inflorescence, in severe cases, the plants may even die” (Aslam and Ahmad, 2001).

“Mustard aphid is the major constraint responsible for low yield as well as low quality seed, which is considered as key factor in reducing mustard production and can cause yield loss up to 90%” (Gupta *et al.*, 2003). “It may cause a yield losses ranging from 35.4 to 96% in favourable conditions and can reduce 5-6% oil content” (Sahoo, 2012).

“Chemical control is more accurate as it controls about 90% of the aphid population, but due to its high rate of reproduction, in a period of 2-3 weeks after treatment of insecticide, the aphid population reaches around the same number as before treatment. That’s why it is mandatory for insecticides to have effective control over mustard aphid population for a longer period of time” (Ahmed *et al.*, 2018).

Objectives:

- 1.To evaluate the chemical insecticides and botanicals against mustard aphid, *Lipaphis erysimi* (Kalt.) in mustard (*Brassica juncea*) during the Rabi season 2021.
2. To Calculate the Crop Economics – Benefit Cost ratio [B:C ratio].

MATERIALS AND METHODS

The experiment was conducted at the experimental research plot of the Department of Entomology, Central Research Farm, Sam Higginbottom University of Agriculture Technology and Sciences, during the Rabi season of 2021, in a Randomised Block Design with Six treatments replicated three times using variety Rohini seeds in a plot size of 2m×2m at a spacing of 45cm × 30cm with a recommended package of practices excluding plant protection. The soil of the experimental site was well drained and medium high. The research field situated at 25°27' North latitude 80°05' East longitude and at an altitude of 98 metres above sea level. The

maximum temperature reaches up to 42⁰C in summer and drops down to 4⁰C in the winter.

Preparation of insecticidal spray solution:

The desired concentration of insecticidal spray solution for each treatment was freshly prepared each and every time at the site of experiment, just before the start of spraying operations. The quantity of spray materials required for the crop was gradually increased as the crop advanced in age.

The spray solution of the desired concentration was prepared by adopting the following formula:

$$V = \frac{(C \times A)}{\% \text{ a.i.}}$$

Where, V=Volume of a formulated pesticide is required.

C= Required concentration.

A= Total volume of solution to be prepared

% a.i. = Given Percentage strength of a formulated pesticide

Data for mustard aphid was recorded on five randomly selected plants at the top 10 cm of the central shoot. When the aphid population reached an economic threshold level, treatment of insecticides was done. Five plants from each treatment were randomly selected, tagged and the mustard aphid population was recorded on them. Insect populations were recorded 24 hours before application. Besides this, the insect pest population was recorded at time intervals of 3 days, 7 days and 14 days after treatment. The experiment was repeated twice to minimise possible errors.

RESULTS AND DISCUSSION

The population of *Lipaphis erysimi* one day before the first spray was in a range of 120.2 to 136.4 aphids/10 cm central shoot prior to the application of treatments during Rabi. The chemical insecticides were more effective than the botanical. Efficacy of botanical and selected insecticides against mustard aphid populations on mustard showed that the data on the population of *Lipaphis erysimi* on three days after the first spray revealed that all the treatments were significantly superior over control. Among all the treatments, the plot treated with Imidacloprid @ 0.5ml/lit (26.46) proved to be the most effective. The next best treatment was Difenthurion at 0.25g/lit (89.53), followed by Fipronil at 0.5ml/lit (90.26), Profenophos at 1ml/lit (94.6), and @ NSKE 5%, 5 ml/lit (118.26), which was the least effective of all treatments, respectively on the 7 th and 14 th day after spray.

The population of *Lipaphis erysimi* one day before the second spray was in a range of 51 to 166.6. Three days after the second spray, all the insecticidal treatments were significantly superior over the untreated control. The lowest aphid population was recorded in the treatment of Imidacloprid @ 0.5ml/lit (22.6) followed by Thiamethoxam @ 0.25g/lit (30.73), which was a standard check. The next effective treatments were Difenthurion @ 0.20g/lit (36.4), Fipronil @ 0.5ml/lit (43.46), Profenophos @ 1ml/lit (48.067) and NSKE 5% @ 5ml/lit (62.46), which least effective as they recorded significantly higher populations than they rest of the treatments. However, it was significantly superior than untreated control.

The difference in yields among the different treatments was significant. The highest yield was recorded in Imidacloprid 17.8% SL (24.6 q/ha) these findings were supported by **Chandra et al., (2014)** with (24.87 q/ha) followed by Thiamethoxam 25% WG (21.8 q/ha) these findings were supported by **Patel et al., (2017)** with (21.69 q/ha) and **Chandra et al., (2014)** with (21.87 q/ha). The next best treatment was Difenthurion 50% WP (18.2 q/ha) these findings were supported by **Sen et al., (2017)** with (16.17 q/ha) followed by Fipronil 5% SC (16.4 q/ha) which is in line with the similar findings of **Maurya et al., (2018)** with (16 q/ha) and **Patel et al., (2017)** with (16.62 q/ha). The next best treatment was Profenophos 50% EC (14.3 q/ha) which is in line with the similar findings of **Jat and Singh (2005)** with (14.10 q/ha) and the lowest yield was recorded in NSKE 5% (13.5 q/ha) with similar findings of **Kumar et al., (2020)** with (13.90q/ha) as compared to control plot (9.2q/ha).

Conclusion

From the critical analysis of the present findings, it can be concluded that Imidacloprid 17.8% SL is more effective in controlling population of mustard aphid followed by Thiamethoxam 25% WG, Difenthurion 50% WP, Fipronil 5% SC, Profenophos 50% EC and NSKE 5% in managing *Lipaphis erysimi*. Among the treatments studied, Imidacloprid 17.8% SL gave the highest cost benefit ratio (1:7.24) and marketing yield (24.6 q/ha) followed by Thiamethoxam 25% WG (1:6.43 and 21.8 q/ha), Difenthurion 50% WP (1:5.24 and 18.2 q/ha), Fipronil 5% SC (1:4.64 and 16.4 q/ha), Profenophos 50% EC (1:3.94 and 14.3 q/ha) and NSKE 5% (1:3.90 and 13.5 q/ha) as such more trials are required in future to validate the findings.

Table 1. Efficacy of botanical and selected insecticides against mustard aphid *Lipaphis erysimi* (Kalt.)

S.No.	Treatments	Population of <i>Lipaphis erysimi</i> /10 cm apical twig									Yield (q/ha)	B:C ratio
		First spray				Second spray						
		3DAS	7DAS	14DAS	Mean	3DAS	7DAS	14DAS	Mean	Overall Mean		
T ₁	Fipronil 5% SC	90.26	74.86	78.33	81.15	43.66	26.33	11.46	27.15	54.15	16.4	1:4.64
T ₂	Difenthurion 50% WP	89.53	72.26	56.53	72.77	36.40	18.80	9.60	21.60	47.18	18.2	1:5.24
T ₃	Profenophos 50% EC	94.60	83.80	90.33	89.57	48.06	29.60	14.33	30.66	60.12	14.3	1:3.94
T ₄	Thiamethoxam 25%WG	58.80	39.33	50.66	49.60	30.73	14.46	7.73	17.64	33.63	21.8	1:6.43
T ₅	Imidacloprid 17.8% SL	26.46	32.86	38.26	32.53	22.60	11.53	5.20	13.11	22.82	24.6	1:7.24
T ₆	NSKE 5%	118.26	96.53	112.46	109.08	62.46	53.40	38.26	51.37	80.23	13.5	1:3.90
T ₀	Control	132.33	140.46	166.66	146.48	182.53	216.26	193.53	197.44	171.96	9.2	1:2.83
	F-test	S	S	S	S	S	S	S	S	S	-----	-----
	S. Ed (±)	0.35	0.39	0.35	5.28	0.36	0.30	0.24	4.77	19.65	-----	-----
	C.D. (P = 0.5)	0.77	0.86	0.76	19.95	0.80	0.66	0.53	18.00	68.03	-----	-----

REFERENCES

- Ahlawat, P., Singh, R., Singh, S. P., Sachan, S. K., Singh, S. P., Spoorthi, G. S. and Gautam, M. P. (2018)** . Efficacy of bio pesticides and novel insecticides for control of *Lipaphis erysimi* (kalt) on mustard crop in western U.P. *Journal of Pharmacognosy and Phytochemistry.*, SP1:1814-1820.
- Ahmed, S., Cheema, S. A., Zubair, M., Abbas, Q., Bashir, M. R., Malik, K., Aslam, A. and Naeem, A. M. (2018)**. Comparative efficacy of insecticides against mustard aphid in *Brassica juncea*. *International Journal of Entomology Research*. **3**(3): 34-37.
- Akter, A., Hossain, M. I., Amin, A. K. M. R. and Liza, M. M. (2021)**. Efficacy of plant derived and synthetic insecticides against mustard aphid for quality seed production. *Journal of Entomology and Zoology Studies*. **9**(4): 48-53.
- Chandra, A., Malik, Y. P. and Kumar, A. (2014)**. Efficacy and economics of new insecticides for management of aphid (*Lipaphis erysimi*) in Indian mustard. *Current Advances in Agricultural Sciences*. **6**(1): 88-90.
- Dwivedi, S. A. and Singh, R. S. (2019)**. Evaluation of the new molecular insecticides and biopesticides against mustard aphid *Lipaphis erysimi* (Kalt.) on yield parameter in mustard. *Journal of Biopesticides*. **12**(2): 203-214.
- Gautam, M. P., Singh, S. N., Kumar, P., Yadav, S. K., Singh, D. P. and Pandey, M. K. (2019)**. Mustard aphid, *Lipaphis erysimi* (Kalt) (Hemiptera: Aphididae). A review *The Pharma Innovation Journal*. **8**(9): 90-95.
- Jat, S. L. and Singh, B. (2005)**. Relative efficacy of some newer insecticides against mustard aphid, *Lipaphis erysimi*(KALT.).*Current Agriculture*. **29**(1-2): 79-82.
- Kalia, A., Shukla, G., Mishra, D., Mishra, B. P. and Patel, R. R. (2021)**. Comparative trend analysis of mustard in Bundelkhand region, Uttar Pradesh, India. *Indian Journal of Extension Education*. **57**(1): 15-19.
- Kumar, A., Yadav, S., Kumar, Y., and Yadav, J. (2020)**. Evaluation of different botanicals for the management of mustard aphid, *Lipaphis erysimi* (Kaltenbach). *Journal of Oilseed Brassica*. **11**(1): 42-48.
- Maurya, N. K., Singh, R., Singh, J., Nigam, R., Hasan, W. and Kumar, A. (2018)**. Efficacy of novel insecticides against mustard aphid *Lipaphis erysimi* (Kaltenbach). *International Journal of Agricultural Invention*. **3**(1): 62-70.
- Patel, S. and Hasan, C. S. W. (2020)**. Relative efficacy of certain insecticides against mustard aphid in mustard ecosystem. *International Journal of Agricultural and*

- Rashid, S., Hussain, S. M., Yaqoob, M. and Ashraf, S. (2021).** Efficacy of insecticides against mustard aphid, *Brevicoryne brassicae* and their safety to coccinellid predators. *Indian journal of entomology*. **67**(10): 44-55.
- Sharma, N., Upadhyaya, S. N., Singh, U. C., Dubey, M. and Ahmad, A. (2020).** Bio efficacy of insecticides against mustard aphid. *Journal of Entomology and Zoology Studies*. **8**(4): 97-102
- Vishvendra., Kumar, S., Sachan, S. K., Singh, G. and Rajendra, S. (2018).** Bio-Efficacy of insecticides and biorational against *Lipaphis erysimi* (KALT.) in mustard crop. *Journal of Pharmacognosy and Phytochemistry*. **5**: 58-61.
- Yadav, S. K., Singh, R. B., Gautam, M. P., Singh, G. and Giri, S. K. (2018).** Bio-efficacy of insecticides against mustard aphid (*Lipaphis erysimi* Kalt.) on mustard (*Brassica juncea* L.). *International Journal of Chemical Studies*. **6**(2): 2704-2708.
- Zorempuii, R. and Kumar, A. (2019).** Efficacy of certain chemicals and botanicals against aphid, *Lipaphis erysimi* (Kaltenbach) on cabbage (*Brassica oleracea* L.). *Journal of Entomology and Zoology Studies*. **7**(5): 89-93.