

Assessment of soil fertility and creation of thematic mapping in the coastal soils of Ramanathapuram block, Ramanathapuram district in Tamil Nadu, India

ABSTRACT

Aims: A view the current analysis used GPS and GIS techniques to evaluate and map the soil fertility status of the Ramanathapuram block in the Ramanathapuram district of Tamil Nadu, India.

Study design: In this study, a systematic set of 100 georeferenced soil samples were collected from 100 selected villages and analyzed for 10 chemical parameters and the data along with GPS readings were used for the preparation of soil fertility maps using GIS.

Methodology: Hundred soil samples were collected from the sixteen revenue villages of Ramanathapuram block, Ramanathapuram district, Tamil Nadu during the pre-monsoon season with lat-long co-ordinates by using GPS to evaluate the soil fertility. Ramanathapuram block is the middle- coastal block of Ramanathapuram district of Tamil Nadu.

Results: In present study, the mean soil values of pH (8.16), EC (0.74 dSm^{-1}), organic carbon content (2.52 g/kg), Available N ($159.36 \text{ kg ha}^{-1}$), Available P (30.83 kg ha^{-1}), Available K ($355.39 \text{ kg ha}^{-1}$) and (Di-Ethylene Triamine Penta Acidic Acid DTPA) extractable micronutrients viz., Fe (7.21), Zn (0.16), Mn (6.46) and Cu (1.36) ppm were recorded respectively in Ramanathapuram block of Ramanathapuram district.

Thematic maps pertaining to soil fertility were prepared using Arc GIS software 10.1.

Conclusion: The georeferenced sampling sites can be revisited with the help of GPS, which helps in monitoring the soil fertility changes over long run. Further, it will be useful to the researchers, planners, policy makers, extension workers of the State Department of Agriculture, fertilizer industries and farmers.

Keywords: pH, EC, organic carbon, available NPK and DTPA extractable micronutrients

Introduction

Agriculture is an imperative sector for sustained growth of Indian economy. About 70 percent of rural households and eight percent of urban household are still dependent on agriculture for employment. Soil is the basic natural resource for agriculture and it supplies essential nutrients for plant growth, the food security and necessary components of human and animal food and the nutritional security of the country. However continuous cropping of high yielding varieties without proper substitution of inorganic fertilizers or organic manures/ composts and non-

addition of micronutrients have caused excessive removal of essential nutrients from the soil solution/ reserves that eventually led to the deficiencies of major, secondary and in particular micronutrients in soils. The deficiency may either be primarily due to their low contents or secondarily by soil factors that reduce the availability (Sharma and Chaudhary, 2007). Global Positioning System (GPS) and Geographical Information System (GIS) helps in collecting a systematic set of geo-referenced samples and generating spatial data about the distribution of nutrients (Sharma, 2004). The estimation, characterization and comparison of spatial variation of micronutrients are important issues in the site-specific crop management, precision farming and sustainable agriculture (Nayak *et al.*, 2006). Soil nutrient maps covering large areas improve understanding of the nature and extent of nutrient problems, and aid in determining their relationships with climate, soil properties, and soil genetic characteristics determined at similar scales. Intermediate scale maps can be useful in delineating specific areas where deficiencies or toxicities are likely for agriculture, and in determining localized soil characteristics that may be associated with such problems. The thematic maps for individual nutrient (Fe, Zn, Mn and Cu) is prepared by using GIS software (Minakshi *et al.*, 2005 and Nayak *et al.*, 2006) and multi micronutrient maps are generated by integrating individual maps of Fe, Zn, Mn and Cu in the GIS (Sood *et al.*, 2004). This will also help in monitoring changes in micronutrient status over a period of time.

Materials and Methods

Hundred soil samples were collected from the sixteen revenue villages of Ramanathapuram block, Ramanathapuram district, Tamil Nadu during the pre-monsoon season with lat-long co-ordinates by using GPS to evaluate the soil fertility. Ramanathapuram block is the middle-coastal block of Ramanathapuram district of Tamil Nadu. It lies between 9.05° 'to 9.50°' North latitudes and 78.10° 'to 79.27°' East longitudes at an elevation of 2 m. The Geographical area of this Ramanathapuram district has an area is 4,123 km² of rural and urban area. The main source of irrigation is Sarugani River, Manimuthar River, Vaigai River and Vaippar River in Ramanathapuram district. The total cropped area of the district/ zone is 1,72,469 ha (as per 2016-17 G-return). The area under irrigated agriculture is 63,800 ha, while 1,37,099 hectare is under rainfed Agriculture. The major food grain crops cultivated are Paddy (*Oryza sativa*), cholam (*Zea mays*), cumbu (*Pennisetum glaucum*), Ragi (*Eleusine coracana*) and Blackgram (*Vigna mungo L.*). Cotton (*Gossypium hirsutum*) is the major non-food crop grown.

To delineate the soil fertility, 100 soil samples were collected during pre-monsoon seasonie., in the month of July, 2019. The samples were collected in such a manner that it represents the soil fertility of all the revenue villages and overall soil fertility of Ramanathapuram block. The soil samples were collected randomly with GPS co-ordinates. From each revenue village, a minimum of two to maximum of twenty four number of soil samples were collected, properly labelled and brought to the laboratory for further chemical analysis. Soil samples collected from Ramanathapuram block, fields were collected from 2 depths i.e. 0-15 cm depth in polythene bags. The soil collected from each depth was mixed, dried, crushed and sieved with a 2 mm sieve. The prepared soil samples were then stored in polyethylene bags for analysis.

The collected soil samples were analyzed for various parameters by adopting the standard procedures viz., pH by Potentiometry (Jackson, 1958), EC by Conductometry (Jackson 1973), Available Nitrogen by Alkaline Permanganate method (Subbiah and Asija, 1956), Available Phosphorus 0.5 M NaHCO₃ extract (Olsen, 1965), Available Potassium by Neutral Normal (1.0 N) Ammonium Acetate extraction method (Stanford and English, 1949), Organic carbon by Chromic acid wet digestion method (Walkley and Black, 1934) and DTPA extractable micronutrients by Atomic Adsorption spectrophotometer (AAS), (Lindsay and Norwell, 1978). Thematic maps pertaining to soil fertility were prepared using Arc GIS software 10.8. Database on soil available nutrient status was generated in Microsoft Excel package at TNAU and the soil fertility maps were prepared at Indian Institute of Soil Science, Bhopal by using Arc-GIS software version 10.8. The thematic maps on available nutrient status were generated by categorizing the fertility status as 'low', 'medium' and 'high' by showing appropriate legend for OC and available N, P and K; 'deficient', 'moderate' and 'sufficient' for available DTPA micronutrients. The analytical results of each soil sample were categorized as low, medium and high categories for OC and macronutrients and as deficient, moderate and sufficient based on the critical limits for available micronutrients as followed in Tamil Nadu.




Fig.1. Location map of study area

Results and Discussion

Soil fertility status of Ramanathapuram block

pH and Electrical Conductivity

The pH (Table 1) of the soil ranged from 6.07 to 9.81 with a mean of 8.16, respectively. Among the revenue villages, the lowest pH of 6.07 has recorded in Therkutharavai which was followed by 6.35 at Ramanathapuram. The highest pH of 9.81 has recorded in peravoor which was followed by 9.65 at kavanoor. Around 73 percent of samples were fallen under Neutral category (pH 6.0-8.5), 27 percent in Alkaline category and 0 percent in Acidic category. The soils of Ramanathapuram block were predominantly neutral to alkaline. The variation in pH may be due to inherent heterogeneity of soils and also due to the nature of parent material and differences in cultural and fertilizer management practices (Vijayakumar *et al.*, 2015).

The EC of the soil ranged from 0.05 to 8.88 dS m⁻¹ with a mean of 0.74 dS m⁻¹, respectively. Among the revenue villages, the lowest EC of 0.05 dS m⁻¹ has recorded in Madakottan which was followed by 0.06 dS m⁻¹ at Therkutharavai. (Table1) The highest EC of 8.88 dS m⁻¹ has recorded in Ramanathapuram which was followed by 6.45 dS m⁻¹ at pullankudi. Around 80 percent of samples were fallen under Non – saline category (<1.0 dS m⁻¹), 13 percent in slightly saline category, 2 percent in moderately saline category and 5 percent in saline category. The soil samples might be due to proper management and inherent properties of soil as also reported by Sharma *et al.* (2008). The soil samples analysed was found to be non-saline in nature, which might be attributed to light textured soils resulting in free drainage (Verma *et al.*, 2005; Vijayakumar *et al.*, 2015).

Organic carbon (g kg⁻¹)

The overall OC status of the soil (Table 2) ranged from 0.60 to 8.70 gkg⁻¹ with a mean value of 2.52 gkg⁻¹. Among the revenue villages, the lowest OC of 0.60gkg⁻¹ have recorded in both Therkutharavai, Ramanathapuram and Kavanoor which was followed by 0.70 gkg⁻¹ at Kusavankudi and Madakottan. The highest OC of 8.70 gkg⁻¹ recorded at Chidharkottai which was followed by Pullangudi (6.60 gkg⁻¹) and Melakottai (6.30 gkg⁻¹). About 95 per cent of the soil samples in the Ramanathapuram block were mostly belongs to the Mandapam Soil series which is having loamy sand texture (Very coarser in texture) and with less cultivation due to less water availability which may cause less organic carbon content in the block. The low organic carbon status is primarily due to high temperature leading to higher rate of organic matter decomposition (Kameriya 1995) and also due to little or no organic matter additions (Rego *et al.* 2003).

Available N, P and K

The overall available N status (Table 3) ranged from 42 to 455 kg ha⁻¹ with a mean value of 159kg ha⁻¹. Similar to OC, a major percentage of soil samples (95%) were under low status. The minimum available N of 42kg ha⁻¹ has recorded in Sathankulam which was followed by 45 kg ha⁻¹ at Kusavankudi. The maximum available N of 455kg ha⁻¹ recorded at Chidarkottai which was followed by Melakottai (328kg ha⁻¹). Around 95 per cent of samples were fallen under low category (<280 kg ha⁻¹), 5 per cent in medium

category and 95 per cent in low category. As majority of soils is alkaline in nature and having light texture, the applied fertilizers would have been subjected to various losses which resulted in low amount of available N in the soil. This might be due to the fact that being alkaline in major area of the district, applied N in soil is lost through various mechanisms like ammonia volatilization, nitrification succeeding denitrification, chemical and microbial fixation, leaching and run off (De Datta and Buresh 1989) which would have resulted in low amount of available N in soil. The flooded condition during the monsoon leads to leaching losses and barren soil after monsoon leads to volatilization losses in the district.

The Olsen-P (Table 3) ranged from 7.2 to 89.6 kg ha⁻¹ with an overall mean value of 30.8 kg ha⁻¹, respectively. Among the revenue villages, the lowest available P of 7.2 kg ha⁻¹ has recorded in Devipattinam which was followed by 9.40 kg ha⁻¹ at Chidharkottai. The highest available N of 89.6 kg ha⁻¹ recorded at Madakottan which was followed by Ramanathapuram and Peravoor (78.4 kg ha⁻¹). Around 68 per cent of samples were fallen under high category (> 22 kg ha⁻¹), 22 per cent in medium category and 10 per cent in low category. The highest P per centage might be due to the continuous application of rice crop cultivation for prolonged period with the application of Di-Ammonium Phosphate (DAP) that would have built-up soil available P status. High status of P in majority of the soils might be attributed to continuous application phosphatic fertilizers to crops which would have resulted in slow build up phosphorus data as the efficiency of applied P is very low (Aulakh and Pasricha, 1999).

The range of available K was 245.60 to 462.20 kg ha⁻¹ with a mean of 355.39 kg ha⁻¹ and the per cent sample category under low, medium and high was nil, 6 and 94%, respectively. The minimum available N of 246 kg ha⁻¹ has recorded in Kavanoor which was followed by 247 kg ha⁻¹ at Ramanathapuram. The maximum available N of 462 kg ha⁻¹ recorded at Ramanathapuram which was followed by Pullangudi (461 kg ha⁻¹). 'High' available K in these soils may be attributed due to the continuous drain of K from the soil reserve over the years with inadequate supply of chemical fertilizers to meet the crop need, mining of K has started appearing in the soils which is a matter of concern. This shows that if sufficient quantity of potassium is not added externally there will be potassium mining from the soil. These results are in confirmation with the findings of Bhangu and Sidhu, (1991), Naidu *et al.* (2011).

Available micronutrients

The available Fe status (Table 4) varied from 1.5 to 19.8 ppm with a mean of 7.2 ppm. Deficient, moderate and sufficient Fe status was noticed in 31, 36 and 33% of the samples, respectively. Among the revenue villages, the lowest DTPA- Fe of 1.5 ppm has recorded in Valantheravai and Ramanathapuram which was followed by 1.9 ppm at Therkutharavai. The highest DTPA-Fe of 19.8 ppm recorded at Thoruvalur and followed by Kavanur (19.5 ppm). This might be due to precipitation of Fe²⁺ in higher pH of the sodic soils in these soils which was also reported by Verma *et al.* (2007). Similar results were also reported for villages of northern Madhya Pradesh by Rajput *et al.* (2015), Karajanagi *et al.* (2016) for



Malaprabha command area of Karnataka, for Patan district by Patel *et al.* (2016), Wagh *et al.* (2016) for Nagpur district of Maharashtra.

The available Zn status (Table 4) ranged from 0.10 to 0.23ppm with a mean of 0.16ppm. About 100 % of the soil samples were deficient in available Zn with an overall soil status is very low, the present results are in line with the findings of Velu *et al.* (2008). The findings of Shyampura and Seghal (1995) and Katyal and Datta (2004) also subscribe to this view. Climatic conditions, parent materials, and management appeared to be largely responsible for the distribution of Zn in the soil. Coarse texture, high pH, diminishing OC and leaching often accentuated the Zn deficiency (Katyal and Rattan 1993). Zinc content in the investigated soils might be due to the low OC values in these blocks. The results of the present investigation are in the conformity with those of Takkar *et al.* (1997) who envisaged that when the soils are low in organic matter and not supplemented by mineral fertilization they are prone to Zn deficiency.


The available Mn status (Table 1) varied from 0.30 to 16.4 ppm with a mean of 6.46 ppm. Deficient, moderate and sufficient Mn status was noticed in 14, 25 and 61% of the samples, respectively. Among the revenue villages, the lowest DTPA- Mn of 0.3 ppm has recorded in Melakottai and Ramanathapuram which was followed by 0.4 ppm at Therpooki. The highest DTPA-Mn of 16.4 ppm recorded at Sathankulam and followed by Valanthuravai (14.7 ppm). The Mn bearing minerals in the parent material of these soils might be the reason for higher Mn content of soils. Which may be due to the formation of insoluble higher valent oxides of Mn at high pH (Naheed *et al.*, 2010).

The available Cu status (Table 4) varied from 0.22to 3.89 ppm with a mean of 1.36ppm. Deficient, moderate and sufficient Cu status was noticed in 51, 33 and 16% of the samples, respectively. Among the revenue villages, the lowest DTPA- Cu of 0.22 ppm has recorded in pullangudi which was followed by 0.23 ppm at Valuthur. The highest DTPA-Cu of 3.89 ppm recorded at Sathankulam and followed by Kavanur (3.87 ppm). The study area which might be due to the fact that decomposition of organic matter releases micronutrients and also reduces the pH of the soil around the plant roots which helps in increasing the solubility of cationic micronutrients (Sharma and Chaudhary 2007). Hence, deficiencies of micronutrients in the soils are reported to affect the chances of vegetation growth (Arvind Kumar Rai *et al.*, 2011). Thus, this shortfall of micronutrients is adjusted by adding the essential nutrients to the soil either naturally or by artificial fertilizers.


Spatial distribution of different quality parameters of soil fertility in Ramanathapuram block of Ramanathapuram district







Status of OC in Ramanathapuram block




Status of N in Ramanathapuram block








Status of Mn in Ramanathapuram block



Status of Cu in Ramanathapuram block



Conclusion

It can be concluded that based on thematic maps, a major area of Ramanathapuram block, Ramanathapuram district was alkaline, nonsaline, low in OC, low, high and medium in available N, P and K, respectively; with regard to available DTPA micronutrients, Zn was predominantly deficient and Cu was moderate while , Fe and Mn were in sufficient status. The georeferenced sampling sites can be revisited with the help of GPS, which helps in monitoring the soil fertility changes over long run. Further, it will be useful to the researchers, planners, policy makers, extension workers of the State Department of Agriculture, fertilizer industries and farmers. Thus, higher crop productivity through maintaining the soil health and fertility conditions can be achieved through sustainable crop management, Ramanathapuram block, Ramanathapuram district, Tamil Nadu.

References

- Arvind Kumar Rai, Biswajit Paul, & Singh, G. (2011). A study on physicochemical properties of overburdendump materials from selected coal mining areas of Jharia coalfields, Jharkhand, India. *International Journal of Environmental Sciences*, **1(6)**, 1350–1360.
- Aulakh, M.S. and Pasricha, N.S. 1999. Effect of rate and frequency of applied P on crop yield, P uptake and fertilizer P use efficiency and its recovery in groundnut- mustard cropping system. *J. Agric Sci., Cambridge*, **132**: 181-188.
- Bhangu, S.S. and Sidhu, P.S. (1991). Potassium mineralogy of five benchmark soils of central Punjab. *J. PotassiumRes.*, **8**: 243-245.
- De datta, S.K. and Buresh, R.J. 1989. Integrated N management in irrigated rice. *Adv. in Agron.* **10**: 143-169.
- Kameriya, P.R. (1995) Characterization of soils of agro climatic zone of transitional plain of inland drainage (Zone II-A) of Rajasthan, *Ph.D. Thesis*, R.A.U., Bikaner.
- Karajanagi MS, Patil PL, Gundlur SS. (2016). GIS Mapping of available nutrients status of Dundur village under Malaprabha command area in Karnataka. *Journal of Farm Sciences*, **29(1)**:37-40.
- Katyal, J.C. and Datta, S.P. (2004). Role of micronutrients in ensuring the optimum use of macronutrients. *Paperpresented at IFA International Symposium on Micronutrients*, held at New Delhi, India duringFebruary 23-25.
- Katyal, J.C. and Rattan, R.K, (1993). Distribution of zinc in Indian soils. *Fertiliser News* **38**, 15-26.

Minakshi, N.S., V.K. Tur Nayar, P.K. Sharma and A.K. Sood. (2005). Spatial distribution of micronutrients in soils of Patiala district - A GIS approach. *J. Indian Soc. Soil Sci.*, **53(3)**: 324-329

Naheed A, Denich M, Goldbach H (2010). Using GIS approach to map soil fertility in Hyderabad district of Pakistan., In: *Proceedings of 19th World Congress of Soil Science*, **1-6** Brisbane, Australia. Soil Solutions for a changing World, pp. 280-282.

Naidu, L.G.K., Ramamurthy, V., Sidhu, G.S. and Dipak Sarkar. (2011). Emerging deficiency of potassium in soils and crops of India. *Karnataka J. Agric.Sci.*, **24 (1)**: 12-19.

Nayak, A.K., A.R. Chyinchamatpure, G. Gururaja Rao, M.K. Khandelwal and N.K. Tyagi. (2006). Spatial variability of DTPA extractable micronutrients in soils of Bara tract of Sardar Sarovar canal command in Gujarat state India. *J. Indian Soc. Soil Sci.*, **42**: 137-145.

Patel JM, Patel BT, Patel IM. (2016). Fertility status of cultivated soils in Patan district of North Gujarat. *Gujarat Agricultural Universities Research Journal*, **41(1)**:23-27.

Rajput B, Trivedi SK, Gupta N, Toma AS. (2015). Status of Available Sulphur and Micronutrients in Mustard Growing Areas of Northern Madhya Pradesh. *Journal of the Indian Society of Soil Science*, **63(3)**:358-361.

Rego, T.J., Rao, V.N., Seeling, B., Pardhasaradhi, G. and Kumar Rao, J.V.D.K. (2003) Nutrient balances a guide to improving sorghum and groundnut based dry land cropping systems in semiarid tropical India. *FieldCrops Research* **81**, 53-68.

Sharma, J.C. and Chaudhary, S.K. (2007). Vertical Distribution of Micronutrient Cations in Relation to Soil Characteristics in Lower Shiwalika of Solan District in North-West Himalayas. *Journal of the Indian Society of Soil Science*, **55**, 40-44.

Sharma, P.K. (2004). Emerging technologies of remote sensing and GIS for the development of spatial data structure. *J. Indian Soc. Soil Sci.*, **52 (4)**: 384-406.

Shyampura, R.L. and Sehgal, J, (1995). Soils of Rajasthan for optimizing land use, *NBSS publication*, 51 (Soil of India series) NBSS&LUP Nagpur 76 + 6 sheets of soil map.

Sood, A., R.K. Setia, R.L. Bansal, P.K. Sharma and V.K. Nayar. (2004). Spatial distribution of micronutrients in soils of Amritsar district using frontier technologies. In: *Proceedings of 7th Punjab Sci., Congress*. February 7-9 held at Guru Nanak Dev. University, Amritsar.

Takkar, P.N., Nayar, V.K., Bansal, R.L., Dwivedi, R.S. and Manna, M.C, (1997). *Annual Progress Report of ICAR Coordinated Micronutrient Scheme 1996-97*, PAU, Ludhiana.

Velu, V., Usha Mathew and Baskar, A. (2008). Scenario of micro and secondary nutrient deficiencies in the states of Tamil Nadu, Kerala and Pondicherry and amelioration practices for increasing crop production and ensuring food security. *Paper presented in the National Seminar on Micro and Secondary Nutrients for Balanced Fertilization and Food Security* held during 11-12 March, pp. 29-30.

Verma V K, Patel L B, Toor G S and P.K. Sharma. (2005). Spatial distribution of macronutrients in soils of arid tract of Punjab, India. *Int. J. Agri. Biol.*, **7(2)**: 295 – 297

Verma VK, Setia RK, Sharma PK, Khurana MPS, Kang GS (2007). Pedospheric distribution of micronutrient cations in soils developed on various landforms in North-East Punjab. *J. Indian Soc. Soil Sci.* **55(4)**:515-520.

Vijaya Kumar, M. Bakiyathu Saliha, B., Kannan, P. and P.P. Mahendran. (2015). Delineation and geographic information system (GIS) mapping of soil nutrient status of sugarcane growing tracts of Theni district, Tamil Nadu, *African Journal of Agricultural research*, **10(33)**: 3281-3291.

Wagh NS, Mandal DK, Sadanshiv NS. (2016). Available micronutrient status of sunflower growing soils of Nagpur district (Maharashtra). *An Asian Journal of Soil Science*, **11(1)**:225-229.

Appendices

Chart 1 : USDA system of Soil fertility classification

Parameter	Ranges	Soil Fertility Classes
pH	<6.0	Acidic
	6.0 – 8.5	Neutral
	> 8.5	Alkaline
EC (dSm ⁻¹)	<1.0	Non – saline
	1.0 – 2.0	Slightly saline
	2.0 – 4.0	Moderately saline
	>4.0	Saline
Nitrogen (kg ha ⁻¹)	<280	Low
	280 – 480	Medium
	>480	High

Phosphorus (kg ha ⁻¹)	<11	Low
	11 – 22	Medium
	>22	High
Potassium (kg ha ⁻¹)	<118	Low
	118 – 280	Medium
	>280	High
Organic carbon (g kg ⁻¹)	<5	Low
	5 – 7.5	Medium
	>7.5	High

Chart 2 : Micronutrient classification

Parameter (ppm)	Low (Deficient)	Medium (Moderate)	High (Sufficient)
Iron (Fe)	<3.7	3.7 – 8.0	>8.0
Zinc (Zn)	<1.2	1.2 – 1.8	>1.8
Manganese (Mn)	<2.0	2.0 – 4.0	>4.0
Copper (Cu)	<1.2	1.2 – 1.8	>1.8

Table1. Range of values for pH, EC of soil samples of Ramanathapuram block

S. No.	Village Name	No. of samples	pH			EC (dSm ⁻¹)		
			Min	Max	Mean	Min	Max	Mean
1	Kusavankudi	4	7.13	8.72	7.64	0.09	0.17	0.13
2	Sathankulam	4	7.45	8.70	8.21	0.19	0.90	0.46
3	Pattinamkathan	4	6.79	8.55	7.60	0.13	1.45	0.61
4	Therkutharavai	5	6.07	8.79	7.54	0.06	1.15	0.35
5	Madakottan	9	6.60	8.69	7.89	0.05	1.03	0.46
6	Melakottai	4	7.89	8.86	8.33	0.17	1.33	0.63
7	Devipattinum	5	7.87	9.47	8.29	0.35	0.73	0.46
8	Valanthanarai	8	6.60	8.79	7.80	0.13	1.05	0.58
9	Valathoor	2	8.30	9.09	8.70	0.21	0.85	0.53
10	Pullankudi	6	7.84	8.66	8.19	0.36	6.45	1.65
11	Ramanathapuram	24	6.35	9.50	8.11	0.11	8.88	1.51
12	Chidharkottai	7	7.31	8.76	7.84	0.12	0.89	0.48
13	Therpooki	5	7.57	8.70	8.23	0.24	2.91	1.49
14	Thoruvallur	3	8.45	8.90	8.67	1.04	1.53	1.25
15	Peravoor	2	8.20	9.81	9.01	0.35	1.50	0.93
16	Kavanoor	8	7.49	9.65	8.56	0.17	0.49	0.36

Minimum	6.07	8.55	7.54	0.05	0.17	0.13
Maximum	8.45	9.81	9.01	1.04	8.88	1.65
Mean	7.37	8.98	8.16	0.24	1.96	0.74
SD	0.72	0.40	0.43	0.24	2.35	0.47
Std. Err	0.07	0.04	0.04	0.02	0.23	0.05

Table2. Range of values for OC of soil samples of Ramanathapuram block

S.No.	Village Name	No. of samples	OC (g/kg)		
			Min	Max	Mean
1	Kusavankudi	4	0.70	3.90	2.23
2	Sathankulam	4	0.80	2.80	1.88
3	Pattinamkathan	4	1.40	4.80	2.33
4	Therkutharavai	5	0.60	1.80	1.12
5	Madakottan	9	0.70	4.20	2.18
6	Melakottai	4	0.90	6.30	2.98
7	Devipattinum	5	1.10	1.80	1.38
8	Valanthanarai	8	1.50	4.30	3.03
9	Valathoor	2	2.70	3.90	3.30
10	Pullankudi	6	2.10	6.60	3.65

11	Ramanathapuram	24	0.60	4.80	1.95
12	Chidharkottai	7	1.10	8.70	3.29
13	Therpooki	5	1.10	4.20	2.32
14	Thoruvalur	3	3.90	5.30	4.70
15	Peravoor	2	1.30	2.10	1.70
16	Kavanoor	8	0.60	4.00	2.35
Minimum			0.60	1.80	1.12
Maximum			3.90	8.70	4.70
Mean			1.32	4.34	2.52
SD			0.90	1.83	0.92
Std. Err			0.09	0.18	0.09

Table 3. Range of values for N, P and K of soil samples of Ramanathapuram block

S. No.	Village Name	No. of samples	N (kg ha ⁻¹)			P (kg ha ⁻¹)			K (kg ha ⁻¹)		
			Min	Max	Mean	Min	Max	Mean	Min	Max	Mean
1	Kusavankudi	4	45	246	148	10.2	44.8	30.6	293	400	332
2	Sathankulam	4	42	188	126	11.2	56.0	33.6	249	376	329
3	Pattinamkathan	4	78	277	134	9.2	56.0	35.9	330	385	346
4	Therkutharavai	5	64	137	94	10.2	33.6	24.4	270	379	327

5	Madakottan	9	50	216	129	9.2	89.6	38.4	317	418	361
6	Melakottai	4	95	328	178	22.4	56.0	42.0	294	365	340
7	Devipattinum	5	95	126	112	7.2	67.2	30.6	318	457	399
8	Valanthanarai	8	106	241	179	11.2	44.8	25.2	267	369	333
9	Valathoor	2	154	204	179	34.8	44.1	39.5	437	452	444
10	Pullankudi	6	115	319	207	10.1	44.8	20.4	250	461	338
11	Ramanathapuram	24	50	266	141	11.2	78.4	32.7	247	462	347
12	Chidharkottai	7	101	455	200	9.4	33.6	17.3	291	426	361
13	Therpooki	5	76	266	157	10.2	22.4	15.5	326	418	364
14	Thoruvalur	3	227	300	263	22.4	44.8	33.6	314	350	337
15	Peravoor	2	112	182	147	33.6	78.4	56.0	337	368	353
16	Kavanoor	8	73	213	156	9.5	33.6	17.9	246	446	375
Minimum			42	126	94	7.2	22.4	15.5	246	350	327
Maximum			227	455	263	34.8	89.6	56.0	437	462	444
Mean			93	248	159	14.5	51.8	30.8	299	408	355
SD			46.91	80.79	41.45	8.83	18.70	10.65	48.71	38.95	30.41
Std. Err			4.69	8.08	4.15	0.88	1.87	1.07	4.87	3.90	3.04

Table4. Range of values for Fe, Zn, Mn, and Cu of soil samples of Ramanathapuram block

S.No.	Village Name	No. of samples	Fe (ppm)			Zn (ppm)			Mn (ppm)			Cu (ppm)		
			Min	Max	Mean	Min	Max	Mean	Min	Max	Mean	Min	Max	Mean
1	Kusavankudi	4	2.8	13.2	7.4	0.11	0.17	0.14	5.0	11.3	8.4	1.23	3.51	1.98
2	Sathankulam	4	3.3	11.7	6.6	0.11	0.20	0.18	4.1	16.4	9.8	0.80	3.89	2.06
3	Pattinamkathan	4	7.5	15.9	12.2	0.20	0.22	0.21	1.9	8.5	5.0	1.13	1.44	1.28
4	Therkutharavai	5	1.9	14.4	6.5	0.11	0.21	0.16	3.1	14.1	10.2	0.59	3.01	1.88
5	Madakottan	9	2.5	12.9	5.6	0.12	0.19	0.15	3.1	13.9	6.9	0.30	1.51	1.06
6	Melakottai	4	6.8	19.0	11.9	0.12	0.17	0.15	0.3	9.5	4.2	0.63	3.60	1.90
7	Devipattinum	5	3.1	7.5	5.2	0.11	0.16	0.13	0.7	12.7	6.7	0.34	2.52	1.35
8	Valanthanarai	8	1.5	7.8	4.0	0.11	0.19	0.15	2.7	14.7	9.1	0.34	1.53	1.00
9	Valathoor	2	2.0	3.2	2.6	0.15	0.19	0.17	0.7	4.5	2.6	0.23	1.11	0.67
10	Pullankudi	6	2.6	14.4	7.8	0.12	0.19	0.16	1.5	12.3	6.5	0.22	1.90	1.08
11	Ramanathapuram	24	1.5	15.2	7.3	0.11	0.19	0.15	0.3	12.5	5.8	0.36	3.82	1.30
12	Chidharkottai	7	2.6	14.8	7.4	0.11	0.19	0.15	1.5	12.9	6.0	0.38	3.01	1.45
13	Therpooki	5	2.4	18.8	7.1	0.11	0.17	0.15	0.4	11.5	3.6	0.55	1.19	0.93
14	Thoruvalur	3	3.8	19.8	9.5	0.10	0.19	0.14	0.5	7.0	3.6	1.15	1.82	1.47
15	Peravoor	2	2.8	3.4	3.1	0.13	0.19	0.16	5.1	10.4	7.7	0.66	1.03	0.85
16	Kavanoor	8	4.7	19.5	11.0	0.12	0.23	0.18	0.5	14.2	7.3	0.59	3.87	1.58

Minimum	1.5	3.2	2.6	0.10	0.16	0.13	0.3	4.5	2.6	0.22	1.03	0.67
Maximum	7.5	19.8	12.2	0.20	0.23	0.21	5.1	16.4	10.2	1.23	3.89	2.06
Mean	3.2	13.2	7.2	0.12	0.19	0.16	2.0	11.6	6.5	0.59	2.42	1.36
SD	1.7	5.3	2.9	0.02	0.02	0.02	1.7	3.1	2.3	0.33	1.09	0.43
Std. Err	0.2	0.5	0.3	0.002	0.002	0.002	0.2	0.3	0.2	0.03	0.11	0.04