

Effects of TA41 on growth, yield and quality of brinjal

Abstract

An experiment was conducted during 2021 to study the effect of TA41 on growth, yield and quality of brinjal cv. Kashi Sandesh at Vegetable farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj (UP). There were 10 treatments 9 are consisting of different attributes of TA41 and water was used as control. The treatment is sprayed at 30 and 45 days after transplanting. All the treatments were replicated three times in a randomized block design keeping the plot size 1mx2m. The treatment T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L) found the best performances. The maximum Plant height (63.67), Number of leaves per plant (59.74), leaf length (36.76 cm), leaf width (26.76cm). days of first flowering (75.85), no of flowers per plant (20.81), fruit length (8.05cm), fruit width(39.74mm), no of fruits per plant (17.22), fruit yield per plant (1449.33g), average fresh weight of fruit (151.97g), yield per hectare (805.35q/ha), TSS (5.67 Brix), shelf life of fruits (4.11), fruit borer (8.67%), leaf miner (1.00%), little leaf (1.00%), Phomopsis blight (8.33%). The significantly higher gross return (Rs 805350/ha), net profit (Rs 649760/ha) and B:C ratio (5.17) was recorded under T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L. Overall results revealed that the application of TA41 as soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L proved to be better for different growth, yield and quality traits in brinjal.

Keywords: Brinjal, Biostimulator, organic growth promoter, TA41.

Introduction

“Vegetables are the important components in the diet of Indian people as majority of the Indians are vegetarian. Vegetables are the rich source of carbohydrates, protein, vitamins, minerals, fat, crude fiber and elemental salts. Besides, vegetables also enhance the test, colour and texture of the diet” Choudhary (1976). “India occupies prime position in the world in production of vegetable crops ranking 2nd next to China and producing about 162.89 million tons from an area of 9.72 million hectares. However, it is not sufficient to meet the requirement of the present population. India will require about 200 million tons of vegetables annually by the year 2022 AD” Choudhary (1976). “Among vegetables, brinjal is one of the most common and principal vegetable grown in India and other parts of the world, however, higher altitudes are not suitable for its cultivation. There are a large number of cultivars which are being grown throughout the country depending upon yield and consumers preference. It belongs to the family Solanaceae and the scientific name is *Solanum melongena* L. some other names are Eggplant and Aubergine etc” (Solanki et al 2010). “The brinjal is much more important in the warm areas, being grown extensively in India, Bangladesh, Pakistan, China, Philippines. Besides, France, Italy and United State are popular country for its cultivation” • Anonymous (2006). “Brinjal is also

famous as a poor man’s crop because it finds the place among the vegetables, where higher production of vegetables is an important observation” (Bhattarai et al 2021). “The brinjal is staple vegetable in almost all tropical countries in the world and liked by both poor and rich consumers. Since there is a common belief that brinjal is not much valuable vegetable for health. However, it is quite high in nutritive value and can be compared with tomato” (Barakart and Gabr, 1998).

“Generally, solanaceous vegetables require large quantity of major nutrients like nitrogen, phosphorus and potassium, in addition to secondary nutrients such as calcium and sulphur for better growth, fruit and seed yield. The cost of inorganic fertilizer has been enormously increasing to an extent that they are out of reach of the small and marginal farmers and also it is continuously harming the ecological niche. The use of TA41 in such situation is therefore a practically paying proposal” (Chojnacka et al, 2014; Popko et al, 2018). TA41 is a multifunctional organic growth promoter that takes care of viral, fungal, and sucking pests. As a result of this, the quality of crop produce is highly improved and increases the growth and yield of crops by 50%. TA41 effectively controls and prevents soft-bodied sucking pests such as whiteflies, aphids, mites, scale insects, thrips, mealy bugs, and planthoppers. It is very effective

& works quickly against sucking pests. TA41 also helps in retaining soil moisture and prevent soil erosion and develops healthy soil for high yield.

Materials and Methods

The details of the various materials used and methods adopted to lay out the experiment are presented below:

Experimental site

The experiment was carried out at Main Experimental Field, Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) india. The area of Allahabad district comes under subtropical belt in the South East of Uttar Pradesh, which experience extremely hot summer and fairly cold winter. The maximum temperature of the location reaches up to 46°C – 48°C and seldom falls as low as 4°C – 5°C. The relative humidity ranged between 20-94%. The average rainfall in this area is around 1013.4mm annually. However, occasional precipitation is also not uncommon during winter months.

Experimental details

The brinjal crop with Kashi Sandesh variety was grown in Randomized Block Design with 10 treatments and each replicated thrice. Treatment details were T₀(control) without any treatment, T1 soil drenching of TA41 @10ml/L, T2 soil drenching of TA41 @ 10 ml/L +Foliar spray 20ml/L, T3 Soil drenching of TA41 @ 15ml/L, T4 Soil drenching of TA41@ 15ml/+Foliar spray 30ml/L,T5 Soil drenching of TA41@ 20ml/L, T6 Soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L, T7 Soil drenching of TA41 @ 25ml /L, T8 Soil drenching of TA41 @ 25ml/l + foliar spray 50ml/L, T9 Foliar spray of TA41 @ 30 ml/L. Each plot comprised of 2m² with a spacing 60 cm row to row and 30 cm plant to plant and each treatment carrying 9 plants per plot. The observations were recorded on vegetative growth, yield , quality and pest and disease incidence of brinjal as plant height(cm), number of leaves per plant , leaf length (cm), leaf width (cm), fruit length(cm), fruit width(mm), no of fruits per plant , fruit yield per plant (g), days of first flowering , no of flowers per plant , average fresh weight of fruit (g), yield per hectare(g/ha), shelf life of fruits, total soluble solids(Brix) were recorded with the help of hand refractometer, fruit borer (%) , Phomopsis blight (%). the data were collected from 9 plants of each treatment and

analysis of variance technique was used to analyses the recorded data.

Results and Discussion

The present investigation entitled “Effect of TA41 on growth yield and quality of brinjal *Solanum melongena* L. under prayagraj agroclimatic condition” cv. Kashi Sandesh was carried during November 2021 to April 2022 in research field. Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom university of Agriculture, Technology and Sciences, Prayagraj (U.P) India. The experiment was conducted in Randomized block design (RBD) with ten treatments and three replications. The mean data of all the traits were subjected to statistical analysis and salient features of experimental findings are mentioned below:

A. Growth parameter

Plant height

At 150 days after transplanting the maximum plant height was recorded in T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L which is (63.67cm) followed by T8 and T7 which were at par to each other while the minimum plant height was recorded in the treatment T₀ (without any treatment) is (54.63) as presented table 1. This might be due to fact that Soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L act as a nutrilink to plants, increases hormonal, nutritional condition and contributed to a considerable extent for better plant height. Similar results were also reported by **saptari et al., (2013)** who reported that TA41 can promote the activity of xyloglucan endotransglucosylase which cause loosening of cell wall and increases cell permeability. Similar result was found in brinjal by **Meena and Dhaka (2003)**

Number of leaves per plant

At 150 days after transplanting the maximum number of leaves per plant was recorded T₆ (Soil drenching of TA41 @ 20ml /L + foliar spray 40ml / 1L spray tank which is (59.74) followed by T8 and T7 which were at par to each other while the minimum was recorded in the treatment T₀ (without any treatment) is (54.41) as presented

table 1. This might be due to fact that soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L act as a nutriliink to plants, increases hormonal, nutritional condition and contributed to a considerable extent for better number of leaves per plant. The marked increase in growth characters might be due to the possible stimulation of meristematic tissues by auxin accelerating greater cell division and cell enlargement in growing portions. Findings are in accordance with the findings of **Viradia (1982) and El-Soad *et al.* (1976).**

Leaf length

At 150 days after transplanting the maximum leaf length (cm) was recorded in T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L which is (36.76) followed by T₈ and T₇ which were at par to each other while the minimum was recorded in the treatment T₀ (without any treatment) is (27.69) as presented table 2. This might be due to fact that Soil drenching of TA41 @ 20ml /L + foliar spray 40ml / 1L spray tank act as a nutriliink to plants, increases hormonal, nutritional condition and contributed to a considerable extent for better leaf length. The marked increase in growth characters might be due to the possible stimulation of meristematic tissues by auxin accelerating greater cell division and cell enlargement in growing portions.

Leaf width

At 150 days after transplanting the maximum leaf width (mm) was recorded in T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L which is (26.76) followed by T₈ and T₇ which were at par to each other while the minimum was recorded in the treatment T₀ (without any treatment) is (21.79) as presented table 1. This might be due to fact that Soil drenching of soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L act as a neutralino to plants, increases hormonal, nutritional condition and contributed to a considerable extent for better leaf width. **Contreras-Cornejo *et al.*, (2015)** revealed that the secretion of IAA by *Trichoderma* spp. can significantly improve plant and lateral root growth. *Trichoderma* spp. can produce volatile and non-volatile secondary metabolites including 6-n-pentyl-6H-pyran-2-one (6PP), viridin, gliotoxin, harziandione, harzianopyridone, and peptaibols which, as plant-growth promoters, have a significant effect.

Days to first flowering

The data mentioned in table 1 showed that Days to first flowering is recorded maximum in Treatment T₀ without any treatment with (81.04 days) followed by T₁ and T₂ while the minimum is found in T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L which is (75.84days). More intensive flower setting was elicited either by improved plant growth through seaweed extract application or by endogenous components, especially cytokinins, which enhance nutrient partitioning in vegetative plant organ and increase in the transport to assimilates to the growing fruits. A similar effect was observed for eggplant treated with seaweed extract by **Abd El-Gawad and Osman (2014)**

No of flowers/plant

The number of flowers per plant with maximum is recorded (20.81) at T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L followed by T₈ and T₇ which is at par to each other While the minimum at control T₀ (without any treatment) (12.41) This might be due to fact that soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L act as a nutriliink to plants and increases hormonal, nutritional condition and contribute considerable extent for earliness in flowering and fruiting. Under the condition of presented experiment, bio stimulant application also increased the number of pollen tubes on the plants resulted in better reproductive effectiveness and increased fruit yield and quality, described in detail by **Gomez Cadenas *et al.*, (2012)** and **Pohl *et al.*, (2019)** Investigated the effect of the bio stimulant product containing macronutrients on citrus fruit set abscission. The beneficial effect of the bio stimulant resulted from an increase in the photosynthetic efficiency which led to better transport of carbohydrates from leaves of fruit sets.

B. Yield Parameters

Fruit length

The fruit length (cm) with maximum is recorded (8.05) at T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L followed by treatment T₈ and T₇ which is significantly superior over all the treatments. While the minimum at T₂ (6.34). Fruit development is highly affected by auxin formation in the growing seeds and other parts of the fruit to supply food reserves in order to increase fruit development. Moreover, microorganisms that produce auxin are VAM and *Azospirillum* sp. as

microbe, which attaches nitrogen and plays as growth regulator.

Fruit width

The fruit width (mm) with maximum is recorded (39.74) at T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L followed by T5 and T7 which is significantly superior over all the treatments. While the minimum at T3 (25.00). Fruit development is highly affected by auxin formation in the growing seeds and other parts of the fruit to supply food reserves in order to increase fruit development. Moreover, microorganisms that produce auxin are VAM and Azospirillum sp. as microbe, which attaches nitrogen and plays as growth regulator.

The number of fruits per plant with maximum is recorded (17.22) at T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L followed by T8 and T6 and T7 which is significantly superior over all the treatments. While the minimum control at T0 (4.70). The number of fruits per plant is an important determination of yield in brinjal due to providing efficiency part and hormonal balance in the plant system (**Anburani and Manivannan 2002**).

The increase in the number of fruits were associated with increased production of flower, coupled with the reduction in flower and fruit drop that ultimately increased the percentage of fruit set (**Prasad et al. 2013**). Similar results were reported by **Choudhury et al. (2013)** and **Akand et al. (2015)** in tomato.

Number of fruits per plant

Table 1: effect of TA41 on plant height (cm) and no of leaves per plant, leaf length (cm), leaf width (cm), days of first flowering, number of flowers per plant, fruit length (cm), fruit width(mm). no of fruits/plant on brinjal

Treatment no	Treatment details	Plant height	No of leaves/ plant	Leaf length	Leaf width	Days to first flowering	No of flowers/ plant	Fruit Length	Fruit Width	No of fruits/ plant
		150 days	150 days	150 days	150 days					
T ₀	Without any treatment of TA41	54.63	54.41	27.69	21.79	81.04	12.41	6.57	30.93	4.70
T ₁	soil drenching of TA41 @10ml/L	55.26	54.15	28.06	22.89	80.67	13.04	7.03	38.96	5.85
T ₂	soil drenching of TA41 @ 10 ml/L +Foliar spray 20ml/L	55.43	53.30	30.76	24.11	80.11	14.07	6.34	27.89	7.70
T ₃	Soil drenching of TA41 @ 15ml/L	56.89	54.44	28.00	22.63	80.04	14.85	6.75	25.00	8.70
T ₄	Soil drenching of TA41@ 15ml/+Foliar spray 30ml/L	58.78	53.78	31.29	23.37	78.67	16.81	6.92	27.26	10.52
T ₅	Soil drenching of TA41@ 20ml/L	59.18	55.26	31.33	23.96	78.41	17.26	6.23	38.04	11.67
T ₆	Soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L	63.67	59.74	36.76	26.76	75.85	20.81	8.05	39.74	17.22
T ₇	Soil drenching of TA41 @ 25ml /L	59.30	59.30	31.75	23.07	77.30	17.81	7.60	38.15	15.30
T ₈	Soil drenching of TA41 @ 25ml/l + foliar spray 50ml/L	59.74	56.78	32.53	25.81	78.59	18.41	7.70	37.15	16.59
T ₉	Foliar spray of TA41 @ 30 ml/L	57.67	56.56	32.20	25.19	79.26	17.37	7.64	37.11	14.70
F-Test		S	S	S	S	S	S	S	S	S
SEm±		0.43	0.74	1.09	0.67	0.85	0.75	0.16	2.24	0.45
C.D. at 0.5		0.90	2.46	2.28	1.41	1.78	1.58	0.34	4.70	0.94
CV		0.91	1.63	4.30	3.45	1.32	5.64	2.78	8.05	4.86

Fruit yield per plant

The fruit yield per plant (g) with maximum is recorded (1449.33) at T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L followed by T8 and T7 (table 2), which is significantly superior over all the treatments. While the minimum control at T₀ (668.15). The increase in fruit yield might have been due to the better performance of the yield attributes. It may be due to better assimilation of plant nutrients through bio fertilizers (Nanthakumar and Veeragavatham 2000).

Average fresh weight of fruit

The average fresh weight of fruit (g) with maximum is recorded (151.97) at T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L and T followed by T8 which is significantly superior over all the treatments. While the minimum control at T₀ (84.94). The application of biofertilizer may increase fresh weight of eggplant 50.32 %. It was due to the content level of nitrogen, phosphorus, and potassium in TA41 have higher nutrients than other treatments. Similar results were reported by Kumaran et al., (2001), Suryanto and Solanki.

Yield per hectare

The yield per hectare (qq/ha) are shown in table 2 with maximum is recorded (805.35) at T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L followed by T8 and T7 which is significantly superior over all the treatments. While the minimum control T₀ (371.26) The possible reason for increased fruit yield might be associated to better inorganic nitrogen utilization in the presence of bio fertilizers, which enhanced biological nitrogen fixation, better development of root system and possible higher synthesis of plant growth hormones Gajbhiye et al., 2003. Similar trend of work has been noted by Anburani and Manivannan (2002), Devi et al., 2002a, Devi et al., 2002b and Wange and Kale 2004 in brinjal

Quality parameter

Total soluble solids

In terms of Total Soluble Solids (Brix) was recorded maximum in treatments T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L) with (5.67Brix) which is statistically at par with treatment T7 and T8 which is significantly

superior over the other treatment. The lowest TSS of fruit was observed in the treatment T₀ (without any treatment of TA41) which is (5.13 brix) as shown in table 2 . Total soluble solids (T.S.S.), quality of solids, dissolved in the liquid part of brinjal were observed to be increased after treatment with TA41. Total soluble solids (T.S.S.), quality of solids, dissolved in the liquid part of brinjal were observed to be increased after treatment with Azospirillum. The increased in quality due to application of bio-fertilizer and nitrogen, phosphorus and potassium could be attributed to the enhanced photosynthetic and metabolic activities, which resulted in the synthesis of higher number of acids, metabolites and glucose. These reserves ultimately contributed synthesis of TSS in tomato (Kumaran and Natrajan, (2001). These results are in conformity with Kamili et al. (2002) in brinjal.

Shelf life of fruits

In Table 2 is shown the shelf life of fruits. It was recorded maximum in treatment (4.11) T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L followed by T8 and T7 which is superior over the other treatments. The low shelf life of fruits was recorded in T₀ (without any treatment of TA41) is (3.33).

D. Pest and disease incidence

Fruit borer

The minimum fruit borer (%) attack (8.67%) was recorded on T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L. While the maximum (13.00%) recorded in Absolute Control. Only four unrecorded post harvested diseases of fruits and vegetables were described by Dhingra and Mehrotra in 1980. Soft rot of eggplant fruit was caused by *Chrysosporium pruinosum* for controlling Continuous cropping of brinjal on the same piece of land should be avoided. As soon as the insect is detected, the affected parts should be clipped along with the insect and destroyed. Fruits showing any boring should be picked and destroyed. Whole of fruit surface become soft and pulpy, later on fruits emitted foul smell. White coloured spore masses were also seen on the rotten fruit surface.

Leaf miner

The minimum leaf miner (%) attack (1.00%) was recorded on T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L. While the maximum (0.03) recorded in T1 and T9. it could be deduced that the *T. asperelloides* TA41 can modulate the response of the plant, increase resistance, and prevent the suppression of defense genes caused by *R. solani*

Little leaf

The minimum little leaf (1.00%) was recorded on T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L. While the maximum (0.03) recorded in T1 and T9. **Mitra and Majumdar(1980)** described that little leaf is an important disease of Brinjal caused by Mycoplasma. Investigations were carried out to study the uptake of ³²P by healthy as well as little leaf infected tissue. Data revealed that no considerable difference in the uptake of ³²P was higher in TMV infected leaf tissue than in healthy ones

Phomopsis blight

The minimum *Phomopsis* blight (%) (8.33%) was recorded on T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L. While the maximum (11.83) recorded in T1 and T2. **Singh and Chakrabarti (1982)** found that *Phomopsis* rot / blight caused by *Phomopsis vexans*, is a serious disease of Brinjal particularly of the seed crop. The losses caused to the seed crop are much more than that reported on the vegetable crops.

Economics

Maximum gross returns, Net Return and Cost Benefit Ratio Rs. 805350/ha, Rs. 649760/ha and (1: 5.17) respectively was recorded in treatment T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L and the minimum Gross Return, Net Return and Cost Benefit Ratio (Rs. 371260/ha, Rs. 251450/ha and 1:3.09) respectively was recorded in treatment T0 (Control) as presented in table. As the economics is the need of the farmers while taking decision regarding the adoption of the techniques and scientific knowledge Hence, T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L gave the highest gross return, net return, and cost benefit is due to higher productivity and enhanced fruit quality, which increase the market value of the fruit.

Conclusion

On the basis of present investigation, it is concluded that the treatment T₆ (soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L is proved as a best combination of TA41 for growth, yeild and quality. where treatment T₆ significantly enhances the growth, yield and quality traits of brinjal and also gave maximum return/ha.

Table 2: Effect of TA41 on fruit yield/plant, average fresh weight of fruit(g), yield per hectare (q/ha), TSS(Brix), shelf life of fruits, fruit borer (%), leaf miner (%), little leaf (%), Phomopsis blight (%) on brinjal

Treatment no	Treatment details	Fruit yield/ Plant (g)	Average fresh weight of fruits(g)	Yield/ Hectare (q/ha)	TSS(Brix)	Shelf life of fruits	Fruit borer (%)	Leaf miner (%)	Little leaf (%)	Phomopsis blight (%)
T ₀	Without any treatment of TA41	668.15	84.94	371.26	5.13	3.33	13.00	2.00	2.00	11.33
T ₁	soil drenching of TA41 @10ml/L	732.22	84.86	406.84	5.39	3.81	12.67	3.00	3.00	11.83
T ₂	soil drenching of TA41 @ 10 ml/L +Foliar spray 20ml/L	856.96	84.94	476.19	5.37	3.67	12.33	2.00	2.00	1.83
T ₃	Soil drenching of TA41 @ 15ml/L	948.44	95.10	527.02	5.61	3.67	11.67	2.67	2.67	10.33
T ₄	Soil drenching of TA41@ 15ml/+Foliar spray 30ml/L	1027.81	95.10	571.12	5.44	3.70	12.33	2.00	2.00	10.00
T ₅	Soil drenching of TA41@ 20ml/L	1017.74	91.70	565.59	5.71	3.63	9.67	2.00	2.00	9.67
T ₆	Soil drenching of TA41 @ 20ml /L + foliar spray 40ml /L	1449.33	151.97	805.35	5.67	4.11	8.67	1.00	1.00	8.33
T ₇	Soil drenching of TA41 @ 25ml /L	1283.00	151.97	734.10	5.57	4.07	9.67	2.00	2.00	9.00
T ₈	Soil drenching of TA41 @ 25ml/l + foliar spray 50ml/L	1426.70	134.65	792.78	5.48	4.07	9.00	2.00	2.00	8.67
T ₉	Foliar spray of TA41 @ 30 ml/L	1223.89	111.41	680.08	5.55	3.07	9.33	3.00	3.00	9.00
F-Test		S	S	S	S	S	S	S	S	S
SEM±		22.28	7.29	8.68	0.03	0.11	0.63	0.15	0.15	0.37
C.D. at 0.5		46.81	15.31	18.23	0.07	0.23	1.33	0.31	0.31	0.78
CV		2.57	8.21	1.79	0.73	3.60	7.17	8.43	8.43	4.54

References

- **Abd, el. Gawed., H.G., and Osman, H.S., (2014)** Effect of Exogenous Application of Boric Acid and Seaweed Extract on Growth, Biochemical Content and Yield of Eggplant 10.5829/idosi.jhsop **6**.3.1147
- **Akand H., Khairul Mazed H.E.M., Pulok Md. A.I., Chowdhury Md., Shah. N. and Moonmoon J.F. (2015)** Intl. J. Appl. Res., **1**, 71-74.
- **Olumetla , Kumar.A., Ajin,M., Karra , S., Swarup , Chakrabarti., Suman,K., Surat, Sharma. R., Sarvjeet,K., Rameshwar , Sharma. P., (1998)** insect-resistant transgenic brinjal plants [*Molecular Breeding*](#) volume **4**, pages33–37
- **Anonymous (2006)** Estimates of brinjal production. Directorate of Economics. Mini Agriculture, New Delhi.
- **AOAC (2000)** Official methods of analysis, 17th edn. Association of Official Analytical Chemists, Gaithersburg. Chemical Science Review and Letters ISSN 2278-6783 DOI:10.37273/chesci.CS205107192 Chem Sci Rev Lett **2020**, **9** (35), 786-791 Article cs205107192 791
- **Anburani, A. and Manivannan, K. (2002).** Effect of integrated nutrient management on growth in brinjal (*Solanum melongena* L.) cv. Annamalai. South Indian J. Hort., **50** (4-6): 377-386.
- **Barakart, M.A.S., and Gabr, S.M., (1998)** Effect of different biofertilizer types and nitrogen fertilizer levels on tomato plants. Alexandria J. Agril. Res. **43**: 149-60.
- **Bhattacharai,B.R, Pal,A.K.and Amgain, L.P. (2021)** Response of varying levels of phyto-hormones and micro nutrients on growth and yield of brinjal (*Solanum melongena* L.) in sub-tropical Terai region of India. Journal of Agriculture and Natural Resources **4**(2): 40-47
- **Choudhary., (1976)** Vegetable (4th Edition). National Book Trust, New Delhi: 50-58.
- **Colla, G., Rouphael, Y., Canaguier, R., Svecova, E.,Cardarelli, M.,(2014)** Bio stimulant action of a plant-derived protein hydrolysate produced through enzymatic hydrolysis. Front. Plant Science, **5** , 448
- **Chojnacka, K., Michalak, I., Dmytryk, A., Wilk, R., Górecki, H., (2014)** Innovative Natural Plant Growth Bio stimulants. In Advances in Fertilizer Technology; Shishir Sinha, Pant, K.K., Eds.; Studium Press LLC: Houston, TX, USA, Volume 21, pp. 451–489.
- **Calvo, P., Nelson, L, Kloepper, J.W., (2014)** Agricultural uses of plant bio stimulants. Plant Soil , **383**, 3–41.
- **Contreras-Cornejo, H.A.; López-Bucio, J.S.; Méndez-Bravo, A.; Macías-Rodríguez, L.; Ramos-Vega, M.; Guevara-García, Á.A.; López-Bucio, J (2015).** Mitogen-activated protein kinase 6 and ethylene and auxin signaling pathways are involved in Arabidopsis root-system architecture alterations by *Trichoderma atroviride*. Mol. Plant Microbe Interact. **2015**, **28**, 701–710
- **Choudhury S., Islam. N., Sarkar M.D. and Ali M.A. (2013)** Intl. J. Sustain. Agric., **5**, 25-28
- **S.R. Chowdhury, T. Sharmin*, M. Hoque, Md. Sumsujjaman, M. Das and F. Nahar (2013)** evaluation of thrombolytic and membrane stabilizing activities of four medicinal plants of bangladesh. ; Vol. **4**(11): 4223-4227.
- **Devi, H.J., Maity, T.K., Thapa, U. and Paria, N.C. (2000a).** Effect of integrated nitrogen management on yield and economics of brinjal. J. Interacademica, **6** (4) : 450-453.

- **Devi, H.J., Maity, T.K., Thapa, U. and Paria, N.C. (2000b).** Response of brinjal (*Solanum melongena* L.) to different sources of nitrogen. *Veg. Sci.*, 29 (1) : 45-47.
- **Du, Jardin. P., (2015)** Plant bio stimulants: Definition, concept, main categories and regulation. *Sci. Horticulture* 196, 3–14
- **Dutta, N.K., Alam, M.S., Nasiruddin, M., Das, A.K., and Munmun, T.S., (2007)** Efficacy of some new chemical insecticides against brinjal shoot and fruit borer (*Leucinodes orbonalis* Guen.). *Journal of Subtropical Agricultural Research and Development* 5 (3): 301-304
- **olumetla, Kumar. A., Ajin. Karra.M., S., Swarup,S., Chakrabarti.k., Suman, R., Surat , Sharma, K., Sarvjeet. P., Rameshwar , Sharma., (1998).** iinsect-resistant transgenic brinjal plants [Molecular Breeding](#) volume 4, pages33–37
- **Singh, S., and Kapoor, K.K., (1998).** Inoculation with phosphate solubilizing microorganisms and a vesicular arbuscular mycorrhizal fungus improves dry matter yield and nutrient uptake by wheat grown in a sandy soil. *Biol. Fertil. Soils.* 28: 139-144.
- **Gianinazzi, S., (1991)**Vesicular-Arbuscular EndoMycorrhizas Technology in Agriculture: from Genes to Byproducts. Birkhauser, Basel.
- **Gómez-Cadenas, A.; Pérez-Santamarina, R.,(2012).** Ghorbel, R. Effect of a biostimulant product containing macronutrients and a carboxylic acid (AMEC®) on citrus fruitlet abscission. *Acta Hort.* 2012, 189–193.
- **Gajbhiye, R.P., Sharma, R.R. and Tewari, R.N. (2003).** Effect of biofertilizers on growth and yield parameters of tomato. *Indian J. Hort.*, 60 (4) : 368-371.
- **Jan, M.T., Shah,p., Hoolinton, P.A., Khan , M.J., and Sohail, Q., (2009)** Agriculture research: Design and Analysis. Dept. Agronomy, KPK Agric. Uni. Peshawar, Pakistan.
- **Kiran, J., Vyakarana, B.S., Raikar, S.D., Ravikumar, G.H., and Deshpande, V.K., (2010)** Seed yield and quality of brinjal as influenced by crop nutrition. *Indian Journal of Agricultural Research*, 44(1): 1-7.
- **Kumaran, S.S. and Natarajan, S. (2001).** Efficiency of soil conditions and recommended fertilizer on quality parameters of rainfed tomato. *South Indian J. Hort.*, 49 (Special) : 199-201.
- **Kamili, I.A., Zargar, M.Y. and Chattoo, M.A. (2002).** Effect of microbial inoculants, chemical nitrogen and their combination on brinjal (*Solanum melongena* L.). *Veg. Sci.*, 29 (1) : 87-89.
- **Mishra, D.K., Paliwal, D.K., Tailor, R.S. and Deshwal, Alok. K. (2009)** Impact of front line demonstration on yield enhancement of potato. *Indian Research Journal of Extension Education* 9(3): 26-28.
- **Mitra, D.K. and Majumdar, M. (1980).** Absorption of 32p by Brinjal and Tobacco leaf tissues infected with little leaf and Tobacco Mosaic Virus respectively. *Indian Phytopathology.* Vol. :33 619-20
- **Moniruzzaman,M., Khatoon,R., Hossain ,M.F.B., Jamil,M.K., and Islam, M.N. (2014)** Effect of GA3 and NAA on physio morphological characters, yeild and yeild components of brinjal (*Solanum melongena* L.)
- **Mahmood,N., Abbasi,N.A., Hafiz,I.A., Ali,I. and Shehna., (2017)** effect of biostimulator on growth, yield and quality of bell pepper cv.yolo wonder ssn 2076-0906 DOI: 10.21162/PAKJAS/17.5653
- **Mehta, D.N., Singh, K.M. and Singh, R.N. (1979)** Note on extent of damage by *Leucinodes orbonalis* Guenee. *Bulletin of Entomology* 20: 115-116.
- **Meena S.S., and Dhaka R.S., (2003),** Economics of plant growth regulators in brinjal (*Solanum melogena* L.) under semiarid condition of Rajasthan. *Annals Agric. Res.*, 24 (2): 273-275.
- **Mahesh, P., and Men, U. B., (2007)** Seasonal incidence of *Leucinodes orbonalis* on brinjal. *Annals of Plant Protection Science* 15(2): 469-539.

- **Nardi, S., Pizzeghello, D., Schiavon, M., Ertani, A., (2016)** Plant bio stimulants: physiological responses induced by protein hydrolyzed-based products and humic substances in plant metabolism. *Science. Agriculture.* **73**, 18–23.
- **Netam, J.L., and Sharma, R., (2014)** Efficacy of plant growth regulators on growth characters and yield attributes in brinjal (*Solanum melongena* L.) *IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS)*.
- **Nanthakumar, S. and Veeragavathatham, D. (2000).** Effect of integrated nutrient management on growth parameters and yield of brinjal (*Solanum melongena* L.) cv. PVR-1. *South Indian J. Hort.*, 48 (1-6) : 31-35.
- **Patra, S.K., Padhi, A.K. and Mishra, S.N. (1989)** Effect of biofertilizers at graded levels of nitrogen on the yield of wheat and toria in the north-eastern ghat region of Orissa. *Environ. Ecol.* **7**: 533-36
- **Prasad R.N., Singh S.K., Yadava R.B. and Chaurasia S.N.S. (2013)** *Veg. Sci.*, 40, 195-197.
- **Paradiškov, N., Vinkovic, T., Vinkovic, Vrcek. I., Tkalec, M., (2013)** Natural bio stimulants reduce the incidence of BER in sweet yellow pepper plants (*Capsicum annuum* L.) *Agricult. Food Science* **22**, 307–317.
- **Popko, M., Michalak, I., Wilk, R., Gramza, M., Chojnacka, K., and Górecki, H., (2018)** Effect of the New Plant Growth Bio Stimulants Based on Amino Acids on Yield and Grain Quality of Winter Wheat *Molecules* **23**, 470; doi:10.3390/molecules23020470
- **Pohl, A., Grabowska, A., Kalisz, A., and Ekara, A.S., (2019)** Bio stimulant Application Enhances Fruit Setting in Eggplant An Insight into the Biology of Flowering
- **Rajesh Waghunde. R., Rahul, M., Shelake. N., Ambalal, Sabalpara., Trichoderma, A., (2016)** significant fungus for agriculture and environment Vol. **11**(22), pp. 1952-1965, 2 June,
- **Radkowski, A., Radkowska, I., (2013)** Effect of foliar application of growth bio stimulant on quality and nutritive value of meadow sward. *Ecol. Chem. Eng. A*, **20**, 1205–1211.
- **Sam, Ruban. J., Priya, M.R., Barathan, G., and Suresh, Kumar. S.M., (2019)** Effect of foliar application of bio stimulants on growth and yield of brinjal (*solanum melongona*.) e-ISSN:2581-6063 (online), ISSN:0972-5210
- **Saptari R.T., and Dewi K., (2013),** Effect of borax and gibberellic acid on the growth and development of red chilli (*Capsicum annuum* L. “gelora”). *The Third Basic Science International Conference.* (B41):1-3.
- **Sharma, O.P., (2003)** Moth bean yield improvement through front line demonstrations. *Agriculture Extension Review.* **15**(5): 11-13.
- **Solanki, M.P., Patel, B.N., Tandel, Y.N. and Patel, N.B. (2010)** Growth, yield and quality of brinjal as affected by use of bio-fertilizers *Asian Journal. Horticulture.*, **5** (2): 403-406
- **Subbarao, S.B., Aftab, Hussain. I.S., Ganesh, P.T., (2015)** Bio stimulant activity of protein hydrolysate: influence on plant growth and yield. *Journal Plant Science Research*, **2**, 125
- **Singh, D., Chakrabati, A.K., (1982).** Chemical control of Phomopsis fruit of Brinjal. *Indian Phytopathology.* Vol. 35:314- 315
- **Thanki, K.V., and Patel, J. R., (1988)** Seasonal incidence of shoot and fruit borer (*Leucinodes orbonalis* Guenee) on eggplant (*Solanum melongena*) in Gujarat. *Indian Journal of Agricultural Sciences* **58**: 867-868
- **Wange, S.S. and Kale, R.H. (2004).** Effect of bio-fertilizers under graded nitrogen levels of brinjal crop. *J. Soils & Crops*, 14 (1) : 9-11.