

Progress in the treatment of Henoch-Schonlein purpura in children

ABSTRACT:Henoch-Schonleinpurpura(HSP) is the most common leukocyte fragmentary vasculitis with IgA immune complex deposition in childhood, which may affect capillaries, arterioles and venules in various systems of the body.Henoch-Schonlein purpuranephritis(HSPN) can be diagnosed when kidney damage occurs in children with HSP, which is the most serious complication of HSP. If not treated in time, more than 20% of children with HSPN will eventually develop renal failure, seriously affecting the prognosis and quality of life of the children.Therefore, early diagnosis and treatment are crucial to the prognosis of children with HSP. This article reviews the treatment of HSP and HSPN.

Key words: Henoch-Schonleinpurpura; Nephritis; Treatment

Introduction

HSP, also known as IgA vasculitis,[1]is one of the most common vasculitis in children. Its characteristic clinical manifestations are rashes, mainly distributed in lower limbs and buttocks. The rashes are symmetrical, with different shapes and sizes, protrusion of the skin surface and non-discoloring under pressure. About 2/3 children with HSP may develop paroxysms of colic (around the umbilicus or in the lower abdomen), blood in the stool, vomiting and other gastrointestinal symptoms, but most of the abdominal signs are only mild tenderness, and a few children may develop more serious complications, such as intussusception, intestinal obstruction, intestinal perforation and other acute abdominal complications. Joint swelling and pain can occur in about 1/3 of the cases, mainly involving the knee, ankle, elbow and other large joints, the lesions are transient, generally will not lead to joint deformity. According to relevant domestic reports, about 30%-50% of children with HSP may develop kidney damage,[2]called HSPN, which is manifested as hematuria, proteinuria, or tubular urine, accompanied by increased blood pressure and edema. HSPN is a self-limited, benign disease, most of which will recover completely with correct and aggressive treatment, but more than 20% of children with HSPN will eventually develop chronic renal failure if treatment is delayed or recurrent.

As for the treatment of HSP and HSPN, there is no unified plan at home and abroad, and the main treatment includes the following aspects:

1. GENERAL TREATMENT

1.1 Rest

Comment [S1]: severe

Comment [S2]: sentence may be unclear or hard to follow

Comment [S3]: the lower

Comment [S4]: of children

Comment [S5]: severe

Comment [S6]: . Consider inserting a comma to separate the elements.

Comment [S7]: And generally

Comment [S8]: Still more

Comment [S9]: primary

the acute phase should rest in bed, avoid large exercise and long time lower limb ptosis, clinical symptoms within 3 months of remission, prohibition of vigorous exercise, the recovery period can be appropriate physical exercise.

Comment [S10]: The acute

Comment [S11]: extensive

Comment [S12]: three

1.2 Diet

Eat light and easily digestible food, and gradually add other kinds of food after the skin purpura and digestive tract symptoms are significantly reduced or the child's condition is completely recovered. The general order of food addition is grain, vegetable→fruit→pork,[3]and keep away from the allergenic environment. If the allergen of the child is not clear, the allergen test should be given.

Comment [S13]: Consider rephrasing

Comment [S14]: The word addition doesn't seem to fit this context.

1.3 Other

Vitamin C can be added to reduce vascular permeability and edema. In addition, attention should be paid to the amount of access to maintain water and electrolyte balance.

2. SYMPTOMATIC TREATMENT

2.1 Rash

Antihistamines and calcium should be given to children with HSP with urticaria or angioneurotic edema. Montelukast sodium can also be given orally, which can reduce vascular permeability and inflammatory response. Montelukast sodium may also play a role in promoting fibrosis in the pathogenesis of HSP.[4] In clinical treatment, more antihistamines and montelukast sodium combined application. According to the principle of randomized control, Lu Zhijian divided 96 children with HSP into two groups, both of which had received basic treatment. Montelukast sodium was added to the control group, and antihistamines and montelukast sodium were added to the observation group. The results showed that the observation group had shorter symptom disappearance time, lower recurrence rate, lower incidence of adverse reactions and higher safety.[5]

2.2 Symptoms of Digestive Tract

Cimetidine, a H2 receptor blocker, can not only inhibit gastric acid secretion, relieve digestive tract symptoms, but also enhance immunity. However, long-term or large doses of cimetidine can cause male breast swelling and lactation, so male children in adolescence and pre-adolescence should be used with caution.[6] The synthesis of coagulation factors II、VII、

Comment [S15]: Consider rephrasing.

Comment [S16]: consider adding the comma

IX、 X depends on vitamin K. The use of vitamin K in small dose has hemostatic effect, and the use of vitamin K in large dose can relieve the pain caused by gastrointestinal spasm. When gastrointestinal bleeding is obvious, strict fasting and intravenous nutritional support should be given.

Comment [S17]: doses

Comment [S18]: spasms

2.2 Symptoms of Joint

When arthritis or joint pain, NSAIDs should be used on the basis of general treatment. When joint symptoms are more obvious and the above treatment effect is not good, hormones should be added.

Comment [S19]: NSAIDs

2.4 Other

Children with helicobacter pylori (HP) infection may be treated with omeprazole in combination with two antibiotics. Chen Cheng randomly divided children with HSP positive into the treatment group and the control group, and compared the two groups, the results showed that the abdominal pain symptoms of children with HSP in the treatment group treated with omeprazole, clindamycin and metronidazole were relieved faster and the recurrence rate was lower than those in the control group without the above-mentioned drugs.[7]Macrolide antibiotics should be used in patients with positive mycoplasma pneumoniae. When accompanied by infection, sensitive antibiotic treatment is given.

Comment [S20]: The word above-mentioned drugs may be considered archaic or unnecessary legal jargon by your readers

Comment [S21]: Check spelling

3.TREATMENT OF HSPN

There is no unified protocol for the treatment of HSPN, which needs to be individualized according to its clinical classification and/or pathological grade.

3.1 Glucocorticoid (GC)

GC is not recommended when purpura is simple or accompanied by slight abdominal pain or joint pain. GC is a basic drug for the treatment of HSPN. It has strong anti-inflammatory effect and can effectively control inflammatory response, relieve severe arthralgia, abdominal pain and other symptoms in children. It can also act on the glomerular basement membrane, reduce hematuria and proteinuria in children with HSPN, and protect renal function. However, its effect is limited, and long-term massive use of GC may lead to aggravation of infection, hypercoagulability of blood, osteoporosis, etc. In addition, GC cannot prevent the occurrence of kidney damage, shorten the course of disease of children with HSPN, nor reduce the recurrence rate. Therefore, GC alone is not recommended at present, and GC combined with immunosuppressive agents is preferred for the treatment of HSPN.

Comment [S22]: Add the

Comment [S23]: Massivelong-term

Comment [S24]: in

Comment [S25]: or

Comment [S26]: treating

3.2 Immunosuppressor

3.2.1 Cytoxan (CTX)

GC combined with CTX shock therapy for HSPN is commonly used in clinical practice, and the curative effect is relatively positive. It has been reported that long-term use of CTX can increase the incidence of alopecia, gonad damage, hemorrhagic cystitis and other complications, and can also cause nephrotoxicity and myocardial damage when the dose is greater than 120mg/kg, so the use of CTX is restricted.[8] According to Liu Fengzhi, the

Comment [S27]: It

Comment [S28]: The word greater is often overused

cumulative dose of CTX in the treatment of HSPN children should not exceed 300mg/kg.[9]

3.2.2 Mycophenolate Mofetil (MMF)

MMF is a new immunosuppressant that can selectively block the proliferation of B and T lymphocytes and rapidly inhibit the proliferation of macrophages and the formation of antibodies, thus exerting immunosuppressive effects.[10] GengHaiyun et al. analyzed 68 patients with macroproteinuria (macroproteinuria: HSPN children with 24h urinary protein quantitative $\geq 50\text{mg/kg}$ or morning urinary protein/creatinine (mg/mg) ≥ 2.0)[11] were randomly divided into CTX group and MMF group. The treatment regimen of CTX group was MP+ACEI+CTX, and MMF group was MP+ACEI+MMF. The complete response rate, response rate, urine protein turning negative time and adverse reactions were compared between the two groups, and the results showed that CTX and MMF had comparable efficacy and safety in the treatment of HSPN children with massive proteinuria.[12] However, MMF has the advantage of convenient administration and no gonadal damage compared with CTX.

Comment [S29]: Add the

3.2.3 Cyclosporin A (CyA)

CyA is a potent inhibitor. CyA combined with GC can treat refractory HSPN. It has also been reported that for children with nephrotic proteinuria who are resistant to prednisone, CTX and AZA at the same time, CyA can significantly reduce urinary protein.[13] CyA can not only lead to changes in glomerular structure and function, but also cause adverse reactions such as hepatotoxicity, nervous system lesions and gastrointestinal reactions. Time and dose of medication are related to the occurrence of adverse reactions, so the above side effects can usually disappear after dosage reduction or withdrawal.

3.2.4 Azathioprine (AZA)

AZA is a highly effective immunosuppressant, which produces immunosuppressive effects by inhibiting the proliferation of B and T lymphocytes. The main adverse reactions are bone marrow suppression and liver damage. AZA combined with GC is a common treatment for HSPN. Fotis et al. showed in a study that AZA+GC could effectively control the long-term symptoms of children with HSPN, while allowing earlier GC discontinuation without serious adverse reactions.[14]

3.2.5 Tacrolimus (FK506)

FK506 is highly immunosuppressive and has a similar mechanism of action to CyA, but the immunosuppressive effect is 100 times stronger than CyA. In the treatment of children with HSPN, FK506 combined with GC is superior to CTX combined with GC, and the incidence of adverse reactions is lower.[15]

3.2.6 Mizoribine (MZR)

MZR is a metabolic immunosuppressant that can inhibit cell proliferation in the lymphatic system.[16] Clinical studies have shown that MZR has mild gastrointestinal symptoms and bone marrow suppression, which is positively dose-dependent and sometimes leads to increased uric acid. Studies have reported that MZR combined with GC has a higher remission rate and a smaller incidence of adverse reactions in children with severe HSPN compared with CTX intravenouspercussion therapy.[17]

3.2.7 Leflunomide (LEF)

LEF is an immunomodulator. A meta-analysis of LEF treatment for HSPN showed that LEF treatment for HSPN has certain advantages and is safer.[18] Another meta-analysis of LEF in the treatment of refractory nephropathy showed that the effectiveness of LEF was comparable to CTX and LEF had fewer adverse reactions.[19] The study of Zhang Miaoge et al showed that LEF had significant effect on the disappearance of hematuria and urinary protein and the improvement of renal function [20].

Comment [S30]: Add a

4.RITUXIMAB (RTX)

RTX is a monoclonal antibody. It is now used to treat a variety of refractory and recurrent kidney diseases. Yin Jing et al reported 2 cases of refractory IgAV in children with recurrent rash, abdominal pain and gastrointestinal bleeding. High-dose hormone therapy was effective. Although immunosuppressants were also used, the symptoms of hormone reduction were repeated, and the symptoms disappeared rapidly after RTX administration, and hormone withdrawal was successfully stopped within 2 months and continued remission.[21] The sample size of this study is small, so how to select suitable IgAV patients for RTX treatment, timing of RTX use, appropriate dose and course of treatment, and the impact on long-term prognosis need to be observed in a larger sample and for a longer period of time.

Comment [S31]: IgA

Comment [S32]: two

Comment [S33]: Add the

5.NGIOTENSIN CONVERTING ENZYME INHIBITOR (ACEI)/ANGIOTENSIN RECEPTOR ANTAGONISM AGENT (ARB)

Comment [S34]: Check the word

The main functions of ACEI and ARB are to dilate glomerular arteries, reduce glomerular pressure, protect renal function, prevent kidney damage, reduce proteinuria, relieve hypertension and edema. ACEI/ARB was routinely used for children with proteinuria.[12]

6. ANTIPLATELET AGGLUTINATION DRUGS/ANTICOAGULANTS

Antiplatelet agglutination drugs/anticoagulants Aspirin and dipyridamole can inhibit platelet adhesion and aggregation, and reduce the formation of coagulation and microthrombus. Dipyridamole is often used clinically. Low molecular weight heparin can not only reduce the hypercoagulability of children with HSPN, but also regulate the complement level and antagonize the type IV allergy mediated by immune cells [22]. Other studies have reported that

the addition of thrombolytic urokinase therapy for severe HSPN can reduce glomerular lesions.[23]

7.GAMMAGLOBULIN

GC combined with gamma globulin can significantly relieve gastrointestinal symptoms of children with severe HSP and reduce secondary kidney damage, with good long-term efficacy.[24] High-dose gamma globulin shock therapy can effectively relieve symptoms of HSPN children with hormone dependence or hormone resistance.[25]

8.OTHER DRUGS

8.1 Hydroxychloroquine(HCQ)

HCQ reduces the damage of autoantibodies to human cells. A foreign clinical trial confirmed that HCQ can relieve the disease of children with HSP.[26] Due to its effectiveness, safety and extremely low cost, as well as its anti-thrombosis and anti-lipid disorders, it may be used as an immune modulator to treat HSP in the future, but its potential side effects on children with HSP need further study.

Comment [S35]: could

8.2Dapsone

Dapsone is an anti-inflammatory drug, which can reduce the permeability of capillaries and provide a basis for the treatment of vascular inflammatory diseases. Dapsone can significantly reduce the expression of IL-6 and IL-8 in serum of children with HSP, and has a significant effect on regulating the cascade reaction of cytokines in serum, and has a significant therapeutic effect on children with HSP.[27] However, its specific side effects on children with HSP remain to be further studied.

8.3 Anisodamine (654-2)

654-2 can be used to treat refractory HSP. It is an M choline receptor blocker, which has the effects of relieving vasospasm, anti-septic shock, analgesia, etc. Clinically, other drugs combined with 654-2 are used to treat CHILDREN with HSP.

8.4 Alfalcitol

Alfacalcitol is a calcium modulator that acts in the liver in the formation of 1, 25-(OH)2D3. 1, 25-(OH)2D3 can regulate calcium and phosphate metabolism, promote bone mineralization, and regulate immune function and inflammatory response. Fu Qiang et al showed that the addition of alfacalcitol to children with HSP can increase the level of 1, 25-(OH)2D3 in serum, reduce the level of inflammatory factors in vivo and improve cellular immune function, and reduce the recurrence rate and the incidence of renal damage in children with HSP.[28]

9. NON-DRUG TREATMENT

9.1 Blood Purification

Commonly used blood purification techniques include plasmapheresis (PE), hemoperfusion (HP) and hemodialysis (HD), which are mainly used for the treatment of HSPN children with acute nephritis or HSP children with severe complications. PE is to extract the blood from the body of children with HSP, and then separate the pathogenic components in the extracted plasma and discard them. Finally, the non-pathogenic components in the plasma and the supplemental albumin or balance liquid are transfused back to the body of children. The mechanism of action of HP is similar to that of PE, and compared with PE, HP can avoid the occurrence of blood infections and the loss of plasma components. HD is the use of semi-permeable membrane principle, complete blood purification, correct water and electrolyte disorder, maintain acid-base balance. Studies have reported that HP combined with HD has a more significant effect on children with severe HSP[29] Blood purification technology can rapidly relieve the symptoms of children, reduce proteinuria, alleviate kidney injury, and improve the long-term prognosis of children, but whether it can prevent and treat kidney injury remains to be supported by clinical evidence-based medical evidence.

Comment [S36]: It seems that there is an article usage problem here.

9.2 Tonsillectomy

Tonsil is the first defense barrier of human body. When stimulated by antigen, tonsil can stimulate B lymphocytes to produce IgA1, accelerate the deposition of IgA1 immune complex in the glomerular basement membrane, and damage renal function. Tonsillectomy can reduce pharyngeal infection, reduce hematuria and proteinuria, maintain the stability of renal function, and reduce the recurrence of the disease. This is surgical treatment, and clinical indications should be strictly grasped, especially for children with recurrent tonsillar infection who still have hematuria and proteinuria after routine treatment. Tonsillectomy should be performed after 2-5 weeks of local inflammation control. However, there is no clear conclusion about the effect of HSPN on tonsillectomy in children at home and abroad.

Comment [S37]: Add the

Comment [S38]: renal damage

9.3 Kidney Transplantation

If kidney failure eventually develops, kidney transplantation may be used, but recurrence is still possible.

10、CONCIUSION

Apart from the above treatment, Chinese medicine in the treatment of HSP and in children with HSPN, obtains the good curative effect, also can be adjusted as a whole with the body's immune, studies have shown that tripterygiumglycosides in the treatment of HSPN has made good effect, but its main toxic ingredients is also one of effective components, and its use to make clear on the manual banning the use of children, Therefore, the 2016 evidence-based guidelines for the diagnosis and treatment of Purpura nephritis no longer recommended the use of Tripterygium wilfordiipolyglycosides in the treatment of HSPN.

Comment [S39]: a good

Comment [S40]: add and

Comment [S41]: add the

REFERENCES

1. SaulsburyFT. Clinical update. Henoch—Schonlein purpura. Lan—cet, 2007, 369(9566): 976—978.
2. Hu Yamei, Jiang Zaifang, Shen Kunling, et al. Zhu Futang Practical Pediatrics [M]. 8th edition. Beijing: People's Medical Publishing House,2015:733-744.
3. Xu Hailing, Chen Xiaoyu, Fang Liyan et al. Clinical significance of dietary management in children with henochschonlein Purpura [J]. Chinese Medical Science,2019, 4: 52—54,138.
4. Lu ZJ. Clinical efficacy and safety of montelukast sodium combined with loratadine in the treatment of pediatric allergic purpura [J].2021, 27(15): 55—57.
5. Wu SH,LiaoPY,YinPL,etal.Inverse temporal changes of lipoxin A4 and leukotrienes in children with Henoch-Schonlein purpura[J].Prostaglandins,leukotrienes,and essential fatty acids,2009,80(4): 177-183.
6. Immunology Group, Pediatrics Branch, Shanghai Medical Association. Expert consensus on the clinical use of immunomodulators in children (Shanghai) [J]. Chinese Journal of Clinical Pediatrics,2018, 33(9): 651—664.
7. Chen Cheng, Ouyang Guodong, Gong Changfu. Efficacy of sequential therapy against Helicobacter pylori in children with henochschonlein Purpura abdominal type [J]. Chinese Modern Medicine Application,2014, 8 (7) : 177-178.
8. Santeusanio AD, Menon MC, Liu C, et al. Influence of patient characteristics and immunosuppressant management on mortality in kidney transplant recipients hospitalized with coronavirus19 (COVID) [J]. Clin Transplant, 2021, 18 (47) : 236-239.
9. Liu FENGzhi. Value analysis of high-dose cyclophosphamide and methylprednisolone in the treatment of allergic purpura nephritis [J]. Electronic Journal of Cardiovascular Diseases of Integrated Traditional And Western Medicine, 2018,6 (19) : 181.
10. Dan YU, Nephrology DO, Hospital WC. Clinical Efficacy of Mycophenolate Mofetil Combined with Low-Dose Hormone in the Treatment of Minimal Change Nephrotic Syndromes[J]. China J PharmaceuEcono, 2019, 12 (17) : 236-239.
11. Nephrology Group, Pediatrics Branch of Chinese Medical Association. Evidence-based guidelines for the diagnosis and treatment of common kidney diseases in

children (-) : diagnosis and treatment of hormone-sensitive, recurrent/dependent nephrotic syndrome Evidence-based Guidelines (Trial)[J]. Chinese Journal of Pediatrics,2009,47(3):167-170.

12. GengHaiyun, Chen Chaoying, Li Huarong et al. Mycophenolate and cyclophosphamide in the treatment of children with proteinuria type allergic purpura nephritis: a prospective randomized controlled study [J]. Chinese Journal of Current Pediatrics,2021,23 (4) : 338-342.

13. Nephrology Group, Pediatrics Society of Chinese Medical Association. Evidence-based guidelines for diagnosis and treatment of Purpura nephritis (2016)[J]. Chinese Journal of Pediatrics,2017, 9(55): 647—650.

14. FotisL,TuttlePV,BaszisKW,et al. Azathioprine therapy for steroid-resistant Henoch-Schönlein purpura : A report of 6 cases[J]. PediatrRheumatol Online J,2016,14 (1) : 37.

15. Yu Juan, Zhang Yumei and Zhao Lijun. Clinical observation of low-dose prednisone combined with tacrolimus in treatment of children with henoch-schonlein purpura nephritis [J]. Chinese pharmacists,2019, 22 (11) : 2055-2057.

16. Yan Qiang, Sui Weiguo, Zou Hequn. New progress in clinical application of immunosuppressive imizolibin [J]. International Journal of Urology,2008,28(03):418-421.

17. Li Liangzhi, Liu Lei, He Xianhong et al. Treatment of severe allergic purpura nephritis with mizolibin: a clinical observation of 21 cases [J]. Chongqing Medical Science,2010, 39(14) : 1904—1906.

18. Ni Jinyu, ZHANG Hanfang, LI Bing, ZhaiWensheng. Leflunomide in the treatment of Purpura nephritis: a meta-analysis [J].2020, 24 (6) :1061-1066.

19. Wang Fei, Hu Shunjin, Wang Hongyu, et al. A meta-analysis of leflunomide and cyclophosphamide in the treatment of refractory nephrotic syndrome [J]. Anhui Medicine,2018, 22 (6) : 1182-1186.

20. Zhang Gemiao, Cui Jianjun, Fang Qiwei. Clinical observation of leflunomide in the treatment of children with henoch-Schonlein Purpura nephritis [J]. Chinese Medicine and Clinical,2015, 15 (11) : 1616-1618.

21. Gong Q, Li Q. Clinical study of gamma globulin in the treatment of severe abdominal allergic purpura in children [J]. Modern digestion and interventional therapy,2016, 21 (5) : 738-740.

22. Yin Jing, Zhao Qianqian, Ma Jijun et al. Rituximab in the treatment of refractory immunoglobulin A vasculitis in children: two cases [J]. Chinese Journal of Rheumatology,2021,25(12):823-826.

23. Ohara S, Kawasaki Y, Miyazaki K eta1. Efficacy of cyclosporine A for steroid—resistant severe Henoch-Schonlein purpura nephritis [J].Fukushima J Med Sci. 2013. 59(2): 102—107.

24. Yao Yao. Application and dose analysis of low molecular weight heparin calcium in the treatment of children with Purpura nephritis [J]. Systems medicine,2018,3 (14) : 87-89.

25. Xia Liping, Chen Xu, JIANG Yi. Therapeutic effect of gamma globulin shock therapy on children with abdominal allergic purpura [J]. Chinese Journal of ContemporaryPediatrics,2016,18(10):988-990.

26. Casian A, Sangle S, D' Cruz DP. New use for an old treatment: Hydroxychloroquine as a potential treatment for systemic vasculitis. Autoimmun Rev, 2018, 17: 660-664.

27. Duan Zhijuan. Clinical observation on the effect of dapsone on serum IL-6 and IL-8 in children with henochschonlein Purpura [J]. China's medical frontier,2010,5 (16) : 55-56.
28. Fu Qiang, SHI Mingfang, Chen Ying. A prospective randomized controlled study of alfalacccitol in the treatment of children with henoch purpura [J]. Chinese Journal of Contemporary Pediatrics,2021, 23 (8) : 797-801.
29. Huang Xin-Ying, XIONG Yun, XIONG Hui-ling. Nursing experience of hemoperfusion tandem hemodialysis in treatment of severe allergic purpura in children [J]. Practical clinical medicine,2009,10(4):122-126.