

# Factors Affecting Pediculosis Capitis Transmission Among Primary School Children

## Abstract

**Introduction:** Human head lice infest people worldwide and are most prevalent in children. Outbreaks of head lice more commonly affect children 3 to 12 years old, with girls being affected more commonly than boys. The aim is to study the factors affecting transmission of the disease among the school children in villages in Benghazi - Libya

**Patients and methods:** In a cross-sectional study in western region of Benghazi, 556 students in three schools from Alhlys, Altry and Abofkra regions were enrolled. The ages of children were ranged from 6 to 13 years. Data collected include socio-demographic characteristics of the students, sex, age, school and hair was examined individually for head lice infestation in a separate room. Descriptive analysis and Chi-square test was used to determine the statistical significance of  $p$  value  $\leq 0.05$ .

**Results:** The data collected on 556 students from the 3 villages, 39.7% from Al-Helys, 28.8% from Al-Tarya and 31.5% from Bo-Fakra regions. The age of children ranged from 6 to 13 years. Female children were more than 50%. The mean of family size was  $7 \pm 2$  persons, number of bedroom in houses was ranging from 1-5 rooms. Number of person in each bed room ranged from one person to eight persons, 34.7% was two person and 33.3% for three persons. Sharing of personal equipment as hair comb and linen was reported in 33.5% of families. Pediculosis capitis manifestation was reported in 26.3% of the participants. The prevalence of pediculosis capitis in Al-Helys 19.5%, in Al-Tarya was 35.6% and in Bo-Fakra 26.3%, this difference was statistically significant and  $P$  value was  $.05 >$ . The prevalence of pediculosis capitis in males was 10.2% while in females was 41.8%, this difference was highly statistically significant ( $P = 0.0001$ ). The highest prevalence of pediculosis capitis was in age 11 years (42.9%) followed by age 10 year (36.6%). There was no reported cases at the age of 6 and 13 years. There was no effect of parent occupation or education level on pediculosis infestation transmission.

**Conclusion:** The present study revealed that, pediculosis capitis was prevalent in school children, more in females. Furthermore, pediculosis infestation was higher in certain village than others.

**Keywords:** pediculosis capitis, school children, prevalence, transmission

## Introduction

Pediculosis capitis is an infestation with the human head louse (*Pediculus humanus capitis*). They are ectoparasites where their infestation clinically present with scalp pruritus, excoriations, cervical lymphadenopathy, conjunctivitis and hypersensitivity rash or pediculid that mimic a viral exanthema (1,2,3). Head lice infestation crosses all economic and social boundaries (4,5). Head lice are 1 mm to 3 mm long. The head louse is an obligate parasite that spends its entire life on the human host. Head lice feed exclusively on blood, unable to jump or fly, and transmission requires close contact. Transmission of head lice is thought to occur by head-to-head contact,

sharing of headgear, or other direct contact with fomites (6,7). The life cycle of the head louse has three stages: egg, nymph, and adult. Nits are head lice eggs hard to see and often confused for dandruff or hair spray droplets. Nits are laid by the adult female and are cemented at the base of the hair shaft nearest the scalp (1,2,8). They are 0.8 mm by 0.3 mm, oval and usually yellow to white. Nits take about 1 week to hatch (range 6 to 9 days). Viable eggs are usually located within 6 mm of the scalp (2,3). The aim of this study was to assess the prevalence of head pediculosis capitis among school children in the villages in Benghazi – Libya and to study factors affecting transmission of the disease among this age group.

## **Materials and Methods**

In a cross-sectional study, 556 students in three schools from three villages in west region of Benghazi (Alhlys, Altry and Abofkra). The age of children in this study were ranged from 6 to 13 years. Data collected regarding the selected socio-demographic characteristics of the students, sex, age, school, class, medical history, complaint if any. The hair of each child was examined individually for head lice infestation in a separate room, and pediculosis capitis was defined as the finding of living adult, nymph, or egg (nits) for all students. Questionnaire had been sent to parents of the children includes demographic data for mothers and fathers. The observation of nits and nymphs or adult lice was considered as a positive.

Data was analyzed using (SPSS) statistical package of social science program version 23. The statistical analysis included: I. Descriptive Statistics: Including (Mean value, Standard deviation, Number and Percentage). II. Inferential Statistics: was used when needed as t- test and Chi-square, P-value will be considered significant when  $\leq 0.05$ .

Ethical statement: This study was approval by the local ethics committee. Informed written consent was obtained through a consent form that was given to the participants along with the questionnaire.

## **Results**

The data collected on 556 children from three schools located in three different region in western of Benghazi-Libya (Al-Helys, Bo-Fakra and Al-Tarya) represented as 39.7%, 31.5% and 28.8% respectively. The percentages of age of participants were high in 10-13 years and 6-9

years groups and presented as 50% each group. Females were slightly higher than male (50.7% Vs 49.3%). Pediculosis capitis manifestation was reported in 26.3% of the participants ( Fig.1).

The diagnosis was depending on demonstration of nits and rarely on live louse ( Fig.2,3).

Pediculosis capitis was more prevalent in Al-Tarya region ( $P=0.002$ ) (Table 1). Furthermore, there was significant difference between types of pediculosis infestation and genders ( $P=0.0001$ ) in which girls had more pediculosis infestation than boys (Table 2). Pediculosis increased with children ages and reaches the peak at age 11 years old ( $P=0.0001$ ) (Table 3).

Parents occupations and education levels have been investigated and showed no statistically significant relation with pediculosis infestations ( $P > 0.05$ ) (Table 4,5). The other variables such as family sizes, number of bed rooms, number of person in each bed room, charring of personal equipment as hair comb and linen. There was no significant relationship between these variables and pediculosis capitis (Table 6,7).

## **Discussion**

The prevalence of pediculosis capitis in this study was 26.3%, which was lower than the study conducted in Sabha city that shown the head lice infestation was 38.6% (9). In literature, there are various studies which have reported prevalence rates of pediculosis capitis throughout the world among school children. In the Americas, prevalence varies from 3.6 % to 61.4 % and is higher in females (10). In Europe, prevalence oscillates from 0.48 to 22.4 %. In Asia, prevalence ranges from 0.7 to 59 %, being higher in girls and women. In Africa, the majority of studies were applied in Egypt and prevalence varied from 0 % to 58.9 % and was higher in females as well (11). In the present study, the in our country, like a study applied by the same authors in 2015 that determined a prevalence of 11.5 % in a similar institution (12). In Yemen, the overall prevalence of head lice infestation was 13.3%; 18.9% in girls and 8.6% in boys )(13). In Argentina, the overall prevalence of head lice infestation in primary school students was 27.9%, which was significantly higher in girls than in boys (14). Despite the progress in medical science and development of the civilization, pediculosis capitis is still an important health issue worldwide (15). A large variation in the global spread of head lice is observed. As demonstrated by the available data, its prevalence reaches even 64.1% depending on the examined population (16,17).The symptoms in the current research pediculosis capitis was more common in girls than boys and long hair seems to be an important risk factor. Although long hair has been widely referred in the literature (18,19) even though, there are some authors who disagree (20)..While in other study in Houn city, (21.9%) were found infested with nits, immature or adult of pediculus humanus capitis (21). In Jordan the prevalence of pediculosis capitis was 26.6% (22). Al-Helys area19.5%, in Al-Tarya 35.6% and in Bo-Fakra 26.3% and this difference was statistically significant ( $p < 0.05$ )Nits was present in all male students and 87.2% of females at Al-Helys school, mixed was recorded in 12.8% of female students in the same school, these differences were not statistically significant ( $p = 0.0919$ ).. Nits was present among 88.9% of male students and 68.8% of females at Al-Tarya school, Mixed was recorded in 11.1% of male students and 31.3% of females in same school, these differences were not statistically significant ( $p = 0.407$ ).. In a similar study recorded that the infestation rate among girls varied from 26.07% (12 years group) to 55.89% (8 years group)(15).Nits was present in100% of male students and 76.2% of females in Bo -Fakra school, Mixed was recorded in 23.8% of female in same school, these difference was not statistically significant p value was 0. 0.639. Prevalence of pediculosis capitis in males was10.2% while in females was 41.8% , this difference was highly statistically significant ( $p = 0.0001$ ) the result was with agreement of other study the prevalence was significantly lower in boys (27.1%) than in girls (55.0%); ( $p < 0.0001$ ). (9) Also there was agreement with other study in Houn city, male children had a lower rate of infestation (6.27%) than females (38.66%) (6)..In study in Jordan there were significant differences in the prevalence between girls [34.7%] and boys [19.6%] (22). The highest prevalence of pediculosis capitis was in age 11 years(42.9%) followed by age 10 year (36.6%), this difference was statistically significant ( $p = 0.0001$ ).In other study the infestation rate among school children was significantly varied from 51.8% in children aged 7 years to 27.5% in children aged 10 years ( $p = 0.0001$ )>In Houn study found that school children exhibited decreased in infestation rate with age groups, the children aged 7-8 years were the most frequently affected (6).A study in Iran recorded that the children aged 10-11 years were the most frequently affected (23). In another study there was significant difference between age ( $p < 0.001$ )(22)..Evidence by cases of transmission between siblings recorded in 68.5% of the surveyed schools. Having siblings as one of the factors increasing the probability of P. humanus capitis infestation has been indicated by other authors as well (24-26). Dissemination of head louse is facilitated by staying in large groups of people.

Our analysis has shown that pediculosis capitis is more frequent in schools with greater numbers of children . This was similar to study conducted in Poland (27)..There were no effect of fathers occupation in having pediculosis capitis ( $p=0.419$ ).Also there was no effect of fathers level of education ( $p=0.697$ ). Also pediculosis was not effected with mothers occupation or education level ( $p=0.320$  and  $0.740$ ) respectively .In other study there was a significant relationship between head louse infestation, family income and parents education level (24)..Prevalence of Pediculosis capitis was seen in 17.4% in family size  $\leq 5$  persons and 23.9% in family size  $5 <$  person, although there was no significant difference ( $p=0.255$ ). In study in Jordan found that there was significant difference for family size and income ( $p < 0.001$ ) (22).

## **Conclusion**

Pediculosis capitis constitute a major health problem in school children in the three study villages (Alhlys, Altrya and Abofkra).The present study revealed that, the prevalence of pediculosis was lower than the other studies particularly those done in Libya. Girls have significantly higher rate of pediculosis capitis infestation than boys and socio-demographic factors including parents education, jobs, family size, number of rooms have not been shown to play a significant role in overspread or increased the infestation.

## **CONSENT**

Informed consent form all participant parents were taken through the three school administrations before starting the study.

## **ETHICAL APPROVAL**

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

## **References**

- 1) Tohit NFM, Rampal L, Mun-Sann L. Prevalence and predictors of pediculosis capitis among primary school children in Hulu Langat, Selangor. *Med J Malaysia*. 2017;12(1):72-17
- 2) Dasetima DA, Obinna CE, Bliss MM, and et al . Pattern of Skin Disorder among School Children in Port-Harcourt, Rivers State. *Asian Journal of Research in Dermatological Science*. 2020;14-24.
- 3) Rukke BA, Birkemoe T, Soleng A, and et al . Head lice prevalence among households in Norway: importance of spatial variables and individual and household characteristics. *Parasitology*. 2022. 138:1296–1304
- 4) Ebrahim H. Infestation of head lice, pediculus humanus capitis, in primary school children at Houn City, Libya. *J Acad Res*. 2019;13:38-52.
- 5) Rassami W, Soonwera M. Epidemiology of pediculosis capitis among school children in the eastern area of Bangkok, Thailand. *Asian Pac J Trop Biomed*. 2012;2(11):901-4.

- 6) Center for Disease Control and Prevention. Biology of head lice. Clifton Road: U.S. Department of Health & Human Services; 2015
- 7) Doroodgar A, Sadr F, Paksa A, and et al. The prevalence of pediculosis capitis and 1 relevant factors in primary school students of Kashan, Central Iran. *Asian Pac J Trop Dis.* 2014; 4 (6):500-4.
- 8) Szymanek M, Wojnowska D, Krasowska D. Pediculosis still an up to date clinical problem. *Przegląd Lekarski.* 2009; 66: 206- 8.
- 9) Hasan I & Hind M. Prevalence and associated factors of *Pediculus humanus capitis* infestation among primary school children in Sebha, Libya. *Journal of Pure & Applied Sciences.* 2020;19 . 5:132-38
- 10) McCue JD & Kahan S. In a Page Infectious Disease. 1 st Edition, Lippincott Williams & Wilkins: USA, 2006
- 11) Mimouni D, Ankol OE, Gdalevich M, and et al. Seasonality trends of pediculosis capitis and *Phthirus pubis* in a young adult population: follow-up of 20 years. *Journal of the European Academic Dermatology and Venereology.* 2002; 16: 257-9.
- 12) Rafie A, Kasiri H, Mohammadi Z, and et al. Pediculosis capitis and its associated factors in girl primary school children in Ahvaz City in 2005 -2006. *Iran J Infect Dis Trop Med* (2009);45:41-5.
- 13) Al-Maktari MT. Head louse infestations in Yemen: Prevalence and risk factors determination among primary school children, Al-Mahweet Governorate, Yemen. *Journal of Egyptian Society of Parasitology.* 2008; 38: 741-8.
- 14) Toloza A, Vassena C, Gallardo A, and et al. Epidemiology of Pediculosis capitis in elementary schools of Buenos Aires: Argentina. *Parasitology Research.* 2009; 104: 1295-8
- 15) Watcharawit R , Mayura S. Epidemiology of pediculosis capitis among school children in the eastern area of Bangkok, Thailand. *Asian Pac J Trop Biomed* 2012; 2(11): 901-904.
- 16) Costa DL, de Sousa DS, da Silva RM, and et al. Preventive Actions against *Pediculus capitis humanus* in Children of a Community Riverside's Metropolitan Region of Belém-Pará in Eastern Amazon. *International Journal of Tropical diseases & Health.* 2018: 7:1-3.
- 17) Feldmeier H, Jahnke C. Pediculosis capitis. *Epidemiologie, Diagnose und Therapie, Pädiatrische Praxis.* 2020. 76:359–370
- 18) Feldmeier H. Treatment of pediculosis capitis: a critical appraisal of the current literature. *Am J Clin Dermatol.* 2014;15(5):401-12.
- 19) Rassami W, Soonwera M. Epidemiology of pediculosis capitis among school children in the eastern area of Bangkok, Thailand. *Asian Pac J Trop Biomed.* 2012;2(11):901-4
- 20) Mumcuoglu KY, Meinking TA, Burkhart CN, and et al. Head louse infestations: the “no nit” policy and its consequences. *Int J Dermatol* 2006; 45(8):891–6.

- 21) Hana E. Infestation of Head Lice, *Pediculus humanus capitis*, in Primary School Children at Houn City, Libya. *Journal of Academic Research*, Issue 13 January 2019.38-52
- 22) Al Bashtawy, M & .Hasna, F. Pediculosis capitis among primary-school children in Mafraq Governorate, Jordan. *EMHJ - Eastern Mediterranean Health Journal*. 2012. 18 (1): 43-48.,
- 23) Drali R, Shako JC, Davoust B, and et al. A New Clade of African Body and Head Lice Infected by *Bartonella quintana* and *Yersinia pestis*-Democratic Republic of the Congo. *Ame J Trop Med Hyg*. 2015;93(5):990-3.
- 24) Willems, S.; Lapeere, H.; Haedens, N.; and et al. The importance of socio-economic status and individual characteristics on the prevalence of head lice in schoolchildren. *Eur. J. Dermatol*. 2005, 15, 387–392.
- 25) Dehghanzadeh, R.; Asghari-Jafarabadi, M.; Salimian, S.; and et al. Impact of family ownerships, individual hygiene, and residential environments on the prevalence of pediculosis capitis among schoolchildren in urban and rural areas of northwest of Iran. *Parasitol. Res*. 2015; 114: 4303 - 4295.
- 26) Birkemoe, T.; Lindstedt, H.H.; Ottesen, P and et al .Head lice predictors and infestation dynamics among primary school children in Norway. *Fam. Pract*. 2015, 33, 23–29.
- 27) Bartosik K, Janczaruk M, Zajac Z, and et al. Head Lice Infestation in Schoolchildren, in Poland—Is There a Chance for Change?. *Journal of Clinical Medicine*. 2022; 11 (3): 783




Fig. 1: Percentage of infested students with pediculosis capitis

Figure.2: Diagnosis of pediculosis capitis : Nits attached to hair shaft



Figure.3: Diagnosis of pediculosis capitis: Live lice

Table 1: Distribution of students pediculosis capitis  
In relation to school names

Name of school	Pediculosis capitis			
	Yes		No	
	No	%	No	%
Al-Helys	43	19.5	178	80.5
Al-Tarya	57	35.6	103	64.4\
Bo-Fakra	46	26.3	129	73.7
Total	146	26.3	410	73.7

$$X^2 = 12.529=2, P= 0.002$$

Table.2: Distribution of students pediculosis capitis in relation to sex

Sex	Pediculosis capitis			
	Yes		No	
	No	%	No	%
Male	28	10.2	246	89.8
Female	118	41.8	164	58.2
Total	146	26.3	410	73.7

$$X^2 = 71.779 \text{ df}=1, p= 0.0001$$

Table 3 : Distribution of students pediculosis capitis in relation to age

Age /year		Pediculosis capitis		
		Yes	No	
	No	%	No	%
6	0	0	2	100
7	18	22.8	61	77.2
8	14	13.7	88	86.3
9	23	24.7	70	75.3
10	34	36.6	59	63.4
11	36	42.9	48	57.1
12	21	20.8	80	79.2
13	0	0	2	100
Total	146	26.3	410	73.7

$$X^2 =28.909 \text{ df}=7, p= 0.0001$$

Table 4: Distribution of students pediculosis capitis in relation to fathers education

Fathers level of education		Pediculosis capitis		
		Yes	No	
	No	%	No	%
Illiterate	0	0	3	100
Primary	10	29.4	24	70.6
Preparatory	20	20.2	79	79.8
Secondary	21	23.6	68	76.4
University and above	17	21.5	62	78.5
Total	68	22.4	236	77.6

$$X^2 = 2.213 \text{ df}=4, p= 0.697$$

Table 5: Distribution of students pediculosis capitis in relation to mothers occupation

Mothers occupation		Pediculosis capitis		
		Yes	No	
	No	%	No	%
House wife	44	23.8	141	76.2
Teacher	18	28.1	46	71.9
Employee	1	5.9	16	94.1
Doctor	0	0	5	100
Nurse	2	16.7	10	83.3
Student	1	50	1	50
Lab. technician	0	0	2	100
Total	66	23	221	77

$$X^2 = 7.013 \text{ df}=6, p= 0.320$$

Table 6: Distribution of students pediculosis capitis in relation to family size

Family size		Pediculosis capitis			
		Yes		No	
	No	%	No	%	
≤ 5	12	17.4	57	82.6	
>5	58	23.9	185	76.1	
Total	70	100	242	100	

$$X^2 = 1.296 \text{ df}=1, p= 0.255$$

Table 7: Distribution of students pediculosis capitis in relation to number of person in each bedroom

Number of person in each bedroom	Pediculosis capitis			
	Yes		No	
	No	%	No	%
One	2	16.7	10	83.3
Two	19	19	81	81
Three	21	21.9	75	78.1
Four	15	27.8	39	72.2
Five	8	34.8	15	65.2
Seven	0	0	1	100
Eight	1	50	1	50
Total	66	22.9	222	77.1

$$X^2 = 4.876 \text{ df}=6, p= 0.560$$

