

# **Original Research Article**

## **CONTRACEPTIVE USE AMONG STUDENTS OF A HEALTH TRAINING INSTITUTION IN THE NORTH-EAST REGION OF GHANA**

### **Abstract**

#### **Background**

Despite contraceptive awareness and availability to the general population, including higher education institutions, unplanned pregnancies continue to increase.

#### **Aim/objective**

The study assessed contraceptive usage among female health professional students in the North-East Region, Ghana.

#### **Method**

A descriptive cross-sectional design was used to assess contraceptive use among 143 female health professional students. The data was collected using an online survey, analysed descriptively, Chi-square test and logistic regression analysis were performed using Statistical Package for Social Sciences version 25. A p-value of  $<0.05$  was considered statistically significant.

#### **Results**

The age ranged from 18 and 36 years, with a mean age of 23.7 (Sd±2.82) years, and the majority (69.9%) were in the 21-25 years age category. The mean age of first sexual intercourse was 17.8

(Sd  $\pm 5.65$ ) years. The overall prevalence of contraceptive use was 59.0%, with ever and current users being 47.6% and 35.7% respectively. The male condom was a commonly used contraceptive (53.6%). Using the Chi-square test, we found significant differences between current contraceptive use and age at first sex ( $X^2=15.42, p=0.009$ ), protection during first sexual intercourse ( $X^2=7.30, p=0.026$ ), current relationship status ( $X^2=15.12, p<0.001$ ), and multiple sexual partners ( $X^2=10.25, p=0.001$ ). Being in a sexual relationship (OR=4.49, 95%CI: 1.61, 12.50,  $p=0.004$ ) was positively associated with current contraceptive use. Prevention of unwanted pregnancy/birth was the common reason for contraceptive use, with fear of perceived side effects as the reason for non-use.

### **Conclusion/Interpretation**

Overall, the study found high contraceptive use with significant discontinuation among ever and current users. There is a need for public health education and promotion programmes to increase the acceptance and use of contraceptives among the study population and the public.

**Keywords:** Modern Contraceptives, Sexual intercourse, Students, Pregnancy, Higher institutions, and Ghana.

### **What is known**

- Contraceptives reduce pregnancy-related morbidity and mortality
- Contraceptives reduce unwanted pregnancies
- Contraceptives benefit the education of women and empower them for higher opportunities

### **What is added**

- Early initiation of sexual activities

- Substantial use of contraceptives
- Significant discontinuation of ever users and current users of contraceptive
- Barriers to contraceptive use correspond with reports of other studies

#### What are the Global and Public Health Policy Implications?

- Teenagers should be considered in contraceptive interventional programmes.
- Health promotional activities on contraceptive usage should be intensified, especially among health professionals and students.
- Barriers to contraception should be of global and national concern

## Introduction

Contraceptives are key public health interventions to address unmet needs for family planning and meeting the target of ensuring gender equality and women empowerment of the Sustainable Development Goal five (SDG-5)[1]–[4]. Universally, contraceptives use is reported to be increasing. However, there is a decline among Sub-Saharan African countries [4]. Regional variations in contraceptive use exist in Ghana. However, some previous studies reported prevalence rates of 21.5%, 31.0% in 2014 and 2018 in national representative data, and 18.5% in the Northern Region of Ghana in 2018 [4], [5].

Similarly, the estimated unmet need for birth spacing in the Northern Region is reported to be the highest (21.7%) in Ghana [5]. Ghana faces increasing medico-social challenges, including teenage pregnancy, unsafe abortion, unwanted pregnancies and sexually transmitted infections such as HIV/AIDs [6]–[12] due to the limited use of contraceptives. There is generally a global increase in the incidence of unplanned pregnancies among female students in higher learning institutions [13], potentially adversely impacting students on learning outcomes, high dropout rates and unattained goals among women [14], [15]. Addressing these challenges through the use of contraceptives is highly recommended [2], [4], [16]–[18] to empower women and address the unmet needs of family planning. Therefore, a conscious effort to promote contraceptive use, especially in women of reproductive age, is necessary. This study investigated contraceptives use among female health professional students in the North-East Region of Ghana. Health professionals play a critical role in the dissemination of advances in healthcare and medicine, including contraceptive use. Understanding contraceptive use and associated factors in health professional students who would soon transition to practice are essential to encourage utilisation among other populations. The North-East Region was recently created out of the Northern

Region of Ghana. There has been no reference to contraceptive use combined with associated factors conducted in the region. The study provides evidence on the significance of contraceptive use for addressing the region's unmet needs for family planning.

## **Materials and Method**

### **Study setting and design**

The study was conducted at the only tertiary health training institution, the College of Nursing and Allied Health Sciences, Nalerigu, in the North-East Region of Ghana. The North-East Region is among Ghana's six newly created regions, with Nalerigu as the regional capital. The College trains students in programmes including nursing, midwifery, medical laboratory, nutrition and dietetics, and physiotherapy. A cross-sectional design was used applied in the study. The study was conducted between September and October 2021.

### **Sampling technique**

A purposively institution-based online survey was used to collect data on contraceptive use among female health professional students of the College of Nursing and Allied Health Sciences, Nalerigu. We employed the online survey to reduce physical contact with participants as the study was conducted in the COVID-19 pandemic era. The online survey allowed participants to be reached remotely, and responses were received to meet the study's objective. A total of 143 female health professional students attempted the questionnaire out of 3095 (College of Nursing and Allied Health Sciences, 2021). A link to the online survey questions was shared via female health professional student's means of contact, including email addresses and mobile phone numbers.

Before participation, study participants were asked to indicate their willingness to participate after explaining the study's rationale. Participants who stated unwillingness to participate were automatically excluded from the study. Study participants who consented to participate continued to attempt items in the survey questionnaire.

### Inclusion and Exclusion Criteria

Female students qualified to participate in the study if they were 18 years and older and pursuing any academic programme at the College of Nursing and Allied Health Sciences in Nalerigu. Students who were below 18 years of age and male students were excluded from this study

### Study instrument and data collection

A modified survey questionnaire on contraceptive usage from previous studies was used to collect the data [4], [17]–[19]. An English version of the electronic google form questionnaire platform was created. Independent assessors from the Department of Population and Health, School of Public Health, University for Development Studies, Tamale Campus, reviewed the questionnaire. Subsequently, a pilot study was conducted on 25 female students at a nursing training college in the Northern Region, Ghana, to assess the reliability of the questionnaire. Modifications were made where necessary to the questionnaire. The questionnaire was designed to capture data on sociodemographic characteristics, sexual history, contraceptive use, and factors associated with contraceptives use. Data quality was maintained by designing the questionnaire to allow participants to complete the form before submission. The design was done to exclude incomplete forms. The questionnaire platform also allowed only one submission by a participant

### Data analysis and presentation

The online data were extracted and converted to Microsoft Excel File (version 16) and cross-checked for distortions and accuracy. The data were transferred onto Statistical Package for Social Sciences (SPSS) (version 25) for further data management and analysis. The string data were categorised into numeric data. The outcome variable used in this study was current contraceptives use and was a dichotomous variable, "yes (1) or no" (2). The explanatory variables were age at first sexual intercourse, protected first sexual intercourse, sexual relationship status, and multiple sexual partners. The data was analysed descriptively, and chi-square and logistic regression were performed. A p-value < 0.05 was considered statistically significant at 95% confidence interval. The data summary was presented in frequency, percentages, charts, and tables.

## **Results**

### **Sociodemographic characteristics**

The study consisted of 143 female participants between 18 and 36 years. The mean age was 23.7 (Sd±2.82) years, and the majority (69.9%) were in the 21-25 age category. The majority (65.0%) of the participants were in the first year of training, as shown in **Table 1**. A higher proportion of the participants were single (85.3%), Christians (67.1%) and received a monthly income of GHC ≤ 500.0 (55.2%).

**Table 1. Sociodemographic characteristics of participants**

Variables	Categories	Frequency (n= 143)	Percentage (%)
Age	≤20	14	9.8
	21-25	100	69.9
	26-30	25	17.5
	30≥	4	2.8
Marital Status	Co-habiting	3	2.1
	Married	18	12.6
	Single	122	85.3
Religion	Christian	96	67.1
	Muslim	45	31.5
	Traditionalist	2	1.4
Monthly income (GHC)	>500.00	64	44.8
	≤500.00	79	55.2
Year	First-year	93	65.0
	Second-year	47	32.9
	Third-year	3	2.1

*GHC; Ghana Cedis (Currency), Sd; Standard deviation*

### Sexual history and first contraceptive use among participants

The mean age of first sexual intercourse was 17.8 (Sd ±5.65) years, with the majority (61.5%) being 16-20 years. A high proportion (62.9%) of the participants indicated that they protected their first sexual intercourse. Common contraceptives used during first sexual intercourse included the male condom (32.9%), emergency contraceptive (9.8%) and contraceptive pills (9.1%). Most participants indicated that they were currently in a sexual relationship (67.8%). Of this figure, a low proportion has multiple sexual partners (19.6%), as shown in **Table 2**.

**Table 2. Sexual history and first contraceptive use among study participants**

Variables	Categories	Frequency	Percentage (%)
<b>Age at first sexual intercourse</b> (years)	≤10	4	2.8
	11-15	8	5.6
	16-20	88	61.5
	21-25	32	22.4
	26≥	2	1.4
	None	9	6.3
<b>Mean age (years) at first sexual intercourse</b>	17.80 (Sd ±5.65)		
<b>Protection during first sexual intercourse</b>	Yes	90	62.9
	No	43	30.1
	None	10	7.0
<b>Contraceptive used for first sexual intercourse (n=90)</b>	Contraceptive pills	13	9.1
	Emergency contraceptives	14	9.8
	Female condom	2	1.4
	Implant	1	0.7
	Injection	1	0.7
	Male condom	47	32.9
	Withdrawal method	12	8.4
<b>Current relationship status</b>	In sexual relationship	97	67.8
	Not in a sexual relationship	46	32.2
<b>Multiple sexual relationship (n=97)</b>	Yes	19	19.6
	No	78	80.4

*Sd; Standard deviation*

### **Prevalence of contraceptive use among study participants**

The prevalence of contraceptive use (ever and current users) is high among the study participants (59.0%), as shown in **Figure 1**.

### **Figure 1. Prevalence of contraceptive use among participants**

#### **Pattern and usage of contraceptives among participants**

The number of used and current contraceptive users was 47.6% and 35.7%, respectively. Current contraceptive users who have multiple **sexual** partners were 68.5%. The male condom (53.6%), emergency contraceptive (52.4%), and contraceptive pills (35.7%) were the most commonly used contraceptives. The reasons accounting for contraceptive use included prevention of unwanted pregnancy/births (78.4%), prevention of sexually transmitted infections (33.3%) and partners' choice (15.5%). The major reason for the non-use of contraceptives is fear of associated **perceived** side effects (37.0%), as shown in **Tables 3a and 3b**.

**Table 3a. Pattern and usage of contraceptives among participants**

<b>Variables</b>	<b>Categories</b>	<b>Frequency</b>	<b>Percentage (%)</b>
Ever contraceptive users	Yes	68	47.6
	No	75	52.4
Current contraceptive users	Yes	51	35.7
	No	92	64.3
Current contraceptive users who have multiple sex partners (n=19)	Yes	13	68.4
	No	6	31.1
Age distribution of current contraceptive users (n=51)	≤20	5	9.8
	21-25	35	68.6
	26-30	9	17.6
	31≥	2	4.0
Marital status and contraceptive use (n=51)	Co-habiting	0	0.0
	Married	5	9.8
	Single	46	90.2
Commonly used contraceptives by both current and previous users (Multiple response types of the question) (n=84)			
Male condom	Yes	45	53.6
	No	39	46.4
Female condom	Yes	5	6.0
	No	79	94.0
Emergency contraceptives	Yes	44	52.4
	No	40	47.6
Contraceptive pills	Yes	30	35.7
	No	54	64.3
Withdrawal method	Yes	18	21.4
	No	66	78.6
Injection	Yes	10	12.0
	No	74	88.0
Diaphragm	Yes	1	1.2
	No	83	98.8
Rhythm method	Yes	2	2.4
	No	82	97.6
Implant	Yes	4	4.8
	No	80	95.2
Intrauterine Device (IUD)	Yes	1	1.2
	No	83	98.8

**Table 3b. Continuation: Pattern and usage of contraceptives among participants**

<b>Variable</b>	<b>Categories</b>	<b>Frequency</b>	<b>Percentage (%)</b>
Reasons for using contraceptives among current users (n=51)			
Easily available	Yes	10	20.0
	No	41	80.0
Easy use	Yes	9	17.6
	No	42	82.4
Partner's choice	Yes	13	15.5
	No	38	84.5
Less expensive	Yes	7	13.7
	No	44	86.3
Spacing of birth	Yes	3	3.6
	No	48	96.4
Prevention of unwanted pregnancy/birth	Yes	40	78.4
	No	11	21.6
Prevention of sexually transmitted infections	Yes	17	33.3
	No	34	66.7
Reasons for not using contraceptives among current non-users (n=92)			
Bad perception of society toward its uses	Yes	5	5.4
	No	87	94.6
I am a Christian, and therefore I am forbidden to use	Yes	9	9.8
	No	83	90.2
I am a Muslim, and therefore I am forbidden to use	Yes	2	2.2
	No	90	97.8
It is not readily available	Yes	0	0.0
	No	92	100.0
Fear of side effects	Yes	34	37.0
	No	58	63.0
Not sexually active	Yes	9	9.8
	No	83	90.2
My partner does not allow me to use	Yes	9	9.8
	No	83	90.2
Stigmatization	Yes	4	4.3
	No	88	95.7
Costly	Yes	5	5.4
	No	87	94.6

### Association between sexual history and current contraceptive use of participants

A chi-square was performed to determine the association level between the dependent variable (current contraceptive use) and the independent variables, including sociodemographic characteristics and the sexual history of study participants. The Chi-square test showed significant differences between age at first sexual intercourse ( $X^2=15.42, P=0.009$ ), protection during first sexual intercourse ( $X^2=7.30, P=0.026$ ), and current relationship status ( $X^2=15.12, P<0.001$ ) and multiple sexual partners ( $X^2=10.25, P=0.001$ ) as shown in **Table 4**.

**Table 4. Association between sexual history and current contraceptive use of participants**

Variables	Categories	Current contraceptive use		p-value ( $X^2, df.$ )
		Yes	No	
Age at first sex (years)				0.009 (15.42, 5)
	≤10	2	2	
	11-15	2	6	
	16-20	41	47	
	21-25	6	26	
	26≥	0	2	
	None	0	9	
Protection during first sexual intercourse				0.026(7.30, 2)
	Yes	21	42	
	No	30	40	
	None	0	10	
Current relationship status				<0.001 (15.12,1)
	In a sexual relationship	45	52	
	Not in a sexual relationship	6	40	
Multiple sexual partners				0.001(10.25,1)
	Yes	13	6	
	No	38	86	

## Predictors of contraceptive use among study participants

We performed a logistic regression analysis to determine the association between independent and dependent variables of current contraceptive use. We found that being in a sexual relationship (OR= 4.49, 95%CI (1.610-12.498),  $p=0.004$ ) was positively associated with contraceptive use among study participants, as shown in **Table 5**.

Table 5. Predictors of contraceptive use among participants

Variables	P-Value	OR (95%CI)
<b>Protection during first sexual intercourse</b>		
Yes	0.894	0.89 (0.159-4.984)
No	0.376	0.44 (0.070-2.734)
None (RC)		-
<b>Relationship status</b>		
In a sexual relationship	0.004	4.49 (1.610-12.498)
Not in a sexual relationship (RC)		-
<b>Multiple sexual partners</b>		
Yes	0.119	2.67 (0.777-9.195)
No (RC)		-

*RC; Reference Category, OR; Odds Ratio*

## Discussion

Our study provides the findings on institutional-based contraceptive use in the North-East Region. We found the overall contraceptive use to be high (59.0%), with relatively higher proportions of ever (47.6%) and current users (35.7%). Similar to **previous** studies, contraceptive use was reported to be high among health professional students in Ghana [20], female tertiary students in Botswana [21] and low among other tertiary students in Ghana [13]. The results in our study are **significant** considering our participants, who are health professional students. Their **contraceptive usage** may affect the type of education and recommendation they offer to the public **and potential users**. The high contraceptive use among participants may be partly due to their health background, **making them well informed about the important** benefits of contraceptives. Comparatively, there was significant discontinuation of contraceptive use among ever and current users. The discontinuation raises a **significant** public health challenge, **considering** recent reports of increasing unplanned pregnancies among tertiary students globally [13], [22]. Understanding the factors accounting for the discontinuation among contraceptive users **offers essential** data for public health education and promotion activities and **for preventing** unplanned pregnancies. Among the reasons **reported** by participants for non-use of contraceptives included; fear of **perceived** side effects, partners not encouraging contraceptive use and **a wrong** perception of society towards contraceptive use. These reasons have been equally identified by other researchers [5], [23], [24]. These reasons should be addressed if Ghana and the subregion are to meet **their** family planning needs. The enablers of contraceptive use among current users **provide** public health experts in family planning services **with** valuable information to **address** unmet family planning needs. Unavailability of contraceptives was not cited as a reason for **the** non-use of contraceptives among non-users; this was confirmed by

current users indicating that contraceptives were readily available. The latter may highlight Ghana's progress towards achieving universal family planning and the SDGs 3 and 5 by 2030. Our study found that partners can influence the use or disuse of contraceptives. Therefore, contraceptive use interventions should be bidirectional, targeting females and males to achieve optimum acceptance and use.

Teenagers should be a target for contraceptive use to reduce sexually transmitted infections, including HIV/AIDS, teenage pregnancies, and attendant consequences. Most participants engaged in sexual activities during their adolescent period, and it was high among persons within the age groups of 16-20 years. Other recent studies in the northern region of Ghana reported sexual activities among 16 years and above [25]–[30]. Important in this study is the finding of early use of contraceptives among our study participants, re-emphasising a need to target teenagers in family planning services in Ghana. The knowledge and use of contraceptives among senior high school students were relatively low in a northern region study [23]; such individuals are typically teenagers. Our findings corroborate the findings of Abdul-wahab et al. [25], knowledge, attitudes and practices of sexual and reproductive health among teenagers were poor in the Tamale Metropolis, Ghana. These findings stress the need to intensify the efforts in addressing the unmet needs of family planning, including knowledge and the use of contraceptives. More significantly, public health interventions on contraceptive use should consider factors such as age at first sexual intercourse, protection during first sexual intercourse, current relationship status and persons with multiple sexual partners.

### **Study limitation**

The data used in this study was self-reported and subjected to information biases combined with our inability to confirm the responses. The study was limited to only females and may not

express **males' perspectives** on contraceptive use. However, our study provides essential information on contraceptive use among significant subpopulations as healthcare students who would soon transition to practice. Understanding contraceptives use provides vital information that could be factored into health education and promotion programmes at the various healthcare facilities to increase contraceptive uptake and prevent unwanted pregnancies among tertiary students.

## **Conclusion**

Overall, the study found high contraceptives use and a relatively higher proportion of current contraceptive users. Comparatively, contraceptive dropout was significant among ever users and existing users. An essential reason for discontinuation of contraceptive use included fear of side effects which can be relied on for public health education and promotion programmes. **Educating the public on contraceptives** should be re-**emphasised** to increase acceptance and use.

## **Ethical Approval**

**The Research and Innovation Unit** of the College of Nursing and Allied Health **Sciences**, Nalerigu, **approved the study** (MOH/NMTC/RES21-9/003).

## **Consent**

As per international standard or university standard, Participants' written consent has been collected and preserved by the author(s).

## **References**

- [1] V. Kantorová, M. C. Wheldon, A. N. Z. Dasgupta, P. Ueffing, and H. C. Castanheira, "Contraceptive use and needs among adolescent women aged 15-19: Regional and global estimates and projections from 1990 to 2030 from a Bayesian hierarchical modelling study," *PLoS One*, vol. 16, no. 3 March, pp. 1–28, 2021, doi: 10.1371/journal.pone.0247479.
- [2] N. Cahill *et al.*, "Modern contraceptive use, unmet need, and demand satisfied among women of reproductive age who are married or in a union in the focus countries of the Family Planning 2020 initiative: a systematic analysis using the Family Planning Estimation Tool," *Lancet*, vol. 391, no. 10123, pp. 870–882, 2018, doi: 10.1016/S0140-6736(17)33104-5.
- [3] U. Women, "Gender equality and women's empowerment – United Nations Sustainable Development," *Online*. 2020, [Online]. Available: <https://www.un.org/sustainabledevelopment/gender-equality/>.
- [4] P. A. Aviisah *et al.*, "Modern contraceptive use among women of reproductive age in Ghana: Analysis of the 2003-2014 Ghana Demographic and Health Surveys," *BMC Women's Health*, vol. 18, no. 1, pp. 1–10, 2018, doi: 10.1186/s12905-018-0634-9.
- [5] A. A. E. Biney *et al.*, "Being ready, willing and able: understanding the dynamics of family planning decision-making through community-based group discussions in the Northern Region, Ghana," *Genus*, vol. 77, no. 1, pp. 1–23, 2021, doi: 10.1186/s41118-020-00110-6.
- [6] K. Atuahene, "Current status of HIV in Ghana," *13th Interes. Conf. 14th -17th May, 2019*, pp. 1–14, 2019, [Online]. Available: <http://regist2.virology->

education.com/presentations/2019/13INTEREST/04\_Atuahene.pdf Accessed 23/04/2020.

- [7] J. Abesig, Y. Chen, H. Wang, F. Mwekele, S. Irene, and X. Y. Wu, "Prevalence of viral hepatitis B in Ghana between 2015 and 2019: A systematic review and meta-analysis," *PLoS One*, vol. 15, no. 6, pp. 1–16, 2020, doi: 10.1371/journal.pone.0234348.
- [8] A. Y. Owusu, "A gendered analysis of living with HIV/AIDS in the Eastern Region of Ghana," *BMC Public Health*, vol. 20, no. 1, pp. 1–15, 2020, doi: 10.1186/s12889-020-08702-9.
- [9] C. B. Polis *et al.*, "Estimating the incidence of abortion: Using the Abortion Incidence Complications Methodology in Ghana, 2017," *BMJ Glob. Heal.*, vol. 5, no. 4, 2020, doi: 10.1136/bmjgh-2019-002130.
- [10] A. S. Yussif, A. Lassey, G. Y. K. Ganyaglo, E. J. Kantelhardt, and H. Kielstein, "The long-term effects of adolescent pregnancies in a community in Northern Ghana on subsequent pregnancies and births of the young mothers," *Reprod. Health*, vol. 14, no. 1, pp. 1–7, 2017, doi: 10.1186/s12978-017-0443-x.
- [11] C. A. Atakro *et al.*, "Contributing factors to unsafe abortion practices among women of reproductive age at selected district hospitals in the Ashanti region of Ghana," *BMC Womens. Health*, vol. 19, no. 1, pp. 1–17, 2019, doi: 10.1186/s12905-019-0759-5.
- [12] S. H. Nyarko, "Unintended Pregnancy among Pregnant Women in Ghana: Prevalence and Predictors," *J. Pregnancy*, vol. 2019, 2019, doi: 10.1155/2019/2920491.
- [13] F. Y. Gbagbo, "Family planning among undergraduate university students : a CASE study of a public university in Ghana," vol. 3, pp. 1–9, 2019.

- [14] A. A. E. Biney and P. Nyarko, "Is a woman's first pregnancy outcome related to her years of schooling? An assessment of women's adolescent pregnancy outcomes and subsequent educational attainment in Ghana," *Reprod. Health*, vol. 14, no. 1, pp. 1–15, 2017, doi: 10.1186/s12978-017-0378-2.
- [15] R. Mohr, J. Carbajal, and B. B. Sharma, "The Influence of Educational Attainment on Teenage Pregnancy in Low-Income Countries: A Systematic Literature Review," *J. Soc. Work Glob. Community*, vol. 4, no. 1, pp. 19–31, 2019, doi: 10.5590/jswgc.2019.04.1.02.
- [16] K. Grindlay *et al.*, "Contraceptive use and unintended pregnancy among young women and men in Accra, Ghana," *PLoS One*, vol. 13, no. 8, pp. 1–13, 2019, doi: 10.1371/journal.pone.0201663.
- [17] P. A. Apanga, M. T. Kumbeni, E. A. Ayamga, M. B. Ulanja, and R. Akparibo, "Prevalence and factors associated with modern contraceptive use among women of reproductive age in 20 African countries: A large population-based study," *BMJ Open*, vol. 10, no. 9, 2020, doi: 10.1136/BMJ open-2020-041103.
- [18] P. Beson, R. Appiah, and A. Adomah-Afari, "Modern contraceptive use among reproductive-aged women in Ghana: Prevalence, predictors, and policy implications," *BMC Womens. Health*, vol. 18, no. 1, pp. 1–8, 2018, doi: 10.1186/s12905-018-0649-2.
- [19] J. Agyemang, S. Newton, I. Nkrumah, J. M. Tsoka-Gwegweni, and S. N. Cumber, "Contraceptive use and associated factors among sexually active female adolescents in Atwima Kwanwoma District, Ashanti region-Ghana," *Pan Afr. Med. J.*, vol. 32, pp. 1–13, 2019, doi: 10.11604/pamj.2019.32.182.15344.
- [20] E. Paul *et al.*, "Contraceptive Methods : The Level of Awareness and Patronage among

- Ghanaian Health Profession Trainees," vol. 5, no. 4, pp. 5609–5616, 2018, doi: 10.26717/BJSTR.2018.07.001435.
- [21] M. E. Hoque and T. Ntsipe, "Methodology Study design and setting," *J. Soc. Asp. HIV/AIDS*, vol. 0, no. 0, pp. 1–6, 2013, doi: 10.1080/17290376.2013.869649.
- [22] E. V. Maja TMM, "Contraceptive practices in northern Tshwane, Gauteng Province," *Heal. SA Gesondheid*, 2004.
- [23] F. M. Patricia, A. Nelsi, and E. K. Dzantor, "Assessing the Knowledge , Use and Perception of Contraceptives among Senior High School Students in the Kumbungu District of Northern Ghana," vol. 4, no. 4, pp. 1–12, 2021.
- [24] E. Kofuor and M. Darteh, "in Ghana," *J. Community Health*, no. March, 2016, doi: 10.1007/s10900-015-0057-6.
- [25] I. Abdul-wahab, A. M. Nungbaso, R. N. Nukpezah, and E. K. Dzantor, "Adolescents Sexual and Reproductive Health : A Survey of Knowledge , Attitudes and Practices in the Tamale Metropolis , Ghana," vol. 6, no. 1, pp. 31–47, 2021.
- [26] Q. Ma *et al.*, "Early initiation of sexual activity: A risk factor for sexually transmitted diseases, HIV infection, and unwanted pregnancy among university students in China," *BMC Public Health*, vol. 9, pp. 1–8, 2009, doi: 10.1186/1471-2458-9-111.
- [27] A. R. Alhassan, K. Abdulai, and M. A. Alhassan, "Early Sexual Debut among Ghanaian Women: Correlates and Psychological Effect," *Biomed Res. Int.*, vol. 2021, 2021, doi: 10.1155/2021/5838510.
- [28] E. Turi, B. T. Merga, G. Fekadu, and A. A. Abajobir, "Why too soon? Early initiation of

- sexual intercourse among adolescent females in Ethiopia: Evidence from 2016 ethiopian demographic and health survey," *Int. J. Womens. Health*, vol. 12, pp. 269–275, 2020, doi: 10.2147/IJWH.S244621.
- [29] R. Stephenson, C. Simon, and C. Finneran, "Community factors shaping early age at first sex among adolescents in Burkina Faso, Ghana, Malawi, and Uganda," *J. Heal. Popul. Nutr.*, vol. 32, no. 2, pp. 161–175, 2014, doi: 10.3329/jhpn.v32i2.2610.
- [30] E. Abebe, A. Addis, A. Asmamaw, A. Addisu, and B. Ayanaw, "Factors associated with early sexual initiation among preparatory and high school youths in Woldia town , northeast Ethiopia : a cross-sectional study," *BMC Public Health*, pp. 1–8, 2019, [Online]. Available: <https://bmcpublikealth.biomedcentral.com/articles/10.1186/s12889-019-6682->