

## Case study

# Study of Insulin Hormone Level and $\beta$ -Cell Function in Children with $\beta$ -Thalassemia Major

### Abstract

**Background:** Diabetes and Impaired Glucose Tolerance are a major complication of iron overload in patients with beta thalassemia major.

**Aim:** To study the serum level of Insulin hormone and beta cell function in  $\beta$ -Thalassemia major patients.

**Methods:** This study was carried out on 50 children that were divided into: **Group (1):** thirty child with beta thalassemia with age ranged between (5 to 10 years), **Group (2):** Twenty apparently healthy matched age children were also included as control group. Both groups were subjected to history taking, full clinical examination, anthropometric measures, laboratory investigation that included: complete blood picture, serum iron, serum ferritin, fasting serum glucose, 2h postprandial serum glucose, fasting serum insulin by ELISA (Sandwich assay procedure), IRI, BFI.

**Results:** There was highly statistical significant difference between the patient and control groups in the insulin resistance index (IRI), but there was decrease in beta cell function index (BFI) in patients group with no significant difference.

**Conclusion:** Thalassemic child is at risk of IR which is responsible for abnormalities in glucose metabolism so proper management and regular follow up is necessary to early diagnosis and prevention of complications.

**Key words:** thalassemia, diabetes mellitus, impaired glucose tolerance and oral glucose tolerance test

### Introduction

Thalassemia is inherited blood defect having lifelong implications in the life of the patients and their relatives. This hereditary condition results in defective production of the quaternary protein "haemoglobin-Hb". Hemoglobin, a component of red blood cells. Hb is made up of a pair of  $\alpha$  and  $\beta$  globin chains. Thus, the levels of both the globin chains should be equal (1).

Beta thalassemia is characterized by reduced or absent  $\beta$  globin chain synthesis. The gene encoding for  $\beta$  globin is located on chromosome 11. More than 200 point mutations and rarely some deletion can cause  $\beta$  thalassemia (2).

However, blood transfusions lead to chronic iron overload that frequently cause endocrine problems, especially glucose metabolism abnormalities including diabetes mellitus (DM) and impaired glucose tolerance (IGT) (3).

Levels of glucose in blood are regulated by two endocrine hormones secreted by the organ pancreas: insulin and glucagon. Insulin is required to lower the blood glucose levels. Hence, it helps in the breakdown of glucose. Glucagon on the other hand is increasing the blood glucose levels. Both the hormones work together and maintain the normal glucose concentration in the blood (3).

Accumulation of iron in the pancreas causes the defective insulin secretion and leads to insulin deficiency. However, other studies also showed the presence of hyperinsulinemia and IR (rather than insulin deficiency) suggesting the involvement of insulin resistance in producing impairment to glucose homeostasis. Hyperinsulinemia may result from reduced hepatic clearance, which may be associated to iron overload (4).

The causes for glucose metabolism abnormalities are not fully understood; IGT with hyperinsulinemia and reduced insulin

Small letters : [1ks] Comment

a major : [2ks] Comment

small letter : [3ks] Comment

into two groups : [4ks] Comment

children : [5ks] Comment

including : [6ks] Comment

ferritin : [7ks] Comment

delet : [8ks] Comment

insulin resistance index (IRI) and beta cell function index (BFI) : [9ks] Comment

delet : [10ks] Comment

patient's : [11ks] Comment

write either haemoglobin or hemoglobin in all your article : [12ks] Comment

sensitivity are indicative of insulin resistance in these patients, while insulin deficiency secondary to iron induced pancreatic  $\beta$ -cell defects are considered as another possible cause of glucose abnormalities (5).

**Aim of the work:**

The aim of this work was to evaluate  $\beta$ -cell function of pancreas and glucose metabolism abnormalities in  $\beta$ -Thalassemia major patients.

**Patients and Methods:**

**A-Patients:**

This study was conducted on 30 children with beta thalassemia major who were admitted to Hematology unit, Pediatric department, Tanta University Hospitals, and 20 healthy children of matched age and sex as a control group. This study was conducted after taking a consent from parents of studied patients and approval from ethical committee of research center in Tanta university hospitals.

**\* Inclusion Criteria:**

Children between (5-10) years old diagnosed with  $\beta$ -Thalassaemia by clinical features and hemoglobin electrophoresis who are receiving regular blood transfusion (over period of 12 month) to maintain Hb level not more than 9.5g /dl and receiving Desferoxamine as achelator by intravenous infusion or Exjade orally.

**\* Exclusion Criteria:**

1. Children with hepatitis B or C.
2. Family history of DM in 1<sup>st</sup> degree relatives.
3. Children who receiving medication known to cause glucose intolerance such as steroids.

**B- Methods:**

All children included in this study were subjected to:

**I-History taking with a special account on:**

- Age of patients.
- Age of diagnosis of  $\beta$ -thalassemia
- Age of the first blood transfusion.
- Duration of regular transfusion therapy.
- Age of start of chelation therapy.
- Type of chelation used.
- Dose and regularity of chelation therapy.

**2-Thorough clinical examination with a special account on:**

- Weight, height, span, upper and lower body segment.
  - Head and neck for: Pallor, Jaundice, mongoloid facies.
  - Abdominal examination for: hepatomegaly and splenomegaly.
- 3- Laboratory investigation including:**
- Complete blood count.
  - Serum iron.
  - Serum ferritin.
  - Fasting serum glucose and 2-h post prandial serum glucose.
  - Fasting serum insulin.

**Statistical analysis:** Results were analyzed using SPSS (ver. 25.0; IBM, Chicago, IL, USA). Quantitative data was displayed in the form of mean  $\pm$  standard deviation (SD). Qualitative data was demonstrated through figures of frequency and percentage. Charts were used to illustrate data and relations where appropriate and  $p < 0.05$  was accepted as indicating statistical significance.

**Results:**

This case control study was conducted in Hematology Unit, Pediatric Department, Tanta University Hospital after being approved by the department ethical Committee. The study population included 50 children who were divided into two groups:

**Control group:** This included twenty (20) healthy children of matched age and sex.

**Patient group :** This included thirty (30) children diagnosed with  $\beta$ -Thalassaemia since more than two years by clinical features and hemoglobin electrophoresis and receiving regular blood transfusion (over period of 12 month).

**Table (1): Demographic data of the Patient groups.**

	Patient group (n=30)	Control group (n=20)	Test value	P value
Age <sub>(years)</sub>	7.5 $\pm$ 1.83	7.5 $\pm$ 1.73	0.00	1.00 <sup>1</sup>
Gender	Male	19(63.3%)	0.347	0.556 <sup>2</sup>
	Female	11(36.7%)		
Family history of consanguinity	Positive	9(30%)	3.182	0.003 <sup>3</sup>
	Negative	21(70%)		
Height <sub>(cm)</sub>	124.95 $\pm$ 11.5	127.5 $\pm$ 16.8	0.598	0.553 <sup>1</sup>
Weight <sub>(kg)</sub>	31.7 $\pm$ 7.3	32.8 $\pm$ 7.4	0.481	0.632 <sup>1</sup>
BMI <sub>(kg/m<sup>2</sup>)</sub>	18.8 $\pm$ 2.5	20.4 $\pm$ 1.97	2.147	0.037 <sup>4</sup>

1. Abbreviations: BMI: body mass index.

This paragraph should be replaced by anthropometric measures of only weight and height to calculate body mass index (BMI), while the rest parameters should be deleted because there is no results presented about them

committee :[13ks]Comment

This paragraph transfer to the beginning of patients and methods section :[19ks]Comment

Patient's age at interview :[14ks]Comment

Patient's age at :[15ks]Comment

Patient's age at :[16ks]Comment

Patient's age at :[17ks]Comment

It is not necessary to use abbreviation meaning or number to explain the statistical test, also don't put the unit of parameters as subscript, and why family history result not appear in the text, so either delete it from table or explain what does its meaning :[20ks]Comment

2. Independent t-test used; 3. Chi-square test used; 4. Fisher exact test.

\*Statistical significant when p-value <0.05.

There was no statistically significant difference between studied groups as regard age and sex while there was significant lower BMI in patients group  $P > 0.05$  in comparison to control group table(1).

**Table (2): Comparison between studied groups as regard complete blood picture:**

	Patient group (n=30)	Control group (n=20)	Test value	p-value
<b>Hemoglobin (g/dl)</b> Range	7.3 ± 0.8 (5.6-10.2)	13.9 ± 1.4 (11.5-16)	5.73	<0.001* ↓
<b>MCV (fl)</b> Range	53.2 ± 2.5 (46-60)	72.4 ± 3.2 (62-83)	4.84	<0.001* ↓
<b>MCH (pg)</b> Range	18.8 ± 1.8 (12-25)	22.3 ± 1.2 (18-28)	3.53	<0.001* ↓
<b>MCHC (g%)</b> Range	35.2 ± 3.0 (26-44)	48.3 ± 2.7 (45-57)	2.98	<0.001* ↓
<b>WBCs (10<sup>3</sup>/μL)</b> Range	9.817 ± 1.765 (4-11)	10.514 ± 0.816 (4-11)	0.986	0.412 <sup>1</sup>
<b>Platelets (10<sup>3</sup>/μL)</b> Range	179.342 ± 23.645 (150-230)	218.752 ± 19.824 (160-280)	1.06	0.308 <sup>1</sup>

Independent t test

\* Statistical significant when p-value <0.05.

As regard Hb, MCV, MCH and MCHC were significantly lower among Patient group ( $p < 0.001$ ), while WBCs and platelets count were lower among Patient group but with statistically insignificant differences as  $p > 0.05$ .

**Table (3): comparison between studied groups as regard Liver and kidney function tests :**


	Patient group (n=30)	Control group (n=20)	Test value	p-value
<b>Creatinine (mg/dL)</b> Range	0.79 ± 0.15 (0.42-1.1)	0.70 ± 0.14 (0.40-1.0)	2.158	0.36 <sup>1</sup>
<b>BUN (mg/dL)</b> Range	14.53 ± 2.7 (7-22)	14.18 ± 3.05 (6.8-21.5)	0.427	0.671 <sup>1</sup>
<b>AST (U/L)</b> Range	48.4 ± 19.2 (39-110)	24.7 ± 3.12 (14-35)	5.473	<0.001* <sup>1</sup>
<b>ALT (U/L)</b> Range	68.7 ± 27.7 (42-115)	24.8 ± 3 (15-36)	7.032	<0.001* <sup>1</sup>

BUN: blood urea nitrogen; AST: aspartate aminotransferase; ALT: alanine aminotransferase;

Independent t test

\* Statistical significant when p-value <0.05.

$\beta$  -Thalassemia major patients had significantly higher levels of ALT (68.7 ± 27.7 vs. 24.8 ± 3;  $p < 0.001$ ), and AST (48.4 ± 19.2 vs. 24.7 ± 3.12;  $p < 0.001$ ) in comparison to the control group. fig(1).



**Figure (1): Liver enzymes among study groups.**

Leave this figure as :[23ks]Comment mentioned in table 3

**Table (4): Serum insulin levels among study groups**

	Patient group (n=30)	Control group (n=20)	Test value	p-value
<b>Plasma insulin (me/dL)</b> Range	10.23 ± 1.76 (7-14)	8.01 ± 0.68 (6-10)	5.572	<0.001* <sup>1</sup>
<b>FBS (mg/dL)</b> Range	97.6 ± 8.75 (81-119)	90.5 ± 5.37 (70-105)	3.216	0.002* <sup>1</sup>
<b>2hpp (mg/dL)</b> Range	97.2 ± 8.27 (80-116)	96.1 ± 16.4 (78-110)	0.326	0.031* <sup>1</sup>
<b>IRI</b> Range	2.73 ± 0.26 (2-3.5)	1.76 ± 0.14 (1.3-2)	17.39 9	<0.001* <sup>1</sup>
<b>BFI</b> Range	117.8 ± 11.8 (80-150)	123.5 ± 10.9 (90-155)	1.742	0.08 <sup>1</sup>
<b>OGTT</b>			6.156	<0.001* <sup>2</sup>
Normal	20(66.7%)	19(95%)		
IFG	7(23.3%)	1(5%)		
IGT	0(0%)	0(0%)		
DM	3(10%)	0(0%)		

Abbreviations: OGTT: oral glucose tolerance test; FBS: fasting blood sugar; 2hpp: 2-hour postprandial plasma glucose; IRI: insulin resistance test; BFI: b cell function index

Independent t test; 2.Fisher exact test.

\*Statistical significant when p-value <0.05.

As regard Patient group had significantly higher levels of FBS (97.6 ± 8.75 vs. 90.5 ± 5.37;  $p = 0.002$ ) in comparison to the control group. They also have a significantly higher insulin resistant index than the control group (2.73 ± 0.26 vs. 1.76 ± 0.14;  $p < 0.001$ ).

As regard BFI, was reduced in  $\beta$  -Thalassemia patients but the difference was not significant. The OGTT of the 30  $\beta$  -Thalassemia patients, showed that 10 (33.3%) had abnormal results, three (10%) were diabetic and 7 (23.3%) had IFG. Fig(2&3&4&5&6)

It was :[25ks]Comment

Data in this table include liver :[22ks]Comment function and kidney function tests which are not illustrated as a goal of your study, I prefer to leave these data and published in next article

It is not necessary because its :[26ks]Comment value is mentioned in table 4

**Figure (2): Serum insulin level among study groups**



Figure (3): FBS level among study groups




Figure (4): 2hpp level among study groups




Figure (5): IRI level among study groups




Figure (6): OGTT results among study groups

As regard, Patient group had significantly higher levels of serum iron and ferritin ( $p < 0.001$ ) with mean and standard deviation of  $160.8 \pm 22.12$  vs.  $79.6 \pm 11.24$ ,  $3008.03 \pm 279.2$  vs.  $69.8 \pm 34.3$ , respectively).

Table (6): Correlation between IRI, BFI and laboratory data

	IRI		BFI	
	r	p-value	r	p-value
BMI	-0.282	0.047*	-0.130	0.369
AST	0.523	<0.001*	0.339	0.139
ALT	0.637	<0.001*	0.330	0.818
Serum iron	0.806	<0.001*	0.254	0.075
Serum ferritin	0.906	<0.001*	0.230	0.108
IRI	---	---	0.807	0.035*
BFI	0.807	0.035*	---	---
Age at thalassemia diagnosis	0.092	0.322	0.113	0.298
Mean duration of transfusion	0.107	0.210	0.098	0.427
Mean age at start of chelation	0.084	0.427	0.121	0.335

Abbreviations: BMI: body mass index; AST: aspartate aminotransferase; ALT: alanine aminotransferase; IRI: insulin resistance test; BFI: b cell function index

As regard BFI had significant direct correlation with IRI. IRI had significant direct correlations with AST, ALT, serum iron and serum ferritin, while had significant negative correlation with BMI. Fig(7,8)




Figure (7): Correlation between IRI and BMI.

Table (5): Iron levels among study groups

	Patient group (n=30)	Control group (n=20)	Test value	p-value
Serum Iron	160.8±22.12	79.6±11.24	17.06	<0.001* <sup>1</sup>
Serum Ferritin	3008.03±279.2	69.8±34.3	57.01	<0.001* <sup>1</sup>

Independent t test

\* Statistical significant when p-value < 0.05.

It is not necessary because its value mentioned in table 4 :[27ks]Comment

It is not necessary because its value mentioned in table 4 :[28ks]Comment

Data of correlation with AST and ALT should be left from table and text :[32ks]Comment

It is not necessary because its value mentioned in table 4 :[29ks]Comment

It is not necessary because its value mentioned in table 6 :[33ks]Comment

It is not necessary because its value mentioned in table 4 :[30ks]Comment

The same comments as mentioned in table 1 :[31ks]Comment




Figure (8): Correlation IRI and serum ferritin.

## Discussion

$\beta$ -Thalassemia major is a serious genetic disease that is often treated with frequent blood transfusions (6). The use of regular blood transfusions in  $\beta$ -TM has improved the quality of life and increased survival of these patients; however, blood transfusions lead to chronic iron overload that frequently cause endocrine problems, especially glucose metabolism abnormalities including diabetes mellitus (DM) and impaired glucose tolerance (IGT) (7).

The aim of this work was to evaluate  $\beta$ -cell function, Insulin resistance and glucose metabolism abnormalities in  $\beta$ -Thalassemia major patients.

The current study was carried out on Thirty (30) child with packed RBCs transfusion dependent  $\beta$ -Thalassaemia Major who were admitted in Hematology Unit, Pediatric Department, Tanta University Hospital and

Twenty (20) healthy child of matched age and sex served as a control.

In the current study the mean age of thalassaemic children in our study was  $7.5.1 \pm 1.83$  years. Our study population had a relatively younger age group when compared to other studies, As in, **Chen, et al**(8) where the mean the cases were  $14.8 \pm 6.9$ .

In the current study male predominance was found between Patient groups males represent (63.3%) while female represent (36.7).

This is in agreement with **Islam et al** (9) with which reported a male to female ratio of 1.26.

Our result is not in agreement with a similar study carried out in Hong Kong that found that the male to female ratio was equal among their studied thalassaemic cases.(10)

In the current study it was found that BMI was significantly lower in Patient group than control group

This is in agreement with **Dey, Konwar, Sarkar**(11) who found that underweight (Low BMI) was a common finding in thalassaemia patients specially when they were older than 10 years of age.

The cause of under nutrition in thalassaemia major patients is multifactorial. Endocrinopathies including hypothyroidism, hypo-gonadotropic gonadism, growth hormone deficiency, hypo-parathyroidism are the major causes of under nutrition in these patients.(11).

In the current study we founded that mean age of start of chelation therapy was  $4.32 \pm 2.52$ .

This is in agreement with **Al-Kherbash, Al-Awdi, Hasan**(12) who found that earlier initiation of chelating therapy in 19.7% of the cases at age less than 3 years led to the best outcome of the cases, with no mortality. Furthermore, out of the 19 (28.9%) cases in whom the first chelating therapy was started between 3 and 6 years of age, 16 (90%) patients remained alive and only 10% of these patients died. The mortality rate was high (50%) among those in whom the first chelating therapy was started at the age of 11 years and older. Thus, earlier initiation of chelating therapy led to better outcomes and reduced the mortality rate, with a highly statistically significant difference ( $P=0.0190$ ).

In the current study, pallor, jaundice, mongloid facies, hepatomegaly and splenomegaly were the main presenting clinical manifestations in almost patient group as there is 66.7% had pallor while 40% had mongoloid facies.

Splenomegaly appeared in 50% of patient group, while hepatomegaly appear in 36.7% of patient group. Also, history of splentectomy appear in 50% of Patient group.

This is in agreement with **Galanello**(13) who found that patients with thalassaemia major have a severe

It is not necessary because its value mentioned in table 6 :[34ks]Comment

found :[37ks]Comment

therapy :[38ks]Comment

In which table this result was represented? I prefer either discard this result or showing it in a new table :[39ks]Comment

Don't repeat previous information :[35ks]Comment

All these parameters are not illustrated in your results, please either left them to be used in other research or showing their values in a new table :[40ks]Comment

Why bold and italic font? Delete the word "with" :[36ks]Comment

microcytic hypochromic anemia with low mean corpuscular volume and mean corpuscular hemoglobin.

In the current study, it was found that mean serum iron and ferritin levels were significantly higher in Patient group than control group.

This in agreement with **Kurtoglu et al** (14) and **Taher et al** (15) who found significantly higher serum iron and serum ferritin and significantly lower TIBC in patients with thalassemia major compared with normal control individuals.

In the current study it was found that liver enzymes were significantly higher in Patient group with mean of (48.4±19.2 & 68.7±27.7) of AST and ALT respectively than the control group (24.7±3.12 & 24.8±3).

This is in agreement with **Sedigheh et al** (16) who noted the similar findings from study in Iran with significantly raised liver enzymes (ALT, AST) in homozygous thalassemia major patients than in controls.

Also in agreement with **Suman et al** (17) who noted positive correlation was between number of transfusions and serum ferritin level with correlation coefficient (r=+.33). As Iron deposition in liver takes place, its functions are affected which are predicted by raised ALT and AST. There was a positive correlation between serum ferritin and liver enzymes (r=0.87±84).

**Asharaf S et al** (18) who observed during a study that some disturbances occur in liver functions in hepatitis negative thalassemia patients with iron overload.

**Barton et al** (19) noted similar results as serum ferritin increases liver enzymes also increases.

In the current study the Patient group had a significantly higher insulin resistance index with mean of (2.73±.26) while was (1.76±.14) in control group while BFI was reduced in Patient group than control but the difference was not significant.

This is in agreement with **Suvarna, et al** (20) who noted higher fasting plasma insulin levels with increased Insulin Resistance Index and normal plasma glucose, suggesting the presence of insulin resistance

before the onset of frank impaired glucose tolerance test or diabetes. The high insulin level is probably in compensation for the insulin resistance in an attempt to maintain euglycemia. The increase in insulin levels has been postulated due to reduced hepatic insulin extraction rather than an increase in the secretory response.

This is in agreement with **Ghergherehchi and Habibzadeh**, (21) who documented that, β-TM patients had significantly higher IRI and elevated transaminases than the control group; but the decrease in BFI was not significant.

In the current study, it was found that Patient group showed abnormal OGTT results in 26.3% of which is significantly higher than control group (5%). Three was diabetic (10%) and 7 were IFG (23.6)

This is in agreement with **Ghergherehchi and Habibzadeh**, (21) who documented that abnormal OGTT results were observed in 33.3% of b-TM patients, which was significantly higher than in the control group (7.5%). Diabetes mellitus and IFG prevalence in b-TM patients were 7.7 and 25.6%, respectively. The prevalence of abnormal OGTT, DM and IFG in this study is similar to the literature (**Noetzi et al., 2012**) (22).

In another study from Iran, the prevalence of DM and IFG was 5.1 and 7.7%, respectively (**Shams et al., 2010**) (23). It is possible that these differences were mostly related to the geographical location of the studied population, duration of the diagnosis, number of patients and chelation therapy in each study

## References

1. **Varmani, S.G., Mehta, K., Garg, M. and Sadhu, S.D., (2012):** Diabetes mellitus in β-thalassaemia major-pathogenesis and management strategies. *International Journal of Food and Nutritional Sciences*. 3: 127-131.
2. **Pagon, R.A., Adam, M.P., Ardinger, H.H., Bird, T.D., Dolan, C.R., Fong, C.T., Smith, R.J., Stephens, K., Cao, A., Galanello, R. and Origa, R., (2013):** Beta-Thalassemia. *NCBI Bookshelf*.
3. **Bahar, A., Kashi, Z., Sohrab, M., Kosaryan, M. and Janbabai, G., (2012):** Relationship between beta-globin gene carrier state and insulin resistance. *Journal of Diabetes and Metabolic Disorders*, 11:22.

In Italic font wherever found :[42ks]Comment

If you want to remain the :[41ks]Comment results of liver function test, you must talk about them in your abstract, aim of study, as well as in methods and laboratory tests

4. **Garcia-Compean, D., Jaquez-Quintana, J.O., Gonzalez-Gonzalez, J.A. and Maldonado-Garza, H., (2009):** Liver cirrhosis and diabetes: risk factors, pathophysiology, clinical implications and management. *World J Gastroenterol*, 15(3): 280-288.
5. **Simcox JA and McClain DA. (2013):** Iron and diabetes risk. *Cell Metab*. 17(3):329–341.
6. **Rund D, Rachmilewitz EA. (2001).** Pathophysiology of  $\alpha$ - and  $\beta$ -thalassaemia: therapeutic implications. *Semin Hematol*;38 (9): p.343
7. **Tangvarasittichai S, Pimanprom A, Choowet A, Tangvarasittichai O. (2013)** Association of iron overload and oxidative stress with insulin resistance in transfusion-dependent  $\beta$ -thalassaemia major and  $\beta$ -thalassaemia/Hb E patients. *Clin Lab*.;59(7–8):861–868.
8. **Chen W , Zhang X. Shang X, et al.(2010):**The molecular basis of beta thalassaemia intermedia in Southern China: Genotypic heterogeneity and phenotypic diversity. *BM C Med Genet*:11:31.
9. **Islam S . (2017) :**Thalassemsias in South Asia: clinical lessons learnt from Bangladesh. *Orphanet J Rare Dis* 2017; 12:93
10. **Li CK, Luk SW, ling SC, et al.,(2002).** Morbidity and mortality patterns of thalassaemia major patient in pediatric department of three regional hospital: retrospective study. *Hong Kong Med J*; 8:255–260.
11. **Dey P, Konwar G, Sarkar B (2019).**Body mass index in thalassaemia children. *J. Evolution Med. Dent. Sci.* 2019;8(19):1537-1539, DOI: 10.14260/jemds/2019/3
12. **Al-Kherbash HA, Al-Awdi A and Hasan NS.(2017).** Pattern and clinical profile of thalassaemia among pediatric patients attending the Yemeni Society Centers for Thalassaemia and Genetic Blood Disorders in Yemen.*The Scientific Journal of Al-Azhar Medical Faculty, Girls, 1(2)*, p.43.
13. **Galanello, R., and Origa, R. (2010).**Beta-thalassaemia. *Orphanet journal of rare diseases*, 5(1), 11.
14. **Kurtoglu AU, Kurtoglu E and Temizkan AK. (2012):** Effect of iron overload on endocrinopathies in patients with  $\beta$ -thalassaemia major and intermedia. *Endokrynol Pol*. 63(4):260–263.
15. **TaherA ,Musallam KM, Kaarimi M, et al.(2010):**Overview on practices in thalassaemia intermedia management aiming for lowering complication rates across region of endemicity :the optimal care study. *Blood* . 115(10):1886-1892.
16. **Sedigheh S, Mohammed TA, Maryam MM, Leili K, Heshmat I, Fahimah J et al.** Evaluation of serum insulin, glucose, lipid profile and liver functions in beta-thalassaemia major patients and their correlation with iron overload. *Labmedicine*. 2010;41:486-9.
17. **Suman RL et al. Int J Res Med Sci. 2016** Aug;4(8):3271-3274
18. **Asharaf S, Mohammad Y, Fauzia al Y, Lolwa al N, Noora al M, Aml S, Vincenzo de S.** Longitudinal study on liver functions in patients with thalassaemia major before and after desferroxamine therapy. *Mediterranean journal of hematology and infectious diseases*. 2014;6
19. **Barton JC.** Chelation therapy for iron overload. *Curr Gastroenterology*. 2007;23:74-82
20. **Suvarna J, Ingle H, Deshmukh CT. 2006.** Insulin resistance and  $\beta$  cell function in chronically transfused patients of thalassaemia major. *Indian Paediatr.*;43(5):393–400
21. **Ghergherehchi R, and Habibzadeh A (2015):** “Insulinresistance and  $\beta$  cell function in patients with  $\beta$ -thalassaemia major”*Hemoglobin*. 2015; 39(1):69-73.doi: 10.3109/03630269.2014.999081. Epub 2015 Jan9.
22. **Noetzli LJ, Panigrahy A and Mittelman SD.(2012):**Pituitary iron and volume predict hypogonadism in transfusional iron overload. *AM J Hematol*;87:167.
23. **Shams S, HaghiAshtiani MT, Monajemzadeh M, et al. 2010 .**Evaluation of serum insulin, glucose, lipid profile, and liver function in  $\beta$ -thalassaemia major patients and their correlation with iron overload. *Lab Med.*;41(8):486–489.