

PHYTOCHEMICAL ANALYSIS AND *IN-VITRO* ANTIOXIDANT ACTIVITIES OF SOME SELECTED HIGHER FUNGI OF THE POLYPORE ORDER FROM OYO STATE, SOUTH WEST OF NIGERIA

Abstract.

Higher-fungi (Hf) of the polypore mushrooms are considered to have unique secondary metabolites, making them reservoirs of therapeutically significant bioactive compounds. Phytochemical and antioxidant properties of the Hf were accessed in this study.

Four Hf, were collected in the wild from some parts of Oyo state, Nigeria and were identified to species at the University of Ibadan Botany Department Laboratory. Methanol extracts of air-dried and powdered Hf were analysed for the presence of phytochemicals using standard procedures and *in-vitro* antioxidant activities using, 1,1-Diphenyl-2-picrylhydrazyl (DPPH), Ferric Reducing Antioxidant Power (FRAP) and Hydrogen peroxide (H₂O₂) assays. Results were presented as Mean ± SEM , graphs plotted using Microsoft Excel, analysed using one-way ANOVA and p≤0.05 was significant..

These Hf were identified as *Lycoperdon rimlatum* (*Lr*) FFUI1, *Trametes versicolor* (*Tv*) FFUI2, *Ganoderma lucidum* (*Gl*) FFUI3, and *Daedelia quarcina* (*Dq*) FFUI4 and were recorded on the NCBI with accession numbers EU833664.1, JQ621899, JQ520179.1 and KP171209.1, respectively. All the Hf showed positive tests for the presence of saponin, tannin, alkaloid, terpenoid, carbohydrate, tannin and flavonoids. The Hf showed antioxidant activities, the highest DPPH inhibition was by *Tv* (94.48%), FRAP was by *Gl* (0.16 mg/g) and H₂O₂ inhibition was by *Lr* (70.90%).

The antioxidant activities observed were due to the presence of useful phytochemicals making them therapeutically significant.

Key words: Higher- fungi, Phytochemicals, and *In-vitro* Antioxidant activities.

1.0 INTRODUCTION.

Fungi constitute an undescribed, undocumented lineage of eukaryotes with huge ecological and economic effects (Carris, *et al.*, 2012). Higher fungi have been considered as one of the highly diversified biological resources in the world (Zhong, & Xiao, 2009). They are diverse, heterotrophic organisms with unique nutritional and ecological requirements (Okaroh, 2017). These Mushrooms have been reported as foods with the therapeutic value used in the management of hypercholesterolemia, hypertension, and cancer (Kumar, 2015). Natural products produced by microorganisms, plants, and animals that play important roles in defense are secondary metabolites (Tiwari & Rana, 2015).

This metabolite from fungi has been receiving great consideration globally since the 1940s when antibiotics were discovered (Manganyi & Ateba, 2020). Oxidation is essential for the energy production in several living organisms for the fueling of physiological processes (Augusto & Miyamoto, 2011). Although oxygen is essential for life, it can also worsen the harm done by oxidative events within the cell. Oxygen's oxidative property is vital in a range of biological activities. The production of ATP (adenosine triphosphate) by the mitochondria, which is used by the cell to produce energy, results in the synthesis of free radicals. Reactive oxygen species (ROS) and reactive nitrogen species (RNS), which are byproducts of the biological redox process, are both deadly and beneficial molecules in humans. Free radicals can be produced in humans all around the world by environmental factors like noise, radioactivity, smoke, and weedicides. Most endogenous and physiological reactive oxygen species (ROS) are by-products of the oxidative reaction process that occurs in the mitochondrial respiratory chain (Nathan & Cunningham-Bussel, 2013). Reactive oxygen species (ROS) have a variety of consequences on cell physiology, including aiding in the death of invasive bacteria, wound healing, and regenerative processes (Bhattacharyya *et al.*, 2014). Consuming antioxidants on a regular basis can help to reduce free radicals' potential for harm. Unquestionably, the delicate balancing act between these two opposing affects is crucial in life. At low to moderate concentrations, reactive species have positive impacts on immune response and cellular redox signaling; yet, at high concentrations, they can impair cell structures and function by inducing oxidative stress. The hydrogen-based scavenging, regulation of free radicals, and radical peroxide oxidation processes are all part of the mushroom's antioxidant action (Ferreira *et al.*, 2009). By strengthening the immune system, these antioxidants lower the risk of infection, cancer, and cardiac issues. External antioxidants from food supplements are required when endogenous antioxidants are insufficient to sufficiently protect the organisms from free radicals (Sen & Chakraborty, 2011).

God designed plants and other herbal products with specific therapeutic or curative properties, which are known as "phytochemicals," to benefit people (Venkataramaiah, 2020). Phytochemicals, as their name implies, are chemicals produced by plants from their primary or secondary metabolites (Anulika *et al.*, 2019). These phytochemicals come in various kinds depending on their medicinal capabilities. Many plants and herbal items include a number of chemical components known as secondary metabolites that combine to produce therapeutic effects. Given that they have fewer or no side effects as compared to conventional synthetic medications and that many natural therapies have their own curative powers, the therapeutic properties of many plants and herbal items are becoming more and more well-known and preferred (Ogunmefun, 2018). The

phytochemicals that medicinal plants and herbs create for defense are quite abundant, and it is these components that give them their therapeutic properties (Alamgir, 2017). Medicinal plants and herbs may contain a range of phytochemicals, including saponin, which can be used to lower blood cholesterol, nitrogen-rich alkaloids, which can be used as stimulants, tannins, which act as natural antibiotics, anthraquinones, which act as laxatives and dyes, cardiac glycosides for cardiac drugs, flavonoids, and antioxidant phenols, as well as other compounds (Oyugi, 2016). The objective of this research is to assess the phytochemical and antioxidant properties of the selected higher fungi.

2.0. MATERIALS AND METHODS.

2.1 Collection and identification of mushroom samples

In several areas of Oyo State (8°00' N4°00' E), the Southwest of Nigeria, during the rainy season (August to September 2016 and 2017), the fresh fruiting sections of the following macrofungi were taken from the wild: Saki (8.6726° N, 3.3943° E), Ibadan, (7.3775° N, 3.9470° E), Ogbomosho (8.1227° N, 4.2436° E) and Iseyin (7.9765° N, 3.5914° E). They were validated at the University of Ibadan's Botany Department Laboratory (7° 26' 00" N, 3° 54' 00" E). They were identified using descriptions from Alexopoulos *et al.* (1998). The macro-fungal DNA was extracted using hexadecyl trimethyl ammonium bromide (cTAB) (Möller *et al.*, 1992). Using the primer pairs pITS4-F and pITS1-R (5'-TCCGTAGGTGAACCTGCCG-3'), the internal transcribed spacer (ITS) region was amplified. At the International Institute of Tropical Agriculture in Ibadan, Oyo state, Nigeria, the electrophoresis on a % agarose gel was used to analyze the PCR data before sequencing. The produced sequences were examined to identify the closest sequence matches for taxonomic classification using the NCBI's basic alignment search tools (BLAST) (Thomson *et al.*, 1997). Their basidiocarps were gathered and kept in the macrofungi collection owned by Jonathan Gbolagade at the University of Ibadan's Botany Department.

2.2. Extraction of phytochemicals from selected Higher Fungi

To obtain the fraction of the methanol extract, 1.5 kg of the air-dried powdered sample was placed separately into the soxhlet chamber extractor and extracted with 7.5 liters of 95 % methanol for 24 hours at 40°C. As reported by Evans, (2009), the filtrate was evaporated using a rotating electrical evaporator in a vacuum. Crude methanol extract yield was measured and correctly labeled and maintained in hygienic glass containers at room temperature until it was ready for use. The crude methanol yield was calculated after the above extraction, and determined by deducting the final weight of the extract obtained from the initial weight multiply by 100.

% yield = $\frac{W1 - W2}{W1} \times 100$, Where W1 initial weight, W2 =final weight.

2.3. Qualitative Phytochemical analysis

Methanol extracts from medicinal mushrooms were being subjected to numerous chemical tests to classify particular bioactive constituents using standard procedures (Evans, 2009).

2.4. Quantitative determination of phytochemical constituents (Evans, 2009).

Under lower pressure, filtered raw mushroom extracts (200 ml) are concentrated and segmented with 70 percent (V/V) sequential extractions of n-hexane, chloroform, ethyl acetate, and ethanol. These four fractions were tested on secondary metabolites using qualitative phytochemical reactions. Triterpene/steroids, alkaloids, flavonoids, saponins, carbohydrates, tannins, and terpenoids are measured. As an empirical response to those measures, colour intensity was used.

2.5 In-vitro antioxidant assays

2.5.1 DPPH radical scavenging activity method.

Following the protocol established by Shimada, *et al.*, (1992), the scavenging ability of mushrooms was assessed with some modifications. First, 0.5 ml of the aliquot sample extract was put in the test tubes with radical 2.9 ml of 200 μ M DPPH at various ethanol concentrations. The mixture was shaken vigorously and allowed to stand for 30 minutes at room temperature in the dark. Using a UV spectrophotometer, the reaction blend was measured at 515 nm. The solvent extraction was used as a blank, without an extract. The base norm used as ascorbic acid. The scavenging effect was determined based on the following formulation:

$$\text{Scavenging effect (\%)} = 1 - [(\text{Absorbance sample} / \text{Absorbance control}) \times 100]$$

2.5.2 Ferric reducing antioxidant power (FRAP) assay.

(Benzie & Strain, 1996 and Huang, *et al.*, 2005) reported process was used. 1.8 ml of the newly developed FRAP reagent was controlled and incubated for 10 minutes at 30 °C in a water bath. Absorbance was then recorded at 0 min (t^0). 100-500 μ l of mushroom sample extract and 100 μ l of distilled water were applied directly to the test tube at approximately 30 °C for 30 minutes. The absorption was subsequently taken at about 700 nm (t_{30}). Ferrous sulphate was the standard used. The antioxidant potential of the sample extract was measured against a standard ferrous sulfate curve, and the FRAP value was calculated as equal to μ M Fe²⁺ + per gram extract and calculated using the equation:

$$\text{FRAP value} = \text{Absorbance (sample + FRAP reagent)} - \text{Absorbance (FRAP reagent)}$$

2.5.3 H₂O₂ radical scavenging assay.

The extract's ability to scavenge hydrogen peroxide was tested using the Ruch, *et al.*, (1989) method. A solution of hydrogen peroxide (2mol/l) was prepared in the phosphate buffer (pH 7.4). Extracts (1–10 μ g / ml) were applied to the hydrogen peroxide solution (0.6 ml). Using a blank

solution containing a phosphate buffer without hydrogen peroxide and comparing ascorbic acid with the reference compound after 10 min, hydrogen peroxide absorbance at 230 nm was calculated.

$$\text{H}_2\text{O}_2 \text{ activity (\%)} = (\text{Abs control} - \text{Abs sample}) / \text{Abs (control)} \times 100$$

2.6 Statistical Analysis

Results were presented as Mean \pm SEM, graphs plotted using Microsoft Excel, analysed using one-way ANOVA and $p \leq 0.05$ was significant.

3.0 Results and Discussions

These macrofungi were identified as *Lycoperdon rimlatum* FFUI1, *Trametes versicolor* FFUI2, *Ganoderma lucidum* FFUI3, and *Daedelia quarcina* FFUI4 and were recorded on the NCBI with accession numbers EU833664.1, JQ621899, JQ520179.1 and KP171209.1, respectively (Table 1).

In order to properly assign medicinal qualities in research, precise species identification is crucial. The internal transcribe spacer (ITS) region of fungal genes, which have been sequenced, has aided in identifying and classifying related species (White *et al.*, 1990). The Higher fungi used in this research were identified using the internal transcribe spacer (ITS) region. Based on an existing database of ITS sequences, all of the taxa were correctly recognized.

Table 1

The percentage yield of metabolites from the Higher Fungi is represented in Table 2. The highest percentage yield was observed with the organism *Lycoperdon rimulatum* at 46.98%, while the lowest was observed with *Trametes versicolor* at 26.26%.

Table 2

The qualitative phytochemical analysis of the Higher fungi samples is represented in Table 3. In the analysis, it was observed that all the organisms showed positive tests for the presence of saponin, tannin, alkaloid, terpenoid, carbohydrate and tannin. The steroid tested positive with

only *Trametes versicolor* while anthocyanin and phlobatannin tested positive with only *Ganoderma lucidum*. Flavonoids tested positive with all the Higher fungi except *Daedalea quercina*.

The quantitative phytochemical analysis of the Higher fungi samples is represented in Table 4. In the quantitative analysis of carbohydrate content, the highest amount of carbohydrate was observed with *Lycoperdon rimulatum* at 2.38 ± 0.03 , while the lowest was observed with *Trametes versicolor* at 0.19 ± 0.01 . For the flavonoid content, *Ganoderma lucidum* had the highest amount with 0.2 ± 0.01 while the lowest amount was observed with *Trametes versicolor* with 0.11 ± 0.01 . In the quantitative analysis of Terpenoid content, *Lycoperdon rimulatum* has the highest amount of terpenoid with 0.60 ± 0.03 , while *Trametes versicolor* has the lowest with 0.20 ± 0.0 . For saponin content, *Ganoderma lucidum* has the highest amount with 0.18 ± 0.03 while the lowest amount was observed with *Lycoperdon rimulatum* with 0.02 ± 0.01 . For alkaloid, the highest amount was observed with *Trametes versicolor* while the lowest amount was observed with *Daedalea quercina* with 0.02 ± 0.01 . For the quantitative presence of tannin, the highest amount was observed with *Ganoderma lucidum* with 7.73 ± 0.0 while the lowest was observed with *Trametes versicolor* with 2.6 ± 0.02 .

The link between phytochemistry and pharmacology is critical to consider when designing studies on the medicinal potential of plants and herbal medicines. In general, the results showed that phytochemicals were present in all samples, but at varying quantities. Alkaloids, tannins, saponins, and phenols are considered anti-nutrients since they are poisonous when consumed in large doses. However, it has been shown that several of the phytochemicals found in mushrooms have therapeutic properties. The presence of bioactive phytochemical constituents found in all of the Higher fungi studied has been suggested as the reason for their traditional uses in the treatment of inflammation, pains, hemostatic, diuretic, nutrition, antibiotics, and antitumor agents, which is supported by the findings of Edeoga and Erita (2001), who discovered alkaloids' significant pharmacological modulation. The presence of alkaloids in the samples shows that they have analgesics and bactericides medicinal value, confirming Stary's (1998) findings. Furthermore, presence of phenols in the Higher fungi samples makes them antiseptics and antifungal (Gill, 1992). The presence of flavonoids in the Higher fungi suggests that they have antioxidant healing properties, which backs up Okwu, (2004) findings that flavonoids can prevent cancer and oxidative cell damage. External antioxidants from food supplements are required when endogenous antioxidants are insufficient to protect organisms from free radicals. This exogenous antioxidants can be obtained naturally from flavonoids, according to Litescu *et*

al. (2011). The presence of tannin in the sample, on the other hand, is linked to wound healing, supporting Okwu, (2004) results that the concentration of tannin present in mushrooms can draw tissues together to aid wound healing.

Table 3

Table 4

The percentage inhibition of radical of methanolic extract of the selected higher fungi and Vitamin C, using DPPH assay are represented in Figure 1. At 100 µg/ml, the highest percentage inhibition was observed with Vitamin C, a standard antioxidant by 94.63% followed by *Trametes versicolor* by 94.48 % while the least percentage inhibition was observed with *Lycoperdon rimulatum* by 89.44 %, meanwhile at 500 µg/ml, the standard sample, vitamin c had the highest percentage inhibition by 87.1% followed by *Trametes versicolor* by 85 % while the least percentage inhibition was observed with *Lycoperdon rimulatum* by 48.3 %. Overall, it was observed that percentage inhibition reduced with an increase in the concentration of the extracts. The DPPH test provides information on the reactivity of compounds with a stable free radical DPPH that gives a strong absorption band at 517nm in visible region. When the odd electron becomes paired off in the presence of a free radical scavenger, the absorption reduces and the DPPH solution is decolorized as the colour changes from deep violet to light yellow. The degree of reduction in absorbance is reflective of the antioxidant potency of the extract (Barreira *et al.*, 2008).

The ferric reducing power activity of methanolic extract of the selected higher fungi and Vitamin C assay are represented in Figure 2. At 100 µg/ml, the highest ferric reducing power activity was observed with *Ganoderma lucidum* by 0.14 while the least activity was observed with *Lycoperdon rimulatum* by 0.01. .At 500 µg/ml, the standard sample, vitamin c had the highest ferric reducing power activity by 0.37 followed by *Ganoderma lucidum*, by 0.36, while the least was observed with *Lycoperdon rimulatum* by 0.16. Overall, it was observed that ferric reducing power activity increased with an increase in the concentration of the extracts and the highest reducing power was observed with Vitamin C at 500 (µg/ml) with 0.37 while the lowest reducing power was observed with *Lycoperdon rimulatum* with 0.01.at

100 (µg/ml). The reducing power of a compound is related to its electron transfer ability, and may therefore serve as a significant indicator of its potential antioxidant activity (Abbasi *et al.*, 2013). The reductive ability of the extract occurred in a dose dependent manner. This can be attributed to the electron donating ability of the polyphenols present in the extract.

The percentage inhibition of methanolic extract of the selected higher fungi and Vitamin C using Hydrogen peroxide (H₂O₂) radical assay are represented in Figure 3. At 20 µg/ml, the highest percentage inhibition was observed with Vitamin C, a standard antioxidant by 83.5% followed by *Lycoperdon rimulatum*, *Trametes versicolor* and *Daedelia quarcina* by 70.9 %, 65.3 %, and 65.3 % respectively while the least percentage inhibition was observed with *Ganoderma lucidum* by 55.28 %. At 100 µg/ml, the standard sample, vitamin C had the highest percentage inhibition by 54.1 % followed by *Daedelia quarcina* by 35.2 %, while the least percentage inhibition was observed with *Trametes Versicolor* by 24.1 %. In the overall, it was observed that percentage inhibition reduced with an increase in the concentration of the extracts, and *Lycoperdon rimulatum* gave the highest inhibition at 20(µg/ml) with 70.9 % while the least was observed at 100(µg/ml) with *Trametes Versicolor* at 24.1%. Hydrogen peroxide is a precursor for the production of hydroxyl radicals in cellular components. Hydroxyl radicals are considered to be one of the rapid initiators of lipid peroxidation process, abstracting hydrogen atoms from polyunsaturated fatty acid, which brings about peroxidic reactions of membrane lipids (Lipinski, 2011).

The capacity of a chemical to transport electrons influences its reducing power. Reducing power is frequently used to assess the anti-inflammatory properties of polyphenols, such correlated with reductones' existence, which exerts antioxidant activity by severing the cycle of free radicals by giving atom of hydrogen (Duan and Jiang, 2007). The FRAP value acts as a Fe (II) TPTZ extract's reduction power as measured. A defense mechanism that balances unpaired radicals strengthens living things, and this defense mechanism is made up of oxidative enzymes. Accordingly, antioxidant food supplements can help the body's defense mechanism in neutralizing or mitigating oxidative harm, supporting the findings of the American Dietetic Association (2003) that a healthy diet can provide all of the antioxidants needed and Cadenas' (1997) early findings that dietary antioxidant intake can improve protection against free radicals. All the higher-fungi tested positive for antioxidant properties *in vitro* using DPPH, FRAP, and Hydrogen peroxide assays, which is consistent with Mau *et al.*,(2002) findings, that

natural bioactive products generated by microorganisms, macro-fungi, plants, and animals have antioxidant properties. The hydrogen peroxide scavenging activity of the Higher-fungi can be attributed to the proton donating potential of their phytochemical components. It also corroborated the findings of Chang *et al.* (2007) that hydrogen scavenging is a key component of the antioxidant action of mushrooms, keeping free radicals and radical peroxide oxidation in check.

Figure 1

Figure 2

Figure 3

4.0 CONCLUSION

Finding novel natural products from wild sources could benefit the food business by introducing better and safer antioxidants that offer strong defense against oxidative damage, which happens in both the body and our everyday foods. Therefore, new wild non poisonous mushrooms could be introduced as natural supplies for this purpose. The researched Higher-fungi seems to be viable sources of bioactive substances that might have intriguing antioxidant effects in animal systems. Due to the Fungi's high phenolic content, it exhibits substantial antioxidant action. As a result, *Ganoderma lucidum*, *Trametes versicolor*, *Daedelia quercina*, and *Lycoperdon rumilatum*. are recognized as superb source of bioactive substances that can be turned into medicines to treat oxidative stress.

The current research was able to identify indigenous mushrooms using molecular analysis, enriching and adding to our knowledge of mushroom biodiversity in Nigeria. The methanolic extracts of the studied Higher-fungi (*Trametes versicolor*, *Daedelia quercina*, *Ganoderma lucidum*, and *Lycoperdon rimulatum*) indicated the existence of flavonoids as well as antioxidant activity. Consuming these mushrooms can therefore serve as a source of exogenous antioxidants to supplement endogenous antioxidants in nutritionally supplemented diets, which can be extremely beneficial as protection against cancer, heart disease, boosting immunity, and anti-aging, supporting the findings of Omeonu *et al.*, (2022). Preclinical and clinical studies are also needed to assess the effectiveness of the natural extracts of these mushrooms in the treatment or prevention of a variety of human diseases.

REFERENCES

Abbasi, M. A., Saleem, H., Riaz, T., & Ajaib, M. (2013). Determination of Antioxidant Activity and Phytoconstituent Screening of *Euphorbia heterophylla* Linn.

Alamgir, A. N. M. (2017). Therapeutic use of medicinal plants and their extracts: volume 1. Springer International Publishing AG.

Albus, U. (2012). Guide for the Care and Use of Laboratory Animals (8th edn). Allen, A., & Flemström, G. 2005. Gastroduodenal mucus bicarbonate barrier: protection against acid and pepsin. *American Journal of Physiology-Cell Physiology*, 288.1: C1-C19

Alexopoulos, C. J., Mims, C. W., & Blackwell, M. (1996). Introductory mycology (No. Ed. 4). John Wiley and Sons.

American Dietetic Association. (2003). Position of the American Dietetic Association and Dietitians of Canada: vegetarian diets. *Journal of the Academy of Nutrition and Dietetics*, 103(6), 748.

- Anulika, N. P., Ignatius, E. O., Raymond, E. S., Osasere, O. I., & Abiola, A. H. (2016). The chemistry of natural product: Plant secondary metabolites. *Int. J. Technol. Enhanc. Emerg. Eng. Res*, 4(8), 1-9.
- Augusto, O., & Miyamoto, S. (2011). Oxygen radicals and related species. *Principles of free radical biomedicine*, 1, 19-42.
- Barreira, J. C., Ferreira, I. C., Oliveira, M. B. P., & Pereira, J. A. (2008). Antioxidant activities of the extracts from chestnut flower, leaf, skins and fruit. *Food chemistry*, 107(3), 1106-1113.
- Benzie, I. F., & Strain, J. J. (1996). The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Analytical biochemistry*, 239.1: 70-76.
- Bhattacharyya, A., Chattopadhyay, R., Mitra, S., & Crowe, S. E. (2014). Oxidative stress: an essential factor in the pathogenesis of gastrointestinal mucosal diseases. *Physiological reviews*, 94(2), 329-354.
- Cadenas, R. F., Gil, J. A., & Martin, J. F. (1992). Expression of Streptomyces genes encoding extracellular enzymes in Brevibacterium lactofermentum: secretion proceeds by removal of the same leader peptide as in Streptomyces lividans. *Applied microbiology and biotechnology*, 38(3), 362-369.
- Carris, L. M. (2021). Carris, LM, CR Little, and CM Stiles. 2012. Introduction to Fungi. The Plant Health Instructor. DOI: 10.1094/PHI-I-2012-0426-01. *Phytopathology News*.
- Carris, L. M., & Stiles, C. M. (2012). Introduction to Fungi.
- Chang, H. Y., Peng, W. H., Sheu, M. J., Huang, G. J., Tseng, M. C., Lai, M. T., ... & Chang, Y. S. 2007. Hepatoprotective and antioxidant effects of ethanol extract from Phellinus merrillii on carbon tetrachloride-induced liver damage. *The American journal of Chinese medicine*, 35.05: 793-804.
- Chang, S. T., & Wasser, S. P. (2012). The role of culinary-medicinal mushrooms on human welfare with a pyramid model for human health. *International journal of medicinal mushrooms*, 14.2.
- Duan, X., Wu, G., & Jiang, Y. (2007). Evaluation of the antioxidant properties of litchi fruit phenolics in relation to pericarp browning prevention. *Molecules*, 12(4), 759-771.
- Edeoga, H. O. (2001). Erita: Alkaloids, Tannins and Saponins contents of some medicinal plants. *J Med Aro Sci*, 23, 344-349.
- Ellman, G. L. (1959). Tissue sulfhydryl groups. *Archives of biochemistry and biophysics*, 82.1: 70-77.
- Evans, W. C. (2009). *Trease and evans' pharmacognosy E-book*. Elsevier Health Sciences.

- Ferreira, I. C., Barros, L., & Abreu, R. (2009). Antioxidants in wild mushrooms. *Current Medicinal Chemistry*, 16(12), 1543-1560.
- Hauge, K., Bergene, E., Chen, D., Fredriksen, G. R., & Holmen, A. (2005). Oligomerization of isobutene over solid acid catalysts. *Catalysis today*, 100(3-4), 463-466.
- Ignarro, L. J., Buga, G. M., Wood, K. S., Byrns, R. E., & Chaudhuri, G. (1987). Endothelium-derived relaxing factor produced and released from artery and vein is nitric oxide. *Proceedings of the National Academy of Sciences*, 84.24: 9265-9269.
- Kumar, K. (2015). Role of edible mushrooms as functional foods—a review. *South Asian J Food Technol Environ*, 1(3-4), 211-218.
- Lipinski, B. (2011). Hydroxyl radical and its scavengers in health and disease. *Oxidative medicine and cellular longevity*, 2011.
- Litescu, S. C., Eremia, S. A., Diaconu, M., Tache, A., & Radu, G. L. (2011). Biosensors applications on assessment of reactive oxygen species and antioxidants. In *Environmental Biosensors*. Intech Open.
- Manganyi, M. C., & Ateba, C. N. (2020). Untapped potentials of endophytic fungi: A review of novel bioactive compounds with biological applications. *Microorganisms*, 8(12), 1934.
- Mau, J. L., Lin, H. C., & Song, S. F. 2002. Antioxidant properties of several specialty mushrooms. *Food Research International*, 35.6: 519-526.
- Möller, E. M., Bahnweg, G., Sandermann, H., & Geiger, H. H. (1992). A simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. *Nucleic acids research*, 20(22), 6115.
- Nathan, C., & Cunningham-Bussel, A. (2013). Beyond oxidative stress: an immunologist's guide to reactive oxygen species. *Nature Reviews Immunology*, 13(5), 349-361.
- NRLA, N. 1995. Nutrient Requirements of Laboratory Animals. *Fourth Revised Edition*.
- Ogihara, Y., and Okabe, S. 1993. Effect and mechanism of sucralfate on healing of acetic acid-induced gastric ulcers in rats. *Journal of physiology and pharmacology: an official journal of the Polish Physiological Society*, 44.2: 109-118
- Ogunmefun, O. T. (2018). Phytochemicals—God's Endowment of Curative Power in Plants. *Phytochemicals: Source of Antioxidants and Role in Disease Prevention*, 7.
- Okaroh, B. C. (2017). *Physico-Chemical and Toxicological Profiles of Five Species of Mushroom In Anambra State, Nigeria and Their Potential for Bioremediation of Trace Metal Polluted Soil* (Doctoral dissertation).
- Okwu, D. E. (2004). Phytochemical and vitamin content of indigenous spices of South Eastern Nigeria. *J. Sustain. Agric. Environ*, 6, 30-34.

Omeonu, F. C., Jonathan, S. G., Salami, A. T., Azuh, V. O., & Ado, B. V. Anti-Ulcer and Blood-Boosting Effect of Diet Supplemented with *Daedalea quercina* from Ogbomoso, Oyo State, South West of Nigeria on Indomethacin Induced Gastric Ulcer In Rats.

Oyedemi, S. O., Bradley, G., & Afolayan, A. J. (2010). In-vitro and-vivo antioxidant activities of aqueous extract of *Strychnos henningsii* Gilg. *African Journal of pharmacy and pharmacology*, 4.2: 070-078.

Oyugi, J. O. (2016). *Analysis of Proximate, Micronutrients and Determination of Phytochemicals in Selected Medicinal Plants in Mbita-Homabay County* (Doctoral dissertation, Master's Thesis, Kenyatta University, Nairobi, Kenya).

Rubbo, H., Radi, R., Trujillo, M., Telleri, R., Kalyanaraman, B., Barnes, S., ... & Freeman, B. A. (1994). Nitric oxide regulation of superoxide and peroxynitrite-dependent lipid peroxidation. Formation of novel nitrogen-containing oxidized lipid derivatives. *Journal of Biological Chemistry*, 269(42), 26066-26075.

Ruch, R. J., Crist, K. A., & Klaunig, J. E. 1989. Effects of culture duration on hydrogen peroxide-induced hepatocyte toxicity. *Toxicology and applied pharmacology*, 100.3: 451-464.

Sen, S., & Chakraborty, R. (2011). The role of antioxidants in human health. In *Oxidative stress: diagnostics, prevention, and therapy* (pp. 1-37). American Chemical Society.

Shimada, T., Yun, C. H., Yamazaki, H., Gautier, J. C., Beaune, P. H., & Guengerich, F. P. (1992). Characterization of human lung microsomal cytochrome P-450 1A1 and its role in the oxidation of chemical carcinogens. *Molecular pharmacology*, 41(5), 856-864.

Thomson, E. (1997). Couple childbearing desires, intentions, and births. *Demography*, 34(3), 343-354.

Tiwari, R., & Rana, C. S. (2015). Plant secondary metabolites: a review. *International Journal of Engineering Research and General Science*, 3(5), 661-670.

Varshney, R., & Kale, R. K. 1990. Effects of calmodulin antagonists on radiation-induced lipid peroxidation in microsomes. *International journal of radiation biology*, 58.5; 733-743.

Venkataramaiah, C. (2020). Chemical Constituents of Plants as Promising Drug Candidates: Nature's (God) Benefaction to Mankind. *Chief Editor*, 53.

White, T. J., Bruns, T., Lee, S. J. W. T., & Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods and applications*, 18(1), 315-322.

Wolf, E. (1959). Electromagnetic diffraction in optical systems-I. An integral representation of the image field. Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences, 253.1274: 349-357.9

Zhong, J. J., & Xiao, J. H. (2009). Secondary metabolites from higher fungi: discovery, bioactivity, and bioproduction. *Biotechnology in China I*, 79-150.




Fig. 1. Inhibition of radical by methanolic extract of the selected higher fungi and Vitamin C, using DPPH assay.

T.v - *Trametes versicolor* *D.q*- *Daedalea quercina* *G.l* - *Ganoderma lucidum* *L.r*- *Lycoperdon rimulatum*




Fig. 2. Ferric Reducing Power activity of the selected higher fungi and Vitamin C.

T.v - *Trametes versicolor*, *D.q* - *Daedelia quercina*, *G.l* - *Ganoderma lucidum* *L.r* - *Lycoperdon rimulatum*.




Fig.3. Percentage inhibition of methanolic extract of the selected Higher fungi and Vitamin C using Hydrogen peroxide (H₂O₂) radical assay.

T.v - *Trametes versicolor*, *D.q* - *Daedelia quercina*, *G.l* - *Ganoderma lucidum* *L.r* - *Lycoperdon rimulatum*

List of tables

Strain	ID(NCBI) SUBMISSION	Accession number	Identified name	Blast search similarity
STB112	F1	EU833664.1	<i>Lycoperdon rimlatum</i>	99%
EMB5	F2	JQ621899	<i>Trametes versicolor</i>	99%
IUM4100	F3	JQ520179.1	<i>Ganoderma lucidum</i>	99%
Dai12697	F4	KP171209.1	<i>Daedelia quarcina</i>	99%

Table 1. Identity of the fungi strains

Table 2. Percentage yield of metabolites

ORGANISMS	% YIELD
<i>Trametes versicolor</i>	26.26
<i>Daedelia quercina</i>	38.68
<i>Ganoderma lucidum</i>	36.19
<i>Lycoperdon rimulatum</i>	46.98

TEST	<i>T.v</i>	<i>D.q</i>	<i>G.l</i>	<i>L.r</i>
SAPONIN	+	+	+	+
TANNIN	++	+++	+++	++
STEROID	+	-ve	--ve	-ve
ALKALOID	+	+	+	+
FLAVONOID	+	-ve	+	+
COUMARIN	-ve	-ve	-ve	-ve
EMODIN	-ve	-ve	-ve	-ve
TERPENOID	+	+	+	+

CARBOHYDRATES	+	+	+	+
ANTHRAQUINONE	-ve	-ve	-ve	-ve
ANTHROCYANIN	-ve	-ve	+	-ve
PHLOBATANNIN	-ve	-ve	+	-ve
CARDIAC GLYCOSIDE	-ve	-ve	-ve	-ve

Table 3. Qualitative phytochemical analysis of methanolic extracts of the selected Higher Fungi samples.

KEY; + - POSITIVE

-ve - NEGATIVE

T.v - *Trametes versicolor*, *D.q* - *Daedalea quercina*, *G.l* - *Ganoderma lucidum*, *L.r* - *Lycoperdon rimulatum*.

ORGANISM/TEST	CARBOHYDRATE	FLAVONOID	TERPENOID	SAPONINE	ALKALOID	TANNIN
<i>T.v</i>	0.19±0.01 ^a	0.11±0.01 ^a	0.20±0.0a	0.17±0.02 _a	0.13±0.03 _a	2.6±0.02 ^a
<i>D.q</i>	0.65±0.03 ^b	-	0.30±0.02 ^b	0.15±0.03 _a	0.02±0.01 _b	7.71±0.0 ^b
<i>G.l</i>						
<i>L.r</i>						

Table 4. Quantitative phytochemical analysis of the higher fungi samples

Values are expressed as Mean± SEM. (n = 3). Mean with the same letter in each column are not significantly different at 0.05 probability level.

Key; *T.v* - *Trametes versicolor*, *D.q* - *Daedalea quercina*, *G.l* - *Ganoderma lucidum*, *L.r*-*Lycoperdon rimulatum*

