

Commonness and Risk Factors Associated with Gastrointestinal Parasitic Infections among Inmates of Port Harcourt Children's Home.

ABSTRACT

Key words: Gastrointestinal Parasitic Infection, Risk factors, Children Home, Port Harcourt.

1. INTRODUCTION

Gastrointestinal infections are endemic worldwide and have been described as constituting the greatest single worldwide cause of illness and disease [1]. It has been estimated to affect about 3.5 billion people globally and caused morbidity in approximately 450 million people [2]. Infections are mainly transmitted via ingestion of water, soil, or food contaminated by feces containing the cysts of protozoans or eggs/larvae of helminths [1]. Most common among the intestinal helminths are *Ascaris lumbricoides*, hookworms, *Trichuris trichiura*, and *Strongyloides stercoralis*. About 1.2 billion people are estimated to be infected with *A. lumbricoides* globally while *T. trichiura* and hookworm infect up to 795 and 740 million people respectively [3]. They are most prevalent in regions exhibiting warm and moist climates coupled with poor sanitation and hygiene. Epidemiologically, it is well established that although individuals of all ages harbor worms, the highest rates occur among children in rural areas of the tropical and subtropical areas [4]. The climatic conditions in this part of the world favor the development and survival of these parasites. The high prevalence in a region results in infection and diseases that are the immediate causes of malnutrition and death in young children [5]. Several pathogenic protozoan parasites are responsible for the above health issues including *Entamoeba histolytica/dispar*, *Giardia lamblia* (also known as *Giardia intestinalis* and *Giardia duodenalis*), *Cryptosporidium*, and *Balantidium coli*, which are the most common species associated with significant

illnesses [6,7,8]. *Entamoeba histolytica* infection is considered one of the third most common causes of death after malaria and schistosomiasis [9].

Orphanages are a classical example of such a scenario. Orphans always lack ~~in~~ basic monetary and sanitary assets in their livelihood. Hence their debilitated lifestyle makes them susceptible to infections. Overcrowded places with ~~a~~ lack of cleanliness like orphanages are most dangerous as the infection spreads through them swiftly leading to multiple infections [10]. Orphans are the most vulnerable part of this group and they need the maximum support from the government. Records show that a global estimate of 162 million under-five years old children are documented to be stunted, 99 million underweight and 51 million wasted [11].

The overall prevalence rate of gastrointestinal parasitic infections in Port Harcourt ranges from 24.8-46.1% [12,13,14], due to varying geographical conditions. There is a paucity of information on the prevalence of gastrointestinal parasitic infections among inmates of orphanages of Port Harcourt. In the light of these developments, this present study was aimed at investigating the commonness ~~gastrointestinal~~ ~~gastrointestinal~~ parasites in relation to age, gender, educational background, and risk factors associated with inmates ~~in~~ Port Harcourt.

2. MATERIALS AND METHODS

2.1 Study Area

The study was conducted at Port Harcourt ~~Children~~ ~~Children's~~ Home, Borokiri, Port Harcourt town, Rivers State, Nigeria ~~in from~~ August to October 2019. The Borokiri town is a ~~neighbourhood~~ ~~neighborhood~~ of the city Port Harcourt situated just south of old GRA in Rivers State, Nigeria. ~~Co-ordinates~~ ~~Coordinates~~ for the town are latitude 4.749° N and longitude 7.035° E.

2.2 Study Design

This investigation involved a prospective study conducted on inmates of Port Harcourt ~~children~~ ~~children's~~ Home. Inmates were categorized into sex and age groups. ~~Questionnaire~~ ~~Questionnaires~~ were given to the inmates for information of their sociodemographic factors and personal hygiene.

2.3 Sample Size

An approximate minimum sample size of 63 was therefore estimated using the formula for a cross-sectional study as recommended by Godden [15]; $n = Z^2 \cdot p(1-p) / M^2$, where $n = n =$ sample size for infinite population, $Z = Z$ score (1.96), $p =$ Population proportion (assumed to be 50 % = 0.5) and $M =$ Margin of error (0.05).

2.4 Study Population

The study was carried out on 63 inmates (47 females and 16 males). All the inmates were informed clearly about the objective and procedure of the study and requested to sign a written consent. Research permit and ethical clearance ~~was were~~ sought from the Rivers State Hospitals Management Board and the Rivers State Ministry of Social Welfare and Rehabilitation.

2.5 Isolation and Identification of Gastrointestinal Parasites

2.5.1 Direct Smear Examination for Stool Samples

On a glass microscope slide, about 1-2 mg of stool was emulsified in a drop of normal saline (0.85% NaCl) on the left-hand side of the slide, and in Lugol's iodine on the right side of the slide. A ~~cover-~~

slip cover slip was then placed on each side, and the slides were scanned under 10x and 40x objective lenses of a light microscope, as required. Saline direct smear is used mainly for the detection of motility of intestinal protozoan trophozoites, which are seen in liquid or semi-liquid specimens. Iodine direct smear shows the characteristic features of the diagnostic stages in more details detail [16].

2.5.2 Formol-Ether Sedimentation Concentration Technique Ritchie

Although, this formol ether technique cannot detect trophozoites, it is considered as the best concentration technique used in diagnostic parasitology laboratories for the detection of cysts, ova, and larvae [17,18]. The Ritchie sedimentation technique was performed by emulsifying about 2 g of stool in 10-15 ml of 10% formol saline. The suspension was allowed to stand for 30 minutes, and then strained through two layers of gauze into a 15 ml conical centrifuge tube and centrifuged at 2000 rpm for 5 minutes. When needed, the washing step was repeated until the supernatant becomes clear. The sediment was resuspended with 10 ml of 10% formal saline and allowed to stand for 5-10 minutes. A total of 3 ml of diethyl ether was added, and then the tube was shaken vigorously for 30 seconds and centrifuged at 2000 rpm for 5 minutes. After centrifugation, the fecal debris layer was loosened by a wooden stick, and the tube rapidly inverted to discard the top three layers while the sediment remained at the bottom. One to two drops of iodine were added to the sediment and mixed well. Then, part of the sediment was transferred to a microscope slide, covered with a cover glass, and scanned microscopically under low and high objective lenses [19].

2.6 Ethical Considerations

Institutional ethical clearance and the research permit and authorization letter were obtained from Rivers State Hospitals Management Board (RSHMB). Before sampling, concerned authorities such as the Rivers State Ministry of Social Welfare and Rehabilitation were contacted and a request for permission was made after explaining the objective of the study. Prior to sample collection inmates were informed clearly about the objective and procedure of the study. Participation was very voluntarily voluntary, without any slightest negative consequence, and samples were collected when they fully agreed by signing an informed consent. Potential inmates were told that there were no foreseeable risk or undesirable side effect during fecal sample collection and any information obtained were to remain confidential.

2.7 Data Analysis

The data collected from the study area were entered in Microsoft office excel 2016 before being imported to SPSS version 23 and was were employed for data entry and statistical analysis. Descriptive statistics were mainly used to describe the characteristics of the study population including the prevalence of the GIPIs. Qualitative data were determined and presented as frequencies and percentages. Statistical significance of differences in proportions was evaluated by the Chi-Square test with a significant value of $p < 0.05$ used for all tests. Chi square Chi-square analysis was run to determine the association of variables with GIPIs GIPI infections.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Prevalence of gastrointestinal parasites among inmates of children children's home.

Stool specimens were collected from the inmates over a period of 8 weeks. A total of 63 inmates were examined in this study with 47 females and 16 males of which 19(30.2%) were found to be infected with one or more gastrointestinal parasites species. Various gastrointestinal parasites parasite species isolated were *Ascaris lumbricoides* 10(15.9%), *Trichuris trichiura* 7(11.1%), and hookworms 2(3.2%) respectively.

3.1.2 Sex-Distribution of gastrointestinal parasites among inmates

The distribution of gastrointestinal parasites according to genders were established, of which females had significantly more gastrointestinal parasitic infections 13(68.4%) compared to males 6(31.6%) ($p<0.05$) (Table 2). The distribution of various parasite species patterns was not affected by gender.

Legend: ♀ : Female; ♂ : Male; $p<0.05$

Fig 1: Sex-Related Distribution of Gastrointestinal Parasites

3.1.3 Distribution of gastrointestinal parasitic infections among inmates according to age groups

In this study, a relatively high number of 17(44.8%) of the inmates of age group 11-15 years were infested with gastrointestinal parasites while the age group of 1-5 years had 2(22.2%) infected with gastrointestinal parasites ($p<0.05$). In addition to all the age groups, the predominant gastrointestinal parasites identified were *A. lumbricoides* followed by *T. trichiura* and hookworm (Table 1).

Table 1: Age-related distribution of gastrointestinal parasites

Distribution of Gastrointestinal parasites								Chi square	p-value
Age	AL N(%)	HK N(%)	TT N(%)	Negative cases N(%)	Positive cases N(%)	Total examined N(%)	Chi-square (χ^2)		
1 - 5yrs	2(22.2)	0(0.0)	0(0.0)	7(77.8)	2(22.2)	9(100.0)	10.799	0.005	

Comment [FIMH1]: P-value should be revised

6 - 10yrs	0(0.0)	0(0.0)	0(0.0)	3(100.0)	0(0.0)	3(100.0)
11 - 15yrs	8(21.1)	2(5.3)	7(18.4)	21(55.2)	17(44.8)	38(100.0)
16 - 20yrs	0(0.0)	0(0.0)	0(0.0)	10(100.0)	0(0.0)	10(100.0)
36 - 40yrs	0(0.0)	0(0.0)	0(0.0)	3(100.0)	0(0.0)	3(100.0)
Total	10(15.9)	2(3.2)	7(11.1)	44(69.8)	19(30.2)	63(100.0)

Legends: A.L: - *Ascaris lumbricoides*; HK: - Hookworm; TT: - *Trichuris trichiura*; $p < 0.05$

Among the participant's educational levels, the primary school children had the most prevalent cases of gastrointestinal parasites (52.6%) followed by secondary school children and the pre-school age with statistical significance ($p < 0.05$) (Table 2).

Table 2: Sociodemographic factors Educational level and distribution of gastrointestinal parasites

Socio-demographic Factors	Distribution of Gastrointestinal parasites				Positive cases (%)	Total examined	Chi square (X ²)	p-value
	AL N(%)	HK N(%)	TT N(%)	Negative cases N(%)				
Educational level								
Educated	0(0.0)	0(0.0)	0(0.0)	3(100)	0(0.0)	3(100)	12.438	0.008
Pre-School age	2(22.2)	0(0.0)	0(0.0)	7(77.8)	2(22.2)	9(100)		
Primary	6(31.6)	0(0.0)	4(21.1)	9(47.4)	10(52.6)	19(100)		
Secondary	2(6.9)	6.9	10.3	22(75.9)	7(24.1)	29(100)		
Tertiary	0(0.0)	0(0.0)	0(0.0)	3(100)	0(0.0)	3(100)		
Total	10(15.9)	3.2	11.1	44(69.8)	19(30.2)	63(100)		

Legends: AL: - *Ascaris lumbricoides*; HK: - Hookworm; TT: - *Trichuris trichiura*; $p < 0.05$

Comment [FIMH3]: P-value should be revised

Comment [FIMH2]: Did the authors mean "Non-educated"

Comment [FIMH4]: 2(6.9)

Comment [FIMH5]: 3(10.3)

Comment [FIMH6]: 2(3.2)

Comment [FIMH7]: 7(11.1)

3.2 Gastrointestinal parasites and Possible Risk Factors

The risk factors associated with gastrointestinal parasites were examined. These factors included the use of drug drugs in the last three months, hand wash washing after toilets, footwear, and washing hands after contact with soil.

3.2.1 The use of the drug in the last three months and its association with gastrointestinal parasites

We investigated the effects of some hygienic practices like the use of drug drugs in the last three months. Inmates who took their drugs in the last three months were shown to be more parasitized 14(36.8%) compared to those who did not though were also parasitized. These were found to be significant ($p < 0.05$) (Table 3).

3.2.2 Hand wash after visiting toilets and its association with gastrointestinal parasites

The effects of hand washing was were considered, whether they were washing their hands after visiting the toilet. The inmates who claimed to wash their hands after toileting were found more infected had 17(34.0%) compared to inmates who were infected and did not regularly wash their hands. These variations were statistically significant ($p < 0.05$) (Table 3).

3.2.3 Foot wear Footwear and risk of infection with gastrointestinal parasites

We further endeavored to examine the effect of wearing shoes on gastrointestinal parasite infections among the inmates. Among those who were infected and had their feet covered with shoes 17(28.4%) while other infected inmates do not cover their feet. These differences were not statistically significant ($p>0.05$) (Table 3).

3.2.4 Washing hands after contact with soil and risk of infection with gastrointestinal parasites

In additional addition, the effect of washing hands after contact with soil on gastrointestinal parasitic infections among the inmates were was scrutinized. Among those who were infected washed their hands after contact with soil 16(33.3%) while other infected inmates did not wash after contact with soil. These were statistically significant ($p<0.05$)(Table 3).

Table 3: Association between Risk factors and gastrointestinal parasites

	Distribution of Gastrointestinal parasites							Chi square Chi-square (X ²)	p-value
	AL N(%)	HK N(%)	TT N(%)	Negative Cases N(%)	Positive Cases N(%)	Total examined N(%)			
Use_of_drug_in_the_last_three_months									
Yes	10(26.3)	0(0.0)	4(10.5)	24(63.2)	14(36.8)	38(100)	10.764	0.00	Comment [FIMH8]: P-value should be revised
No	0(0.0)	2(8.0)	3(12.0)	20(80.0)	5(20.0)	25(100)			
Total	10(15.9)	2(3.2)	7(11.1)	44(69.8)	19(30.2)	63(100)			
Handwashing_with_soap									
Yes	8(16.0)	2(4.0)	7(14.0)	33(66.0)	17(34.0)	50(100)	2.156	0.00	Comment [FIMH9]: P-value should be revised
No	2(15.4)	0(0.0)	0(0.0)	11(84.6)	2(15.4)	13(100)			
Total	10(15.9)	2(3.2)	7(11.1)	44(69.8)	19(30.2)	63(100)			
Putting_on_shoes_outside									
Yes	10(16.7)	0(0.0)	7(11.7)	43(71.6)	17(28.4)	60(100)	12.794	0.00	Comment [FIMH10]: P-value should be revised
No	0(0.0)	2(66.7)	0(0.0)	1(33.3)	2(66.7)	3(100)			
Total	10(15.9)	2(3.2)	7(11.1)	44(69.8)	19(30.2)	63(100)			
Washing_hands_after_contact_with_soil									
Yes	10(20.8)	2(4.2)	4(8.3)	32(66.7)	16(33.3)	48(100)	5.242	0.00	Comment [FIMH11]: P-value should be revised
No	0(0.0)	0(0.0)	3(20.0)	12(80.0)	3(20.0)	15(100)			
Total	10(15.9)	2(3.2)	7(11.1)	44(69.8)	19(30.2)	63(100)			

3.3 Discussion

The incidences of gastrointestinal parasitic infection due to the triad of *A. lumbricoides*, *T. trichiura*, and hookworms as observed in this study has have been reported among children by various authors from different parts of the country [20-23]. The study revealed that, the overall prevalence of the associated gastrointestinal parasitic infection was 30.2%. The prevalence rate of gastrointestinal parasites among orphanages varies with different geographical locations. In Ebonyi, Plateau, Edo, and Anambra State States, the prevalence of 27.9% 24.6%, 20.7%, and 13.8% were observed by Achi et al. [24], Okolo

and John [25], Nwaneri and Omuemu [26] and Oluboyo et al [27] respectively. These prevalence rates appeared to be relatively lower than the prevalence rate recorded in this study. On contrary, Al-Shibani et al.[28] had a higher prevalence rate of 62.7% in Sana'a City, Yemeni. This difference in prevalence rate could be as a result of unhygienic habits practiced by these children and lack of care and food handling from some of their attendants.

The overall prevalence according to gender among the inmates, females had significantly more gastrointestinal parasitic infections 13(68.4%) compared to males 6(31.6%) and the difference was statistically significant ($p<0.05$). This observation confirms unequal exposure to helminths/helminth eggs among the study population indicating more female susceptibility to gastrointestinal parasitic infections. However, the higher prevalence of gastrointestinal parasitic infections in female children was an indication of poor personal hygiene among the females. This was maintained by a previous study by Hailegebrie [29], who separately reported high prevalence of gastrointestinal parasites among females than males due to their activities. On the contrary, other studies elsewhere recorded high prevalence in males than in females in Southwest Nigeria and Southern Mali [30,31,32]. Prevalent rates in some studies were not significant statistically as recorded by Achi et al. [24], Abah and Arene [13], Udensi et al. [33], and Nwaneri and Omuemu [26].

Age-related prevalence were/was common among of 11-15 years; had 17(44.8%) of gastrointestinal parasitic infection which was statistically significant ($p<0.05$) and then declined among the oldest age group. This report is not in accordance with studies by Okpala et al. [34], El-Nadi et al. [35], and Mekonnen and Ekubagewargies [36] whose age-related prevalence were among 2-3 years age range. The high level of physical activity exhibited by children of this age group exposes them to a greater risk of infection than the others because of their multiple routes of exposure.

Ascaris lumbricoides was/were found to be the most prevalent 10(15.9%), followed by *T. trichiura* and hookworm 7(11.1%) and 2(3.2%) respectively. There was a statistically significant difference in the prevalence of the gastrointestinal parasites among the age groups ($p<0.05$). High/The high prevalence of gastrointestinal parasites among the inmates is a reflection of reflects the level of personal/community hygiene in the living environment. The presence of *A. lumbricoides*, and *T. trichiura/trichiura* observed at the home was indicative of fecal-oral transmission which may be related to proper washing of fecal contaminated hands by the children after the use of their toilets. Hookworm seen in samples were/was indicative of regular bare body contact with the soil by the children, through which cysts/larvae of the parasite bear through the skin and from fruits and vegetables.

With regards to the educational level of the inmates, children who had primary education were found to have a higher prevalence of GIP infections than those with secondary education. This report was found to be significant ($p<0.05$). This report agrees with the studies by Omotola and Ofoezie [37] among school children in Osun state. This could probably be due to the physical activities of this level of children.

The present study has shown that despite the use of drug/drugs in the last three months by some of the inmates still revealed high positivity with gastrointestinal parasitic infections compared to those who were not given preventive therapy. Similar findings were reported in studies conducted elsewhere in Nepal by Bhattachan et al. [38]. This could be as a result because of a lack of monitoring during drug administration by the care-givers/caregivers. This risk factor was found to be statistically significant ($p<0.05$).

In the aspects of hand washing habits inmates who claimed to carry out this activity presented gastrointestinal parasites than to inmates who had no hand washing habits. This could be as a result because of improper monitoring in the side of the care-givers/caregivers. Our reports were not in connection with that carried out by Pasaribu et al [39] in Indonesia. Statistical significance of this risk factor was attained ($p<0.05$).

The practice of wearing off/foot-wear/footwear by inmates was also high with gastrointestinal parasites than those who do not put on foot-wear/footwear. Similar findings were reported in Malaysia by Ahmed et al. [40]. This activity was not statistically significant ($p>0.05$).

Inmates who had their hands washed after being in contact with soil were infected with gastrointestinal parasites more than inmates who had no such habits. This might be due to the fact that because moist soil creates an environment conducive for a high prevalence of intestinal parasites [41,42] and improper handwashing technique/techniques were missing. This report agrees with the findings of Pasaribu et al. [39]. Statistical significance was accomplished ($p<0.05$).

The prevalence of gastrointestinal parasitic infection in this study was high and was significantly associated with risk factors. The methods of prevention of the infection along with chemotherapy

intervention and training of **care-givers/caregivers** on the importance of monitoring and follow-up should be adopted **in order to** interrupt transmission among the inmates Port Harcourt **children/children's** home.

References

1. Saki J, Khademvatan S, Masoumi K, Chafghani M. Prevalence of intestinal parasitic infections among food handlers in Khuzestan, Southwest of Iran: A 10-year retrospective study. *African J Microbiol Res.* 2012;6(10):2475-80.
2. Tyoalumun K, Abubakar S, Christopher N. Prevalence of Intestinal Parasitic Infections and their Association with Nutritional Status of Rural and Urban Pre-School Children in Benue State, Nigeria. *International Journal of MCH and AIDS.* 2016;5(3):146-152.
3. De Silva N, Brooker S, Hotez P, Montresor A, Engels D, Savioli, L. Soil transmitted helminth infections: updating the global picture. *Trends Parasitol.* 2003;19(12):547- 51.
4. Štrkolcová G, Mravcová K, Barbušínová E, Mucha R, Várady M, Goldová M. Prevalence of intestinal parasites in Children Living in Various Living Conditions in Slovakia. *J Pediatr Perinatol Child Health.* 2019;3(3):174-85. DOI: 10.26502/jppch.74050028.
5. Kiani H, Haghghi A, Salehi R, Azargashb E. Distribution and risk factors associated with intestinal parasite infections among children with gastrointestinal disorders. *Gastroenterol Hepatol Bed Bench.* 2016;9(Suppl. 1):S80-S87.
6. Hajissa K, Abd Elhafiz M, Abd All T, Zakeia M, Eshag HA, Einzer E, et al. Prevalence of *Entamoeba histolytica* and *Giardia lamblia* among schoolchildren in Um-Asher Area, Sudan. 2020. <https://doi.org/10.21203/rs-3.rs-61761/v1>.
7. Castellanos-Gonzalez A, White A Jr, Melby P, Travi B. Molecular diagnosis of protozoan parasites by Recombinase Polymerase Amplification. *Acta tropica.* 2018;182:4-11. <https://doi.org/10.1016/j.actatropica.2018.02.002> PMID: 29452112.
8. Organization WH. Working to overcome the global impact of neglected tropical diseases: first WHO report on neglected tropical diseases: World Health Organization; 2010.
9. Hajissa K, Islam MA, Sanyang AM, Mohamed Z (2022) Prevalence of intestinal protozoan parasites among school children in Africa: A systematic review and meta-analysis. *PLoS Negl Trop Dis* 16(2): e0009971. <https://doi.org/10.1371/journal.pntd.0009971>
10. Opara K, Udoidung Okon D, Edosomwan E, Udoh A. The Impact of Intestinal Parasitic Infections on the Nutritional Status of Rural and Urban School-Aged Children in Nigeria. *International Journal of MCH and AIDS.* 2012;1(1):73-82.
11. Miller L, Chan W, Comfort K, Tirella L. Health of children adopted from Guatemala: comparison of orphanage and foster care. *Pediatr.* 2005;115(6):e710-7.
12. Michael EI, Abah AE, Marcus P. Prevalence of Gastro-Intestinal Parasitic Infection among School Children in Port Harcourt City Local Government Area. *Nigerian J Bacteriol Parasitol.* 2017;8(5):1-4.
13. Abah AE, Arene FOI. Status of intestinal parasitic infections among primary school children in Rivers State, Nigeria. *J Parasitol Res.* 2015;Article ID937096:7 pages.
14. Wokem GN, Chukwu C, Nwachukwu CB. Prevalence of intestinal parasites seen in HIV Sero-Positive subjects in Port Harcourt, Nigeria. *Nigerian J Parasitol.* 2008;29(2):115-120.
15. Godden, W. (2004). Sample size formulas. <http://williamgodden.com/samplesizeformula>.
16. Kuo HY, Chiang DH, Wang CC, Chen TL, Fung CP, Lin CP, et al. Clinical significance of *Blastocystis hominis*: Experience from a medical center in northern Taiwan. *J Microbiol Immunol Infect.* 2008;41(3):222-6.
17. Wakid MH. Distribution of intestinal parasites among food handlers in Jeddah, Saudi Arabia. *J Parasit Dis.* 2006;30(2):146-52.
18. Garcia LS. Washington: ASM Press; 2007. *Diagnostic Medical Parasitology.*

19. Wakid MH, Azhar EI, Zafar TA. Intestinal parasitic infection among food handlers in the Holy City of Makkah during Hajj Season 1428 Hegira. *JKAU Med Sci.* 2009;16(1):39-52.
20. Amaechi E, Ohaeri C, Ukpai OM. Prevalence of helminthiasis among school children in some rural communities of Abia State, Nigeria. *Animal Res Int.* 2013;10(3):1817-25.
21. Asaolu SO, Ofoezie IE. The role of health education and sanitation in the control of helminth infections. *Acta Trop.* 2003;86(2-3):283-94.
22. Ugbomoiko US, Ofoezie IE. Multiple infection diagnosis of intestinal helminthiasis in the assessment of health and environmental effects of development projects in Nigeria. *J Helminthol.* 2007;81(3):227-31.
23. Olaniran O, Ajayi OO, Odetoyin BW, Afolayan DO, Awoyeni EA, Adekunle OT, et al. Prevalence of Soil Transmitted Helminthes among School Children in Central Local Government Area Ile-Ife Osun State. Nigeria. *Int J Novel Res Interdiscip Stud.* 2015;2(2):8-13.
24. Achi EC, Njoku OO, Nnachi AU, Efunshile AM, Mbah JO, Aghanya IN, Nnemelu PO. Prevalence of Intestinal Parasitic Infections Among Under Five Children in Abakaliki Local Government Area of Ebonyi State. *Eur J Pharm Med Res.* 2017;4(4):218-22.
25. Okolo, S. N, and John, C. (2006). Nutritional status and intestinal parasitic infestation among rural Fulani children in Vom, Plateau State. *Nigerian J Paediatr.* 2006;33(2):47-55.
26. Nwaneri DU, Omuemu VO. Intestinal helminthiasis and nutritional status of children living in orphanages in Benin City, Nigeria. *Nigerian J Clin Pract.* 2013;16(2):243-8.
27. Oluboyo BO, Enweani IB Ekejindu IM, Oluboyo AO. Prevalence of some intestinal parasitic infections in relation to body mass index of children resident in orphanages In Anambra State, Nigeria. *Eur Sci J.* 2014;10(36):222-9. ISSN: 1857-7881 (Print) e - ISSN 1857- 7431
28. Al-Shibani LA, Azazi AA, El-Taweel HA. Cryptosporidiosis and other intestinal parasites in 3 Yemeni orphanages: prevalence, risk, and morbidity. *J Egypt Soc Parasitol.* 2009;39(1):327-37.
29. Hailegebrie T. Prevalence of intestinal parasitic infections and associated risk factors among students at Dona Berber primary school, Bahir Dar, Ethiopia. *BMC Infect Dis.* 2017;17:362.
30. Oyewole F, Ariyo F, Sanyaolu A, Oyibo WA, Faweya T, Monye P, et al. Helminthic reduction with albendazole among children in riverine communities of Nigeria. *Southeast Asian J Trop Public Health.* 2002;33(2):214-7.
31. Adeyeba OA, Akinlabi AM. Intestinal parasitic infections among school children in a rural community Southwest Nigeria. *Nig J Parasitol.* 2002;23(1):11-18.
32. Behnke JM, De Clercq D, Sacko M, Gilbert FS, Ouattara DB, Vercauteren J. The epidemiology of human hookworm infections in the southern region of Mali. *Trop Med Int Health.* 2001;5(5):343-54.
33. Udensi JU, Mgbemena IC, Emeka NI, Ugochukwu MG, Awurum IN. Prevalence of Intestinal Parasites Among Primary School Children in Three Geopolitical Zones of Imo State, Nigeria. *Science Journal of Public Health. Special Issue: Who Is Afraid of the Microbes.* 2015;3(5-1): 25-28. doi: 10.11648/j.sjph.s.2015030501.15
34. Okpala HO, Josiah SJ, Oranekwulu MU, Ovie, EG. Prevalence of intestinal parasites among Children in Day Care Centres in Esan West Local Government Area, Edo State, Nigeria. *Asian J Med Sci.* 2014;6(4):34-9.
35. El-Nadi NAF, Omran EK, Ahmed NS, Fadel EF. Current status of gastrointestinal parasites among elementary school children in Sohag, Egypt. *J Adv Parasitol.* 2017;4(2):3340.
36. Mekonnen HS, Ekubagewargies DT. Prevalence and factors associated with gastrointestinal parasites among under-five children attending Woreta Health Centre, Northwest Ethiopia. *BMC Infect Dis.* 2019;19:256
37. Omotola OA, Ofoezie IE. Prevalence and Intensity of Soil Transmitted Helminths among School Children in Ifetedo, Osun State, Nigeria. *J Bacteriol Parasitol.* 2019;10(1):352. doi:10.4172/2155-9597.1000352
38. Bhattachan B, Panta YB, Tiwari S, Thapa Magar D, Sherchand JB, et al. Intestinal Parasitic Infection among School children in Chitwan district of Nepal. *J Inst Med.* 2015;37(2):79-84.
39. Pasaribu AP, Alam A, Sembiring K, Pasaribu S, Setiabudi D. Prevalence and risk factors of soil-transmitted helminthiasis among school children living in an agricultural area of North Sumatera, Indonesia. *BMC Public Health.* 2019;19:1066 <https://doi.org/10.1186/s12889-019-7397-6>

40. Ahmed A, Al-Mekhlafi HM, Choy SH, Ithoi I, Al-Adhroey AH, Abdulsalam AM, et al. The burden of moderate-to-heavy soil-transmitted helminth infections among rural malaysian aborigines: an urgent need for an integrated control programme. *Parasit Vectors*. 2011;4:242.
41. Abera B, Alem G, Yimer M, Herrador Z. Epidemiology of soil-transmitted helminths, *Schistosoma mansoni*, and haematocrit values among schoolchildren in Ethiopia. *J Infect Dev Ctries*. 2013;7(3):253-60.
42. Feleke BE. Nutritional status and intestinal parasite in school age children: a comparative cross-sectional study. *Int J Pediatr*. 2016;Article ID 196212, page 8.