

Demographic profile & pattern of ocular and adnexal injuries from Sloth Bear Maul: A study in tertiary centre of Bilaspur District, Chhattisgarh.

Abstract :

This study was done in all cases of bear attacked patients admitted in our tertiary centre Chhattisgarh institute of medical science (CIMS) who suffered ocular and adnexal injuries. Sloth bear (*Melursus ursinus*) is found to be native in many forest areas of the state of Chhattisgarh including North Bilaspur Forest Division (Nbfd). The study was conducted in the department of ophthalmology, Chhattisgarh institute of medical science (CIMS) Bilaspur Chhattisgarh for the duration of 3 years (January 2018 to December 2020). All patients who were admitted in our institute with ocular and/or adnexal injuries from bear attack have been included in this study. 27 patients were attacked by sloth bear in the forest areas of Bilaspur division and referred to our tertiary centre. These patients had ocular and adnexal injuries. These bears were found roaming in the forest areas and also at places of human and animal habitats. 67% (18 of 27) patients were attacked in the forest areas, while some were attacked in the crop fields (15%), home courtyard (7%) and near water bodies (11%). Sloth bears are one of the most aggressive bears who have attacked human populations. The injuries cause severe visual loss and disfigurement to eyeball and adnexa. Proper measures are needed to be implemented to prevent such cases in future.

Keywords: Sloth Bear, adnexal injuries, animal habitats, **ocular injury**

INTRODUCTION:

Sloth bear (*Melursus ursinus*) is found to be native in many forest areas of the state of Chhattisgarh including North Bilaspur Forest Division (Nbfd).¹ These forest areas of Nbfd includes Marwahi and Pendra range. Sloth bears are also habitats of Koriya and Korba districts of the state of Chhattisgarh. These forest areas are tropical, dry and deciduous. They have patchy and fragmented areas with agricultural field, and also have villages with higher number of human and cattle population in outer protected parts of the forest. Largest population of sloth bear in India (about 10,000)¹ has been reported in these forests (Pendra-Marwahi, Korba and Koriya districts are one of them having bears in higher numbers). These forest areas have hillocks and boulders which offers den sites for the bears.²

The sloth bears are medium sized species with weight of 55 to 145 kilograms and height of 4.7 to 6.3 feet's (figure 1). They have strong claw nails of 3 to 4 inch and long incisors teeth's.³ These bears have been found to be more active in winter season and in early morning hours during which they use to return in their den after spending time in crop fields and water resources.⁴ Human population who habitats

outside the protected areas of forest, use forest materials for their survival. They visit to forest for farming purpose, defecation, water collection and collection of firewood, mahua (*Madhuca longifolia*), bel (*Aegle marmelos*) and tendu (*Diospyros melanoxylon*).¹ And get encountered here by the sloth bears.

Figure 1: sloth bear (*melursus ursinus*).



These sloth bears have been reported aggressive in nature and can attack humans without provocation. They have been found hidden behind trees or climbed over trees before attack and use their claws and teeth to attack. Humans can be attacked by a single bear or group of bears. These attacks to human beings may vary from mild injuries to severe fatal trauma with many reported mortalities. Head, face and eyes have been found to be the most attacked parts of body in many of the injuries caused by bear encounter.

Ocular and adnexal injuries have been seen in many bear attacked patients. These injuries can involve various parts of eye including eyelids laceration, lacrimal apparatus laceration, degloving injuries of adnexa, fractured orbital walls, corneoscleral lacerations and even traumatic enucleation of eyes with optic nerve laceration. It can lead to severe visual loss and also disfigurement of eye and face. This study was done in all cases of bear attacked patients admitted in our tertiary centre Chhattisgarh institute of medical science (CIMS) who suffered ocular and adnexal injuries.

MATERIALS AND METHODS:

Our study was conducted in the department of ophthalmology, Chhattisgarh institute of medical science (CIMS) Bilaspur Chhattisgarh for the duration of 3 years (January 2018 to December 2020). All patients who were admitted in our institute with ocular and/or adnexal injuries from bear attack have been included in this study. These Patients also had injuries to other body parts including: head, oro-maxillo-facial, ENT, neck, chest, abdomen or others. Proper history of the presenting illness was taken in terms of mode, time and place of attack & activity involved during the attack. Total number of bears involved was also noted.

Complete ocular examination was done in terms of type of injury, parts involved, visual assessment, anterior segment and posterior segment examination. X-ray and CT scan of the orbit have been done to look for any bony involvement (fractures of orbital walls/soft tissue entrapments and optic nerve impingement from bony fragment). Patients were managed as per the type of injury. All those patients who had eyelid and canalicular laceration were managed by suturing with 5-0 vicryl and 5-0 silk sutures while those having tissue loss underwent reconstructive surgery. Patients who presented with corneal, scleral or corneo-scleral laceration were repaired with 10-0 nylon suture under poor or nil visual prognoses. Those having fractures of the orbital walls were referred to oculoplastic surgeon for further management. All patients received anti-rabies vaccine as per schedule and rabies immunoglobulin. Those having injuries to other body parts were managed by the concerned departments.

Comment [u1]: In terms of are too many kindly review

Comment [u2]: as par???

Inclusion criteria: All patients of bear attack with ocular or adnexal injuries.

Exclusion criteria:

- 1) Attack from other animals
- 2) Patients with bear attack not involving ocular or adnexal area.

RESULT:

In our present study from January 2018 to December 2020, total 27 patients had been admitted in the tertiary centre Chhattisgarh institute of medical science (CIMS) Bilaspur, CG. All patients included in the present study had ocular and/or adnexal trauma from cloth bear attacks. Patients also suffered injuries to the other body parts (head, neck, oro-maxillo-facial, Ear, Nose, Throat, chest and others).

Mode of attack:

Comment [u3]: Not necessary here

In all of the 27 patients was from bear claw nails either attacked from front or behind the patients. The bears were found to be hidden either behind the trees or climbed over trees before attack so they were not visible to the patients. None of the patients had given history of injury from bear's teeth.

Time of attack:

Comment [u4]: Review this sentence

Out of these 27 patients, 11 (40.74%) persons had been attacked in the early morning hours of 4 am to 7 am, while 13 (48.15%) patients had been attacked in the evening hours of 6 pm to 9 pm. Only 3 (11.11%) patients were attacked in night time of 10pm to 2 am.

Place of attack:

All patients were residents of forest areas of Bilaspur division (Pendra-Marwahi, koriya and korba) and had been referred from these places. The distribution of locations at which patients had been attacked is shown in table 1.

Table 1: Locations of bear attack.

Place	Number	%
Forest	18	67
Crop fields	4	15
Home courtyard	2	7
Near water bodies	3	11
Total	27	100

Human Activities at the site of attack(Table2):

Majority (48.15%) of patients (13 of 27) attacked have visited forest area for collection on firewood, mahua, tendu and bel. Two patients were attacked when they were sleeping at open areas of house. Three patients got injured while protecting their cattle from bear attack.

Table 2: Activities at site of bear attack.

Activity	Number	%
Forest material collection	13	48.15
Water collection	2	7.41
Defecation	2	7.41
Farming	5	18.51
Sleeping at home	2	7.41
Cattle places	3	11.11
Total	27	100

Demographic profile of patients (Table 3):

Majority of the affected patients were males(81%). ~~It can be due to higher outdoor activities in male population in attacked area.~~ Major age group affected was young adults of 31 to 60 years age group (13 of 27)

Comment [u5]: Not for result but in discussion

Table 3: Age and gender distribution of patients with ocular and adnexal trauma from bear attack.

Age (years)	Number of patients (%)	Male	female
<30	4(15)	4(15%)	0
31-60	13(48)	10(37%)	3(11%)
>60	10(37)	8(30%)	2(7%)
total	27(100)	22(81%)	5(19%)

Parts of ocular and adnexa involved (Table 4):

Face has been seen to be the most common part attacked in encounters with sloth bears. Ocular and adnexal involvement has been seen from eyelid lacerations and corneo-scleral perforations to even complete enucleation of eyeball with ~~optic nerve lacerations.~~ These injuries have even caused complete loss of vision in the affected eye.

Comment [u6]: ?????

Table4: Ocular and adnexal parts injured in bear attack.

Part involved	Number	%
Laceration of eyelids.	11	40.74
Laceration of Eyelid and lacrimal apparatus	12	44.44
Eyelid avulsion	4	14.82
Corneal tear with prolapsed uvea.	3	11.11
Corneo-Scleral tear with	9	33.33

prolapsed uvea and vitreous		
Traumatic enucleation with optic nerve injury	3	11.11

Figure2: Eyebrows with eyelids laceration from bear attack.



Figure3: Corneo-scleral laceration with eyelid laceration from bear attack.



Figure4: Traumatic enucleated eye with optic nerve laceration from bear attack:

Comment [u7]: This can actually be called ruptured globe rather than enucleation

Comment [u8]: ?????????



Figure 5: eyelids avulsion both eye and traumatic enucleation left eye with fracture floor orbit from beat attack.



Figure 6: right eyelid multiple laceration(sutured) with fracture floor and medial wall of orbit.



Visual acuity:

Vision of the injured eye was severely affected in cases of corneo-scleral tear with prolapsed uveal tissue and vitreous ranging from 6/60 to no perception of light. Also cases of traumatic enucleation have no perception of light. 25.93% of patients suffered from no perception of light in the affected eye (Table 5). While 18.51% patients had visual acuity reduced to only perception of light. 11.11% of patients had retained visual acuity of 6/60 to hand movement. Cases having better visual acuity were those who had involvement of ocular adnexa only and they had either immature senile cataract or corneal opacities.

Table 5: Visual acuity of the affected eye from bear attack.

Visual acuity	Number	%
---------------	--------	---

6/6 to 6/12	5	18.51
6/18 to 6/36	7	25.93
6/60 to hand movement	3	11.11
Perception of light	5	18.51
No perception of light	7	25.93
Total	27	100

Management of cases: All cases were managed according to the type of injury of eye. All patients received intramuscular anti rabies vaccination as per schedule along with IgG immunoglobulin infiltration locally around the wound. Those patients who had laceration of eyelids were managed by repairing with 5-0 vicryl suture and 4-0 silk suture. Corneo-scleral tears were reconstructed with 10-0 nylon sutures and 8-0 vicryl sutures. Injury of lacrimal apparatus was managed by primary suturing. Severely traumatized eye underwent enucleation.

Comment [u9]: Already mentioned somewhere

Discussion:

In our present hospital based study, 27 patients were attacked by sloth bear in the forest areas of Bilaspur division and referred to our tertiary centre. These patients had ocular and adnexal injuries. All of the 27 patients were attacked by the claws of bear. Sloth bear were found to be hidden either behind the trees or climbed over trees before attack. 48.15% (13 of 27) of patients were attacked in the evening hours of 6 pm to 9 pm followed by 40.77% (11 of 27) patients in the morning hours of 4 am to 7 am.

These bears were found roaming in the forest areas and also at places of human and animal habitats. 67% (18 of 27) patients were attacked in the forest areas, while some were attacked in the crop fields (15%), home courtyard (7%) and near water bodies (11%). Majority of the patients (48.15%) were attacked while they visited forest for collection of firewood, bel, mahua and tendu leaves (table 2).

Apart from this, some were attacked while they went for water collection, defecation, farming and protecting cattle from the attacks. Due to higher outdoor activities males (81%) have been attacked more than females (19%). Age groups of 31-60 (48%) years have been affected more followed by 37% of people beyond 60 years (table 3). Attacks from bear have seen to cause

severe injuries to body parts including blinding ocular and adnexal trauma. In our study 11% of cases suffered from traumatic enucleation of eye with optic nerve injury leading to disfigurement and complete visual loss of the affected eye.

All patients suffered Eyelid injuries with lid avulsions in 14.82% patients and 44% of patients had injuries to lacrimal apparatus along with eyelid injuries. 33% patients had corneo-scleral tear with prolapsed vitreous and uveal tissue (table 4). 15 of 27 patients suffered severe visual loss(table 5)

Injuries from bear attack have been reported previously. Patil SB *etal* in 2015 reported 48 cases that were attacked by sloth bear in central india.⁵ Out of these 48 patients, facial injuries were seen in 36 patients of whom 12 patients had eyelid injury and 4 had eyeball injuries. Four patients underwent enucleation surgeries due to severe ocular trauma.

Nath A *etal* in 2020 ~~have~~ studied pattern of bear attack and behavior in Himalayan region.⁶ They found most of patients were of low socioeconomic status. Highest incidence of attack was seen from June to December. Majority of patients suffered from injuries of face and neck.

Bhat TA *etal* reported 21 patients suffering upper limb and facial injuries from bear attack.⁷ These patients underwent multiple orthopedic surgeries following bear encounter.

Bear attacks has been seen in other areas of the world also. Frank RC *etal* in 2006 has reported 7 cases of bear attacks who suffered from various types of injuries including lacerations, fractures and tissue avulsions.⁸ These areas have habitat of grizzly bears. Post operative complications like infections and psychological trauma.

A study was done by Penjoretal from 2015 to 2019 in Bhutan. It has habitats of sloth bear and Himalayan black bear.⁹ They have reported 34 patients who got attacked from bear and suffered injuries to various body parts. They found face to most common affected part in injury including 85% of cases.

Shah AA *etal* in 2010 has described patterns of Bear maul maxillofacial injuries in Kashmir.¹⁰ They found 200 cases of bear attacks by Asiatic Black bear. Out of 200 cases 159 were males. Majority of cases were of age group 30-40 years (110 of 200) followed by 40-60 years (55 of

Comment [u10]: These are your result find. In discussion you relate to other findings or give reason for your result observation.

Comment [u11]: Review this with your result and discuss

Comment [u12]: In relation to your study ???

Comment [u13]: In a study by Penjoretal.....

200). Facial injuries were most commonly (161 of 200 cases) found out of which 13 patients suffered orbital wall fractures.

Comment [u14]: Review and relate to your findings

Conclusion:

Sloth bears are one of the most aggressive bears ~~whohat~~ have attacked human populations. These attacks may be a result of invasion of humans to their forest habits. They have large claws which are used to injure. These injuries can be lethal and person may suffer mortality. Face is the most common site which got injured. Ocular and adnexal injuries have been seen in majority of cases from eyelid lacerations, eyelid avulsions, and corneo-scleral tears to traumatic enucleation. These injuries cause severe visual loss and disfigurement to eyeball and adnexa. Proper measures are needed to be implemented to prevent such cases in future.

Reference

1. Bargali H, Akhtar N, Chauhan NPs. The sloth bear activity and movement in highly fragmented and disturbed habitat in central India. *World Journal of Zoology* 2012; 7: 312-319.
2. Akhtar N, Bargali H, Chauhan NPs. Sloth bear habitat use in disturbed and unprotected areas of Madhya Pradesh India. *Ursus* 2009; 15:203-211
3. Joshi, A. R., Smith, J. L., & Garshelis, D. L. (1999). *Sociobiology of the myrmecophagous sloth bear in Nepal*. *Canadian Journal of Zoology*, 77(11), 1690–1704.
4. Prajapati, Utkarsh; Koli, Vijay Kumar; Sundar, K.S. Gopi (2021). "Vulnerable sloth bears are attracted to human food waste: a novel situation in Mount Abu town, India". *Oryx*. In press.
5. Patil SB, Mody NB, Kale SM, Ingole SD. A review of 48 patients after bear attacks in Central India: Demographics, management and outcomes. *Indian J Plast Surg*. 2015;48(1):60-65. doi:10.4103/0970-0358.155267
6. Nath A, Gaur NK, Vathulya M, Kapoor A, Mir MA, Mago V, Chattopadhyay D. Bear attack pattern and behavior in the Himalayan region: A study from a tertiary care center. *Injury*. 2020 Sep;51(9):2082-2085. doi: 10.1016/j.injury.2020.06.032. Epub 2020 Jun 20. PMID: 32591216.
7. Bhat TA, Gulzar A, Bhat AA, Bhat TA, Ali Z. A review of upper limb injuries in bear maul victims: Consistent pattern and inverse relation in severity with facial and scalp injuries. *Chin J Traumatol*. 2018 Feb;21(1):38-41. doi: 10.1016/j.cjtee.2017.11.001. Epub 2017 Nov 24. PMID: 29402720; PMCID: PMC5857895.

8. Frank RC, Mahabir RC, Magi E, Lindsay RL, de Haas W. Bear maulings treated in Calgary, Alberta: Their management and sequelae. *Can J Plast Surg*. 2006;14(3):158-162. doi:10.1177/229255030601400301.

9. Penjor D, Dorji T. Circumstances of human conflicts with bears and patterns of bear maul injuries in Bhutan: Review of records 2015-2019. *PLoS One*. 2020 Aug 20;15(8):e0237812. doi: 10.1371/journal.pone.0237812. PMID: 32817716; PMCID: PMC7446927.

10. Shah AA, Mir BA, Ahmad I, Latoo S, Ali A, Shah BA. Pattern of bear maul maxillofacial injuries in Kashmir. *Natl J Maxillofac Surg*. 2010;1(2):96-101. doi:10.4103/0975-5950.79208.