

## Assessment of Antibacterial and Gas Chromatography/Mass Spectroscopic analysis (GC-MS) Profile of Purified Essential Oil From Multi-Phases Solvent Extraction Method From Stem Extract of *Aframomum melegueta* [Roscoe] K. Schum

### Abstract

This research work aims to ascertain the composition and potency of the bioactive component of essential oil derived from *Afr. melegueta* stem extract through multi-phase solvent extraction Gas Chromatography /Mass Spectroscopic analysis and antimicrobial assay against selected clinical isolates. *Afr. melegueta* stem extract is used as a remedy against stomachache, diarrhea, and snakebite. The fresh stem of *Afr. melegueta* plant was collected from Owo forest reserve, Ondo State, Nigeria, 500g of each dried and powdered plant sample was weighed separately into corked containers containing 1500ml each of acetone and ethanol, the mixture was shaken vigorously and left for 9 days, Solvent of extraction are n-Hexane (StAMH)(153g), Dichloromethane,(StAMD(103g), Ethyl acetate, (StAME)(45.5g), Methanol (StAMM)(50g). For the StAMD Elution, StAMD was absorbed in silica gel of 200-400 mesh from ZICO-TEK laboratory, GC-MS analysis of this extract was performed using a Perkin Elmer GC Claurus 500 system and gas chromatograph interfaced to a Mass Spectrometer (GC-MS) equipped with Elite-1 fused silica capillary column (30mx1µl was Mdf. Composed of 100% Dimethylpolysiloxane). 17 and 48 compounds were analyzed in the stem extract of purified Essential oil from Multi-phases solvent extraction of stem extract of *Afr.melegueta*. The composition, chemical structure, molecular weight, and molecular formulation of different essential oil of StD+ and StD++ from purified Essential oil. The antimicrobial assessment were performed using Kirby-Bauer disc diffusion method (Agar well diffusion method). *E.coli* and *Staphylococcus aureus* has the highest zones of inhibition of 19,0 mm at 100mg/ml. In contrast, *Klebsiella planticola* and *Citrobacter diversus* have the lowest zones of inhibition of 3.0mm at 12.5mg/ml for StD+ and *Staphylococcus aureus* (12%) and *E.coli* (12%), has the highest. In contrast, *Staphylococcus aureus* (6%), has the lowest percentage frequency distribution of purified essential oil of *Afr.melegueta* (StD+) against selected clinical isolates. *E.coli* has the highest zones of inhibition of 20mm at 100mg/ml, while *Klebsiella planticola* has the lowest inhibition of 1.0mm at 12.5mg/ml for (StD++). In conclusion, the uses and importance of essential oil fractions from *Afr. melegueta* have limitless potential.

.Keyword;*Aframomum melegueta*, Bacterial,Essential oil, Extraction solvent

## Introduction

This plant is a member of the ginger family (Zingiberaceae) and is cultivated in tropical areas of West Africa. The plant seeds are used to flavor foods and as components of traditional African folk medicine. In medieval Europe, they were a highly prized spice that was eventually replaced by black pepper and other spices. Ethnobotanically, the stem extract is used as a remedy against stomachache, diarrhea, and snakebite [1], [2].

In Addition, there are reported studies on the different medicinal use of *Aframomum melegueta*, they are; Antiulcer, Cytoprotective, and Antimicrobial activities as well as the sexual performance-enhancing effects of grains of *paradise*[3],[4],[5],[6]. The aqueous seed extract has been shown to reduce the frequency of abdominal constrictions induced by acetic acid in mice and has significant anti-inflammatory activity [7]. It was later reported that the same extract has peripheral analgesic activity. Additionally, it was suggested that the extract has membrane-stabilizing activity along with antioxidant effects [8],[9].as well as hypotensive and antihypertensive activity in humans [10],[11]. it has been also found that the extract has an effect on the whole-body energy expenditure and visceral fat in humans[12]. However, in some of the literature studies, it should be mentioned that the seed of *Aframomum melegueta* derived essential oil contains humulene and caryophyllene occurred in higher proportions in the volatile oil[13].

The major components of the leaf oil were found to be myrtenyl acetate, isolimonene while caryophyllene oxide, myrtenyl acetate,  $\beta$ -eudesmene, and  $\beta$ -caryophyllene make up the composition of the stem oil [14] whereas the root essential oil comprised of myrtenyl acetate and pinocarvyl acetate. However, the seed is comprised mainly of  $\alpha$ -humulene,  $\beta$ -caryophyllene.

In literature, the major constituents of the leaf essential oil[15]was identified as sabinene, $\alpha$ -pinene and  $\beta$ -caryophyllene. The seed essential oil of *A. melegueta* presents a characteristic composition with  $\beta$ -caryophyllene,  $\alpha$ -humulene, and their epoxides as main constituents [16]. Eugenol occurred in abundance in the oil [17], $\beta$ -Pinene predominates in the essential oils of the leaves and seeds of *Afr. melegueta* [18]. GC/MS analysis of the hexane and methanol extracts of *Afr. melegueta* seed yielded gingerol, zingiberone, paradol, *trans*-6-shogaol, *cis*-isoelemicin,

$\beta$ -bisabolene,  $\alpha$ -guaiene, aromadendrene, *trans*- $\beta$ -farnesene and geraniol. The essential oil of *A. melegueta* displayed insect repellency against *Rhyzopertha dominica* [19] antimicrobial activity [20], antifungal effect [21], moderate inhibition of acetyl-cholinesterase [22], antioxidant [23], metabolites exhibit polypharmacology against SARS-CoV-2 drug targets [24].

## 2.0 Material and Method

### Collection of fresh stem *Aframomum melegueta* [Roscoe] K. Schum

The fresh stem of *Aframomum melegueta* [Roscoe] K. Schum plant were collected from Owo forest reserve, Ondo State, Nigeria The experimental site is located between coordinates 6.96879 and 5.5626 and an altitude of 415 m on the month June 20<sup>th</sup>, 2020 and the plant samples were authenticated at the Department of Plant Science and Biotechnology, Adekunle Ajasin University, Akungba Akoko, Ondo State, Nigeria

### Preparation of fresh stem *Aframomum melegueta* [Roscoe] K. Schum

For the extraction of each plant part, 500g of each dried and powdered plant sample was weighed separately into corked containers containing 1500ml each of acetone and ethanol, the mixture was shaken vigorously and left for 9 days. The mixture was in a ratio of 1:1. All mixtures were filtered using sterile Whatman No. 1 filter papers, and the filtrates were collected directly into sterile crucibles. The filtrate was extracted using a soxhlet extractor, and the residues obtained were kept at room temperature [23].

### Extraction of fresh stem *Aframomum melegueta* [Roscoe] K. Schum

#### Multi-Phase Solvent Extraction of fresh stem of *Aframomum melegueta* [Roscoe] K.

##### Schum Bioactive Compound (Essential Oil Fraction)


*Afr. melegueta* was serially extracted using the multiphase extraction method at room temperature for 72 hours varying solvents based on their polarity using organic solvents. Solvent of extraction are: n-Hexane,(StAMH)(153g),Dichloromethane,(StAMD(103g),Ethyl acetate, (StAME) (45.5g), Methanol,(StAMM)(50g). For the StAMD Elution, StAMD was absorbed in silica gel of 200-400 mesh from ZICO-TEK laboratory. The following table shows detailed


multiphase extraction of the essential fraction of *Afr. melegueta*. A schematic extraction tree follows the table below.

**Table 1 : Details of Multi-Phase Solvent Extraction Process. (Essential Oil Phase/Fraction)**

n-Hex	DCM	EtOAc	Volume(mL)	Test-tube Numbering
100	-	-	300	500 Conical flask
90	10	-	200	1-5
50	50	-	200	6-14
45	55	-	100	15-20
30	70	-	100	21-27
20	80	-	400	28-57
-	100	-	100	58-67
-	90	10	300	68-79
-	50	50	100	80-86
-	-	100	200	87-97

**Flowchart 1 :Flowchart showing the study processing**





**Chart 1 ;Schematic diagram of multi-phase extraction of bioactive compound (Essential oil,SD<sup>+</sup> and SD<sup>++</sup>)**

d,33 - 35 e,37 - 51 f. a - f were all chlorophyll and were discarded and tagged A.52 - 55(B),56 - 63(C),64 - 69 (D),70 - 74(E),75 - 84(F),85 - 92(G),93 - 97(H). Further bulking as guided by TLC developed by mobile phase DCM: EtOAc 2 : 8; EtOAc 100%. B, C, and D bulked and labeled D<sup>+</sup> pure compound I. E and F bulked as D<sup>++</sup>, G and H bulked to H<sup>-</sup> - discarded due to insufficient quantity.

## **Gas chromatograph and mass spectroscope (GC-MS) Analysis of fresh stem extract of *Afr. melegueta* [Roscoe] K. Schum**

GC-MS technique was used in this study to identify the components present in the extract of entire parts of *Aframomum melegueta* [Roscoe] K. Schum. GC-MS technique was carried out at the School of Chemistry and Physics, Westville Campus, University of KwaZulu-natal Durban, South Africa. GC-MS analysis of this extract was performed using a Perkin Elmer GC Claurus 500 system and gas chromatograph interfaced to a Mass Spectrometer (GC-MS) equipped with Elite-1 fused silica capillary column (30mx1µl was Mdf. Composed of 100% Dimethylpolysiloxane). For GC-MS detection, an electron ionization energy system with ionization energy of 70eV was used. Helium gas (99.999%) was used as the carrier gas at a constant flow rate of 1ml/min. and an injection volume of 2µl was employed (Split ratio of 10:1). The injector temperature was 250°C. The oven temperature was programmed from 110°C (isothermal for 2min.), with an increase of 10°C/min to 200°C, then 5°C /min. to 280°C, ending with a 9min. isothermal at 280°C. Mass spectra were taken at 70eV; a scan-interval of 0.5 seconds and fragments from 45 to 450 Da. Total GC running time was 36 min. The relative percentage amount of each component was calculated by comparing its average peak area to the total area. Software adapted to handle mass spectra and chromatograms was a Turbo mass Ver5.2.0. Compound identification was obtained by comparing the retention times with those of authentic compounds and the spectral data obtained from the library data of the corresponding compounds. The given sample was extracted with ethyl acetate and analyzed in GC-MS for different components [23, 24, 25].

### **Identification of components**

The identity of the components in the extract was assigned by the comparison of their retention time and mass spectra fragmentation patterns with those stored in the computer library and also with published literature. NIST library sources were also used for matching the identified components from the plant material.

### **Standardization of Plant Extracts for antimicrobial assay.**

At aseptic conditions, the extracts were reconstituted by adding 1g of each extract to 2.5ml of DMSO and 7.5ml of sterile distilled water to make 100mg/ml. The serial concentration was prepared to get concentration of 50mg/ml, 25mg/ml and 12.5mg/ml respectively [25,26]

### **Test Organisms**

The selected clinical test isolates used were: *E.coli*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *P.aeruginosa*, *Staphylococcus aureus*, *Coryne bacterium cystitidis*, *Klebsiella planticola*, *Salmonella choleraesuis*, *Citrobacter freundii*, *Citrobacter diversus*

### **Standardization of Test Organisms**

The test organisms used were obtained from the stock culture of the laboratory of the Department of Microbiology, Adekunle Ajasin University, Akungba-Akoko, Ondo State, Nigeria. The test organisms were sub-cultured to obtain pure cultures of the organisms. The broth cultures of the test organism were prepared according to 0.5 McFarland's standard [26].

### **Antimicrobial Screening of Purified Essential oil of *Aframamum melegueta*.(StD+ and StD++) Extracts against selected clinical isolates using Kirby-Bauer method(Agar well diffusion method)**

The agar well diffusion method according to Osuntokun *et al* was used. The overnight broth culture of the respective bacteria strains was adjusted to 0.5 McFarland standard. Mueller-Hinton agar plates were swabbed using sterile cotton swabs with the adjusted broth culture of the respective bacteria strains. Wells (6 mm in diameter) were made equidistance in each of the plates using a sterile cork borer. 100 µl (0.1 ml) of each concentration of the extract were respectively introduced into the wells using sterile automatic pipettes, with the stock solution in the center well with different concentrations of the extracts (50, 25, and 12.5mg/ml). The plates were allowed to diffuse at room temperature for 2 hours and were incubated at 37°C for 24 hours for the bacterial isolates and 24°C for 48 hours for the fungal isolates. The zones of inhibition were measured to the nearest millimeter (mm) using a standard transparent meter rule. The antibacterial activity was expressed as the mean zone of inhibition diameters (mm) produced by the plant extract [23,25].

### 3.0 Result

The following represented the result obtained during the course of this research work. two tables (2 and 3,) state the spectral analysis of *Aframomum melegueta*, figure 1-11) which demonstrate the antibacterial activity zones of inhibition and percentage frequency distribution of purified essential oil against the clinical organisms/isolates.

Table 2; Gas chromatography and mass spectroscopy (GC-MS) analysis of *Aframomum melegueta* stem extract of purified essential oil stem extract (*StD* +), table 3; Gas chromatography and mass spectroscopy(gc-ms) analysis of *Aframomum melegueta* stem extract of purified essential oil stem extract (*StD*++), Fig 1; Antibacterial activity of purified essential oil of *Aframomum melegueta* (*StD*+) stem extracts against selected clinical isolates, Fig 2; Percentage frequency distribution of antibacterial activity of *Aframomum melegueta* (*StD* +) stem against clinical isolates at (*StD* +)100mg/ml, Fig 3; Percentage frequency distribution of antibacterial activity of *Aframomum melegueta* stem against clinical isolates at (*StD*+) 50mg/ml, Fig 4; Percentage frequency distribution of antibacterial activity of *Aframomum melegueta* stem against clinical isolates at (*StD* +) 25mg/ml, Fig 5; Percentage frequency distribution of antibacterial activity of *Aframomum melegueta* stem against clinical isolates at (*StD* +) 12.5mg/ml, Fig 6; Antibacterial activity of purified essential oil of *Aframomum melegueta*.( *StD* ++ )stem extracts against selected clinical isolates, Fig 7; Percentage frequency distribution of antibacterial activity of *Aframomum melegueta* stem against clinical isolates at (*StD* ++ ) 100mg/ml, Fig 8; Percentage frequency distribution of antibacterial activity of *Aframomum melegueta* stem clinical isolates at (*StD* ++ ) 50mg/ml, Fig 9; Percentage frequency distribution of antibacterial activity of *Aframomum melegueta* stem against clinical isolates at (*StD* ++ ) 25mg/ml, Fig 10; Percentage frequency distribution of antibacterial activity of *Afr. melegueta* stem against clinical isolates at (*StD* ++ ) 12.5mg/ml. Spectral 1; Gas chromatography and mass spectroscopy (GC-MS)spectra analysis of *Afr.melegueta* purified essential oil stem extract (*StD*++).Spectral 2; Gas chromatography and mass spectroscopy (GC-MS) spectra analysis of *Afr.melegueta* stem extract of purified essential oil stem extract (*StD* ++).Table 2 and 3 represents the Gas chromatograph and mass spectroscopy (GC-MS) Spectra Analysis of *Afr. melegueta* purified essential oil (*StD*+ and *StD*++),

Figure 1 represents percentage frequency distribution of antibacterial activity against clinical isolates at (StD+) and SD++) at 100,50,25, and 12.5mg/ml.

**Table 2;** Gas chromatography and mass spectroscopy (GC-MS) analysis of *Aframomum melegueta* stem extract purified Essential oil (StD+). It was observed that 17 compounds were analyzed. It revealed the composition, chemical structure, molecular weight, and molecular formulation of different essential oil. This include 1,1-Dimethyl- chloro propanol ( $C_5H_{11}ClO$ , MolWgt,122Da), Alpha.-Hydroxyisocaproic acid ( $C_6H_{12}O_3$ , Mol W, Mol Wgt1,32Da), Dimethylsulfoxonium formyl methylide( $C_4H_8O_2S$ , Mol Wgt 120), Cyclohexasiloxane, dodecane thyl-(  $C_{12}H_{36}O_6Si_6$ ,Mol Wgt 444Da), Cycloheptasiloxane, tetradecamethyl-(  $C_{14}H_{42}O_7Si_7$ .Mol Wgt 518 Da), Heptasiloxane, hexadecamethyl-(  $C_{16}H_{48}O_6Si_7$ .Mol Wgt 518Da), Cyclo octasiloxane, hexadecane thyl-(  $C_{16}H_{48}O_8Si_8$ ,Mol Wgt 592Da), Cyclonona siloxane, octadeca ethyl-(  $C_{18}H_{54}O_9Si_9$ ,Mol Wgt 666Da), Heptasiloxane, hexadecamethyl-(  $C_{16}H_{48}O_6Si_7$ , Mol Wgt 532Da), Cyclononasiloxane, octadecane thyl-(  $C_{18}H_{54}O_9Si_9$ ,Mol Wgt 666Da), Heptasiloxane, hexadecamethyl-( $C_{16}H_{48}O_6Si_7$ , MolWgt532Da), Cyclodecasiloxane, eicosamethyl ( $C_{20}H_{60}O_{10}Si_{10}$ , Mol Wgt 7740Da), Dibutyl phthalate ( $C_{16}H_{22}O_4$ ,Mol Wgt 278 Da ),l-(+)-Ascorbic acid 2,6 dihexadecanoate( $C_{38}H_{68}O_8$  ,Mol Wgt 652Da), Heptasiloxane, hexadecamethyl-( $C_{16}H_{48}O_6Si_7$ , MolWgt532Da) and -Cyclooctasiloxane, hexadeca methyl-( $C_{18}H_{54}O_9 Si_9$  ,Mol Wgt 592Da).

**Table 3:** Gas chromatography and mass spectroscopy (GC-MS) analysis of *Aframomum melegueta* stem extract of purified Essential oil (StD++), this table denote the chemical nature of *Aframomum melegueta* stem extract of essential oil. It was observed that 48 compound were analyzed this include Dimethylsulfoxonium formylmethylide ( $C_4H_8O_2S$  , Mol Wgt 120Da), 2-Hexanol, 2-methyl-(  $C_7H_{16}O$  ,Mol Wgt 116 Da), Ethanol, 2-butoxy-(  $C_6H_{14}O_2$ , Mol Wgt, 118 Da), Cyclopentasiloxane, decamethyl ( $C_{10}H_{30}O_5Si_5$ ,Mol Wgt 370Da), Cyclohexa siloxane, dodeca methyl ( $C_{12}H_{36}O_6Si_6$ ,Mol Wgt 444Da),Hexasiloxane, tetradecamethyl-(  $C_{14}H_{42} O_5Si_6$ , Mol Wgt 458Da), Cyclohepta siloxane, tetradecamethyl-(  $C_{14}H_{42}O_7Si_7$  Mol Wgt 518Da), 1,4,7,-Cycloundecatriene, 1,5,9,9-tetramethyl,-Z,( $C_{15}H_{24}$  Mol Wgt 204 Da), Phenol, 2,4-bis(1,1-dimethylethyl)-(  $C_{14}H_{22}O$  Mol Wgt 206Da), Heptasiloxane, hexadecamethyl-(  $C_{16}H_{48} O_6 Si_7$  Mol Wgt 532 Da),Caryophyllene oxide( $C_{15}H_{24}O$  Mol Wgt 220Da), 2-Oxatricyclo [4.3.1.0 (3,8) ]decane( $C_9H_{14}OMolWgt138Da$ ),Methanol,[6,8,9-trimethyl-4-(1-propenyl)-3-oxabicyclo( $C_{15} H_{24}$

O<sub>2</sub> Mol Wgt 236 Da), Cyclooctasiloxane, hexadecamethyl(C<sub>16</sub>H<sub>48</sub>O<sub>8</sub>Si<sub>8</sub> Mol Wgt 592 Da), Isoaromadendrene epoxide(C<sub>15</sub>H<sub>24</sub>O mol Wgt Tricyclo[3.2.1.0<sup>2,7</sup>]oct-3-ene, 2,3,4,5-tetra methyl (C<sub>12</sub>H<sub>18</sub> Mol Wgt 162 Da), 1-Oxaspiro [2.5] octane,5,5-dimethyl-4-(3-methyl (C<sub>14</sub> H<sub>22</sub>O Mol Wgt 206Da), alpha.-Cadinol(C<sub>15</sub>H<sub>26</sub>O Mol Wgt 222Da),Heptasiloxane, hexa deca me th yl(C<sub>16</sub>H<sub>48</sub>O<sub>6</sub>Si<sub>7</sub> Mol Wgt 532Da), Longifolenaldehyde (C<sub>15</sub>H<sub>24</sub>O mol Wgt220 Da), Alloaro mad endrene oxide-(1)( C<sub>15</sub>H<sub>24</sub>O Mol Wgt 220 Da),Cyclononasiloxane, octadecamethyl-(C<sub>18</sub>H<sub>54</sub> O<sub>9</sub> Si<sub>9</sub>Mol Wgt 666 Da), 6-epi-shyobunol (C<sub>15</sub>H<sub>26</sub>O Mol Wgt 22Da), Tetradecanoic acid, ethyl ester (C<sub>16</sub>H<sub>32</sub>O<sub>2</sub> Mol Wgt 256Da), 4-Hexen-1-ol, 6-(2,6,6-trimethyl-1-cyclohexen(C<sub>16</sub>H<sub>28</sub>O Mol Wgt 236 Da),Heptasiloxane, hexadecamethyl(C<sub>16</sub>H<sub>48</sub>O<sub>6</sub>Si<sub>7</sub> Mol Wgt 532Da), 2-Penta decanone, 6,10,14-trimethyl-( C<sub>18</sub>H<sub>36</sub>O Mol Wgt 268Da), 1,2-Benzene dicarbox ylic acid, bis(2-methylpro (C<sub>16</sub>H<sub>22</sub>O<sub>4</sub> Mol Wgt 278 Da), Cyclodecasiloxane, eicosamethyl-(C<sub>20</sub>H<sub>60</sub>O<sub>10</sub>Si<sub>10</sub> Mol Wgt 740 Da), Hexa decanoic acid, methyl ester(C<sub>17</sub>H<sub>34</sub>O<sub>2</sub> Mol Wgt 270Da), 1,4-Methanoazulene-9-methanol, decahydro-4(C<sub>15</sub>H<sub>26</sub>O Mol Wgt 222Da), Dibutyl phthalate(C<sub>16</sub>H<sub>22</sub>O<sub>4</sub> Mol Wgt 278Da) ,Naphtho (2,3-b)furan-2(3H)-one,decahydro-8 (C<sub>15</sub> H<sub>20</sub> O<sub>2</sub> Mol Wgt 232Da), 1-(+)-Ascorbic acid 2,6-dihexa decanoate(C<sub>38</sub>H<sub>68</sub>O<sub>8</sub> mol Wgt 652 Da) , Menthol, 1'-(butyn-3-one-1-yl)-, (1R,2S,5 R)-( C<sub>14</sub>H<sub>22</sub>O<sub>2</sub> Mol Wgt 222Da), 4-Hexen-1-ol, 6-(2,6,6-trimethyl-1-cyclohexen(C<sub>16</sub>H<sub>28</sub>O, Mol Wgt 238Da) ,Heptasiloxane, hexadecamethyl-(C<sub>16</sub>H<sub>48</sub>O<sub>6</sub>Si<sub>7</sub> Mol Wgt 532Da), Hexadecanoic acid, ethyl ester(C<sub>18</sub>H<sub>36</sub>O<sub>2</sub> Mol Wgt 284Da), (3H)-Benzofuranone, 6-ethenylhexahydro-6-( C<sub>15</sub>H<sub>20</sub>O<sub>2</sub> Mol Wgt232Da),3,7,11,15-Tetramethyl hexadeca-1,6,10,14-tetr(C<sub>20</sub>H<sub>34</sub>O Mol Wgt 290Da), Cyclo octasiloxane, hexadecamethyl-(C<sub>16</sub>H<sub>48</sub>O<sub>8</sub>Si<sub>8</sub> Mol wgt 592 Da), 7a-Isopropenyl-4,5-dimethyl octa hydroinden-(C<sub>15</sub>H<sub>26</sub>O MolWgt 22Da),Linoleic acid ethyl ester C<sub>20</sub>H<sub>36</sub>O<sub>2</sub> Mol Wgt 308Da), Hepta siloxane, hexadecamethyl-(C<sub>16</sub>H<sub>48</sub>O<sub>6</sub>Si<sub>7</sub> Mol Wgt 532Da), Alloaromadendrene oxide-(1)( C<sub>15</sub>H<sub>24</sub>O Mol Wgt 220Da).

**Figure 1** denotes the Antibacterial activity of purified essential oil of *Aframomum melegueta*(StD++) stem extracts against selected clinical isolates (StD<sup>+</sup>). It was observed that all the selected test organisms were susceptible to the purified essential oil of *Aframomum melegueta* at varying degrees of concentration of 100, 50,25, and 12,5mg/ml. respectively. *E.coli* and *Staphylococcus aureus* has the highest zones of inhibition of 19,0 mm at 100mg/ml while *Klebsiella eplanticola* and *Citrobacter diversus* has the lowest zones of inhibition of 3.0mm at 12.5mg/ml. other relatively high zones of inhibition were *Corynebacterium cystitidis* (18.0mm), *Klebsiella pneumoniae* (17.0mm), *Proteus vulgaris* (17.0mm), *Citrobacter freundii* (15.0mm)and *Citro*

*bacter diversus* (13.0mm) and *Salmonella choleraesuis* (12.0mm). The lowest zones of inhibition were observed in *Staphylococcus aureus* and *E.coli*(7.0mm), *Corynebacterium cystitidis* (6.0mm), and *Klebsiella .pneumoniae* and *Citrobacter freundii* (5.0mm), *Proteus vulgaris*, *P.aeruginosa* and *Salmonella choleraesuis*(4.0mm) respectively.

**Figure 2,3,4,5** represents percentage frequency distribution of antibacterial activity against clinical isolates at (StD+).

**Figure 2** depicts the percentage frequency distribution of antibacterial activity against clinical isolates at (StD+) at 100mg/ml. At 100mg /ml, *Staphylococcus aureus* (12%) and *E.coli* (12%), have the highest while *Staphylococcus aureus* (6%), has the lowest percentage frequency distribution of purified essential oil of *Aframamum melegueta* (StD+) against selected clinical isolates. Other isolates were *Citrobacter diversus* (8%), *Proteus vulgaris* (10%) ,*Klebsiella .pneumoniae* (11%), *P.aeruginosa* (7%),*Staphylococcus aureus*( (6%), *Coryne bacterium cystitidis* (11%), *Klebsiella planticola* (7%), *Salmonella choleraesuis* (7%),*Citrobacter freundii* (9%) respectively.

**Figure 3** depicts the percentage frequency distribution of antibacterial activity against clinical isolates at (StD+) at 50mg/ml. At 50mg/ml, *E.coli*(13%) and *Staphylococcus aureus* (13%), have the highest while *Staphylococcus aureus* (6%), has the lowest percentage frequency distribution of purified essential oil of *Aframamum melegueta* (StD+) against selected clinical isolates. Other isolates were *Klebsiella .pneumoniae* (12%),*Proteus vulgaris* (10%), *P. aeruginosa* (7%),*Klebsiella planticola* (7%),, *Corynebacterium cystitidis* (10%),*Salmonella choleraesuis* (7%),, *Citrobacter freundii* (8%),*Citrobacter diversus* (7%) respectively.

**Figure 4** depicts the percentage frequency distribution of antibacterial activity against clinical isolates at (StD+) at 25mg/ml. At 25mg/ml, *E.coli* (12%) and *Staphylococcus aureus*

(12%), has the highest while *Proteus vulgaris* (7%), has the lowest percentage frequency distribution of purified essential oil of *Aframamum melegueta* (StD+) against selected clinical isolates. Other isolates were *Citrobacter freundii* (9%),*Citrobacter diversus* (9%), *Klebsiella .pneumoniae*(10%),*Proteus vulgaris* (7%),*P.aeruginosa* (9%),*Klebsiella planticola* (9%), *Staphylococcus aureus* (6%), *Coryne bacterium cystitidis* (9%), *Salmonella choleraesuis* (8%), *Citrobacter freundii*(9%),

**Figure 5** depicts the percentage frequency distribution of antibacterial activity against clinical isolates at (StD+) at 12.5mg/ml. At 12.5mg/ml, *E.coli* (13%), and *Staphylococcus aureus* (13%) have the highest while *Citrobacter diversus* (6%), has the lowest percentage frequency distribution of purified essential oil of *Aframamum melegueta* (StD++) against selected clinical isolates. Other isolates were *Klebsiella pneumoniae* (9%), *Proteus vulgaris* (8%), *P.aeruginosa* (8%), *Staphylococcus aureus* (9%), *Corynebacterium cystitidis* (11%), *Klebsiella planticola* (6%), *Citrobacter freundii* (9%), *Salmonella choleraesuis* (8%),

**Figure 6** depicts the antibacterial activity of purified essential oil of *Aframamum melegueta* (StD++) stem extracts against selected clinical isolates. *E.coli* has the highest zones of inhibition of 20mm at 100mg/ml, while *Klebsiella planticola* has the lowest inhibition of 1.0mm at 12.5mg/ml. the order of decreasing zones of inhibition as follows *Klebsiella pneumoniae*, *P.aeruginosa*, *P.aeruginosa*, and *Staphylococcus aureus* (17.0mm), *Corynebacterium cystitidis*, and *Citrobacter diversus* (14.0mm), *Citrobacter freundii* (13.0mm), *Salmonella choleraesuis* (12.0mm) and *Proteus mirabilis* (11.0mm) respectively. The decreasing order of zones of inhibition was as follows *Salmonella choleraesuis* (2.0mm), *Klebsiella pneumoniae* (3.0mm), *Klebsiella pneumoniae*, and *Corynebacterium cystitidis* (4.0mm), *Staphylococcus aureus*, *P. aeruginosa*, (5.0mm) and *E.coli* (6.0mm) respectively.

**Figure 7** depicts the percentage frequency distribution of antibacterial activity of purified essential oil of *Aframamum melegueta* (StD++) stem extracts against selected clinical isolates. At 100mg/ml, *E.coli* (12%) and *Staphylococcus aureus* (12%) has the highest while *Klebsiella planticola* (6%), has the lowest percentage frequency distribution of purified essential oil of *Aframamum melegueta* (StD++) against selected clinical isolates. Other isolates were *Citrobacter freundii* (8%), *Citrobacter diversus* (9%), and *Klebsiella pneumoniae* (10%), *Proteus mirabilis* (7%), *P.aeruginosa* (10%), *Staphylococcus aureus* (10%), *Corynebacterium cystitidis* (9%), *Salmonella choleraesuis* (7%) and *Citrobacter freundii* (8%) respectively .


**Figure 8** shows percentage frequency distribution of antibacterial activity of purified essential oil of *Aframamum melegueta* (StD++) stem extracts against selected clinical isolates at 50mg/ml. *E.coli* (15%) has the highest while *Citrobacter freundii* (5%) has lowest percentage

frequency distribution of purified essential oil of *Aframamum melegueta* (StD++) against selected clinical isolates, other isolates were *Citrobacter diversus* (7%), *Staphylococcus aureus* (12%), *Klebsiella.pneumoniae*(11%), *Proteus.mirabilis* (9%), *P.aeruginosa* (9%), *Staphylococcus aureus* (10%), *Corynebacterium cystitidis* (8%), *Klebsiella planticola* (6%), *Salmonella chole raesuis* (8%),

**Figure 9** shows the percentage frequency distribution of antibacterial activity of purified essential oil of *Aframamum melegueta* (StD++) stem extracts against selected clinical isolates at 25mg/ml. *Sta phylococcus aureus* (13%) has the highest while *Citrobacter freundii* (4%) has the lowest percentage frequency distribution of purified essential oil of *Aframamum melegueta* (StD++) against selected clinical isolates. Other isolates were *Klebsiella planticola*(6%), *Salmonella choleraesuis* (7%), *Citrobacter freundii* (4%), *Citrobacter diversus* (7%), *E.coli* (18%), *Staphylococcus aureus* (7%), *Klebsiella pneumoniae* (9%), *Proteus.mirabilis* (8%) , *P. aeruginosa* (10%) and *Coryneb acterium cystitidis*(11%).


**Figure 10** shows the percentage frequency distribution of antibacterial activity of purified essential oil of *Aframamum melegueta* (StD++) stem extracts against selected clinical isolates at 12.5mg/ml. *Staphylococcus aureus* (14%) has the highest while *Klebsiella planticola* (2%) has the lowest percentage frequency distribution of purified essential oil of *Aframamum melegueta* (StD++) against selected clinical isolates. Other isolates were *Citrobacter diversus* (5%), *E.coli* (12%), *Staphylococcus aureus* (12%), *Klebsiella. Pneumonia* (7%), *Proteus. mirabilis* (9%), *P.aeruginosa* (12%), *Corynebacterium cystitidis* (10%), *Klebsiella planticola* (2%) *Salmonella choleraesuis*(5%) and *Citrobacter freundii* (10%).

**Fig 1; Antibacterial Activity of Purified Essential Oil of Afr.melegueta(StD+) Stem Extracts Against Selected Clinical Isolates**




**Fig 1; Antibacterial Activity of Purified Essential Oil of Afr.melegueta(StD+) Stem Extracts Against Selected Clinical Isolates**


**Fig 2;Percentage Frequency Distribution of Antibacterial Activity of Afr. melegueta Stem against clinical isolates at (StD+) 100mg/ml**




**Fig 3;Percentage Frequency Distribution of Antibacterial Activity of Afr. melegueta Stem against clinical isolates at (StD+) 50mg/ml**




**Fig 4;Percentage Frequency Distribution of Antibacterial Activity of Afr. melegueta Stem against clinical isolates at (StD+) 25mg/ml**



**Fig 5;Percentage Frequency Distribution of Antibacterial Activity of Afr. melegueta Stem against clinical isolates at (StD+) 12.5mg/ml**




**Fig 6; Antibacterial Activity of Purified Essential Oil of Afr. melegueta.(SD++)Stem Extracts Against Selected Clinical Isolates**




**Fig 6; Antibacterial Activity of Purified Essential Oil of Afr. melegueta.(SD++)Stem Extracts Against Selected Clinical Isolates**


**Fig 7; Percentage Frequency Distribution of Antibacterial Activity of Afr. melegueta Stem against clinical isolates at (StD++) 100mg/ml**




**Fig 8; Percentage Frequency Distribution of Antibacterial Activity against of Afr. melegueta Stem clinical isolates at (StD++) 50mg/ml**



**Fig 9; Percentage Frequency Distribution of Antibacterial Activity of Afr. melegueta Stem against clinical isolates at (StD++) 25mg/ml**




**Fig 10; Percentage Frequency Distribution of Antibacterial Activity of Afr. melegueta Stem against clinical isolates at (StD++) 12.5mg/ml**










**Figure 11 :Spectral 1;Gas chromatography/mass spectroscopy(GC-MS) Spectra Analysis of *Afr. melegueta* Purified Essential oil stem extract (StD+)**

Chromatogram

















**Table 2; Gas chromatography/mass spectroscope(GC-MS) Analysis of *Afr. melegueta* stem extract of Purified Essential oil stem extract (StD+**

Retention time	Area	Area %	Height	Height%	A/H	Mol weight	Name	Structure formula	Structure
206	64426	0.25	53502	0.49	1.20	122	1,1-Dimethyl- chloropropanol	$C_5H_{11}ClO$	
4.248	356893	1.41	285635	2.61	1.25	132	alpha.-Hydroxyisocaproic acid	$C_6H_{12}O_3$	
4.577	13651447	53.84	4511552	41.18	3.03	120	Dimethylsulfoxonium formyl methylide	$C_4H_8O_2S$	
10.408	990350	3.91	812257	7.41	1.22	444	Cyclohexasiloxane, dodecamethyl-	$C_{12}H_{36}O_6Si_6$	
12.276	1517873	5.99	1209032	11.04	1.26	518	Cycloheptasiloxane, tetradecamethyl-	$C_{14}H_{42}O_7Si_7$	
13.060	173527	0.68	149045	1.36	1.16	518	Heptasiloxane, hexadecamethyl-	$C_{16}H_{48}O_6Si_7$	
13.939	1254825	4.95	1036069	9.46	1.21	592	Cyclooctasiloxane, hexadecamethyl-	$C_{16}H_{48}O_8Si_8$	







14.409	97955	0.39	43036	0.39	2.28	666	Cyclononasiloxane, octadecaethyl-	$C_{18}H_{54}O_9Si_9$	
14.578	437659	1.73	331899	3.03	1.32	532	Heptasiloxane, hexadecamethyl-	$C_{16}H_{48}O_6Si_7$	
15.531		4.85	740050	6.76	1.66	666	Cyclononasiloxane, octadecamethyl-	$C_{18}H_{54}O_9Si_9$	
16.382	493275	1.95	251641	2.30	1.96	532	Heptasiloxane, hexadecamethyl-	$C_{16}H_{48}O_6Si_7$	
17.692	1070365	4.22	414010	3.78	2.59	740	Cyclodecasiloxane, eicosamethyl-	$C_{20}H_{60}O_{10}Si_{10}$	
18.555	273288	1.08	98037	0.89	2.79	278	Dibutyl phthalate	$C_{16}H_{22}O_4$	
18.778	2352943	9.28	639712	5.84	3.68	652	l-(+)-Ascorbic acid 2,6 dihexadecanoate	$C_{38}H_{68}O_8$	
19.139	371477	1.47	131485	1.20	2.83	532	Heptasiloxane, hexadecamethyl-	$C_{16}H_{48}O_6Si_7$	
21.031	1019984	4.02	247666	2.26	4.12	592	Cyclooctasiloxane, hexadecamethyl-	$C_{18}H_{54}O_9Si_9$	

**figure 12 :Summary of Gas chromatography/mass spectrope(GC-MS) Analysis of *Afr. melegueta* stem extract of Purified Essential oil stem extract (StD+)**























**Table 3; Gas chromatography/mass spectroscope(GC-MS) Analysis of *Afr.melegueta* stem extract of Purified Essential oil stem extract (StD++)**







Peak	Retention time	Area	Area %	Height	Height %	A/H	Mol weight	Name	Structure formula	Structure
1	4.619	43145466	15.72	7641497	6.76	5.65	120	Dimethylsulfoxonium formylmethylyde	C <sub>4</sub> H <sub>8</sub> O <sub>2</sub> S	
2	4.674	2493029	0.91	2352888	2.08	1.06	116	2-Hexanol, 2-methyl-	C <sub>7</sub> H <sub>16</sub> O	
3	5.399	353165	0.13	255303	0.23	1.38	118	Ethanol, 2-butoxy-	C <sub>6</sub> H <sub>14</sub> O <sub>2</sub>	
4	8.333	837959	0.31	681373	0.60	1.23	370	Cyclopentasiloxane, decamethyl	C <sub>10</sub> H <sub>30</sub> O <sub>5</sub> Si <sub>5</sub>	
5	10.418	15200809	5.54	9650386	8.54	1.58	444	Cyclohexasiloxane, dodecamethyl	C <sub>12</sub> H <sub>36</sub> O <sub>6</sub> Si <sub>6</sub>	
6	11.369	434180	0.16	396118	0.35	1.10	458	Hexasiloxane, tetradecamethyl-	C <sub>14</sub> H <sub>42</sub> O <sub>5</sub> Si <sub>6</sub>	







7	12.289	19927221	7.26	11446136	10.13	1.74	518	Cycloheptasiloxane, tetradecamethyl-	$C_{14}H_{42}O_7Si_7$	
8	12.453	388191	0.14	234747	0.21	1.65	204	1,4,7,- Cycloundecatriene, 1,5,9,9-tetramethyl-, Z,	$C_{15}H_{24}$	
9	12.926	477280	0.17	199124	0.18	2.40	206	Phenol, 2,4-bis(1,1- dimethylethyl)-	$C_{14}H_{22}O$	
10	13.060	3804513	1.39	3075160	2.72	1.24	532	Heptasiloxane, hexadecamethyl-	$C_{16}H_{48}O_6Si_7$	
11	13.474	836300	0.30	253446	0.22	3.30	220	Caryophyllene oxide	$C_{15}H_{24}O$	
12	13.520	245557	0.09	163184	0.14	1.50	138	2-Oxatricyclo [4.3.1.0 (3,8)]decane	$C_9H_{14}O$	





13	13.696	962949	0.35	317368	0.28	3.03	236	Methanol, [6,8,9-trimethyl-4-(1-propenyl)-3-oxabicyclo	$C_{15}H_{24}O_2$	
14	13.951	18269598	6.66	10747963	9.51	1.70	592	Cyclooctasiloxane, hexadecamethyl	$C_{16}H_{48}O_8Si_8$	
15	14.040	540950	0.20	342696	0.30	1.58	220	Isoaromadendrene epoxide	$C_{15}H_{24}O$	
16	14.089	2073681	0.76	1061617	0.94	1.95	:220	12-Oxabicyclo [9.1.0] dodeca-3,7-diene, 1,5,5,8-	$C_{15}H_{24}O$	
17	14.244	1046953	0.38	286100	0.25	3.66	162	Tricyclo[3.2.1.0 <sup>2,7</sup> ]oct-3-ene, 2,3,4,5-tetramethyl	$C_{12}H_{18}$	
18	14.345	5991489	2.18	1361670	1.20	4.40	206	1-Oxaspiro [2.5] octane, 5,5-dimethyl-4-(3-methyl	$C_{14}H_{22}O$	

19	14.508	1342836	0.49	547372	0.48	2.45	222	.alpha.-Cadinol	$C_{15}H_{26}O$	
20	14.580	8855880	3.23	6434881	5.69	1.38	:532	Heptasiloxane, hexadecamethyl	$C_{16}H_{48}O_6Si_7$	
21	14.804	1309248	0.48	592766	0.52	2.21	220	Longifolenaldehyde	$C_{15}H_{24}O$	
22	14.864	574571	0.21	334118	0.30	1.72	220	Alloaromadendrene oxide-(1)	$C_{15}H_{24}O$	
23	15.547	20440142	7.45	9029655	7.99	2.26	666	Cyclononasiloxane, octadecamethyl-	$C_{18}H_{54}O_9Si_9$	
24	15.737	887291	0.32	256295	0.23	3.46	222	6-epi-shyobunol	$C_{15}H_{26}O$	


25	15.842	522807	0.19	165137	0.15	3.17	256	Tetradecanoic acid, ethyl ester	$C_{16}H_{32}O_2$	
26	16.328	10165724	3.70	4566247	4.04	2.23	236	4-Hexen-1-ol, 6-(2,6,6-trimethyl-1-cyclohexen	$C_{16}H_{28}O$	
27	16.386	10377796	3.78	5313878	4.70	1.95	532	Heptasiloxane, hexadecamethyl-	$C_{16}H_{48}O_6Si_7$	
28	16.542	3556482	1.30	1672036	1.48	2.13	268	2-Pentadecanone, 6,10,14-trimethyl-	$C_{18}H_{36}O$	
29	16.858	424896	0.15	189288	0.17	2.24	278	1,2-Benzene dicarboxylic acid, bis(2-methyl pro	$C_{16}H_{22}O_4$	
30	17.715	20188562	7.36	6714905	5.94	3.01	740	Cyclodecasiloxane, eicosamethyl-	$C_{20}H_{60}O_{10}Si_{10}$	


31	17.909	917069	0.33	285819	0.25	3.21	270	Hexadecanoic acid, methyl ester	$C_{17}H_{34}O_2$	
32	18.325	1483764	0.54	498584	0.44	2.98	222	1,4-Methanoazulene-9-methanol, decahydro-4	$C_{15}H_{26}O$	
33	18.571	3674794	1.34	1229358	1.09	2.99	278	Dibutyl phthalate	$C_{16}H_{22}O_4$	
34	18.705	2343459	0.85	706972	0.63	3.31	232	Naphtho(2,3-b)furan-2(3H)-one, decahydro-8	$C_{15}H_{20}O_2$	
35	18.760	1236194	0.45	452806	0.40	2.73	652	l-(+)-Ascorbic acid 2,6-dihexadecanoate	$:C_{38}H_{68}O_8$	
36	18.837	797999	0.29	227887	0.20	3.50	222	Menthol, 1'-(butyn-3-one-1-yl)-, (1R,2S,5R)-	$C_{14}H_{22}O_2$	

37	18.998	971942	0.35	174673	0.15	5.56	236	4-Hexen-1-ol, 6-(2,6,6-trimethyl-1-cyclohexen	$C_{16}H_{28}O$	
38	19.149	8785281	3.20	2696162	2.39	3.26	532	Heptasiloxane, hexadecamethyl-	$C_{16}H_{48}O_6Si_7$	
39	19.348	11148953	4.06	3221931	2.85	3.46	284	Hexadecanoic acid, ethyl ester	$C_{18}H_{36}O_2$	
40	19.428	4118606	1.50	1328485	1.18	3.10	232	2(3H)-Benzofuranone, 6-ethenylhexahydro-6-	$C_{15}H_{20}O_2$	
41	20.139	528888	0.19	154535	0.14	3.42	290	3,7,11,15-Tetramethylhexadeca-1,6,10,14-tetr	$C_{20}H_{34}O$	
42	21.051	20063075	7.31	4929341	4.36	4.07	592	Cyclooctasiloxane, hexadecamethyl-	$C_{16}H_{48}O_8Si_8$	


43	21.308	652829	0.24	149423	0.13	4.37	222	(7a-Isopropenyl-4,5-dimethyloctahydroindene)n-	$C_{15}H_{26}O$	
44	22.088	925406	0.34	202243	0.18	4.58	308	Linoleic acid ethyl ester	$C_{20}H_{36}O_2$	
45	23.769	5274359	1.92	1025615	0.91	5.14	532	Heptasiloxane, hexadecamethyl-	$C_{16}H_{48}O_6Si_7$	
46	24.064	1051570	0.38	215439	0.19	4.88	220	Alloaromadendrene oxide-(1)	$C_{15}H_{24}O$	

**Figure 14: Summary of Gas chromatography/mass spectroscope(GC-MS) Analysis of *Afr. melegueta* stem extract of Purified Essential oil stem extract (StD++)**







Heptasiloxane,5,5-dimethyl-4-(3-methyl)




Longifolenaldehyde




Alloaromadendreneoxide-(1)



Cyclononasiloxaneoctadecamethyl-



6-epi-shyobunol




Cyclodecasiloxaneicosamethyl-




Hexadecanoic acid, methyl ester dihexadecanoate



1,4-Methanoazulene-9-methanol, decahydro-4




Dibutyl phthalate




Naphtho(2,3-b)furan-2(3H)-one, decahydro-8



Tetradecanoic acid ethyl ester trimethyl-1-cyclohexen




4-Hexen-1-ol, 6-(2,6,6-trimethyl-1-cyclohexen)



Heptasiloxane, hexadecamethyl-



2-Pentadecanone 6.1



1,2-Benzene dicarboxylic acid, bis(2-methyl)



#### 4.0 Discussion

This research work aims to ascertain the composition and potency of the bioactive component of essential oil derived from *Aframomum melegueta* stem extract through multi-phase solvent extraction Gas Chromatography /Mass Spectroscopic analysis and antimicrobial assay against selected clinical isolates. In this present study, it was observed that *Aframomum melegueta* stem extract has overwhelming potential uses in our-to-day human activity especially the antibacterial potency on clinical isolate. It can be observed the *Aframomum melegueta* stem extract has various degrees of antibacterial potency on clinical isolates which is corroborated with different results gathered in the table and figures

In StD+ Fraction,(Antibacterial activity of purified essential oil of *Aframomum melegueta* (StD+) stem extracts against selected clinical isolates), *E.coli* and *Staphylococcus aureus* has the highest zones of inhibition of 19,0 mm at 100mg/ml while *Klebsiella eplanticola* and *Citrobacter diversus* has the lowest zones of inhibition of 3.0mm at 12.5mg/ml. other relatively high zones of inhibition were *Corynebacterium cystitidis* (18.0mm), *Klebsiella .pneumoniae* (17.0mm), *Proteus vulgaris* (17.0mm), *Citrobacter freundii* (15.0mm) and *Citrobacter diversus*(13.0mm) and *Salmonella choleraesuis* (12.0mm). The lowest zones of inhibition were observed in *Staphylococcus aureus* and *E.coli* (7.0mm), *Corynebacterium cystitidis*(6.0mm), *Klebsiella .pneumoniae* and *Citrobacter freundii* (5.0mm), *Proteus vulgaris*, *P.aeruginosa* and *Salmonella choleraesuis* (4.0mm) while In StD++, fraction,(Antibacterial activity of purified essential oil of *Aframomum melegueta*(StD++) stem extracts against selected clinical isolates),*E.coli* has the highest zones of inhibition of 20mm at 100mg/ml, while *Klebsiella planticola* has the lowest of inhibition of 1.0mm at 12.5mg/ml. the order of decreasing zones of inhibition as follows *Klebsiella.pneumoniae*, *P.aeruginosa* *P.aeruginosa* and *Staphylococcus aureus* (17.0mm), *Corynebacterium cystitidis* and *Citrobacter diversus* (14.0mm),*Citrobacter freundii* (13.0mm), *Salmonella choleraesuis*(12.0mm) and *Proteus. mirabilis* (11.0mm) respectively. The decreasing order of zones of inhibition was as follows *Salmonella choleraesuis* (2.0mm), *Klebsiella.pneumoniae*(3.0mm),*Klebsiella. Pneumoniae*, and *Corynebacterium cystitidis* (4.0 mm), *Staphylococcus aureus*, *P.aeruginosa*, (5.0mm) and *E.coli*,(6.0mm). The result shows that *Aframomum melegueta* stem extract possesses some levels of antibacterial potential and is in agreement with previous

findings on oils of *V. danielli* and *Aframomum melegueta* [26,27] and *A. melegueta* [28,29,20].

Some researchers propound that the presence of secondary metabolite i.e Phyto-constituents is the defector to the mechanism of action of essential oil from *Aframomum melegueta* stem extract, like Flavonoids are ketone-containing secondary metabolites found in plants. Flavonoids such as myricetin, quercetin, and morin have been extensively investigated and proven to possess anti-inflammatory, antimicrobial, and antiproliferative activities [30]

Alkaloids are bioactive molecules with nitrogen, constituting a key component of their molecular architecture, with distinct antimicrobial properties example are furoquinolones, indole alkaloids and acridones which is recent, with a plethora of studies reporting the efficacy of alkaloids against protozoan parasites [31]. The aglycone and hydrophilic sugar moieties are active biosurfactants that have several therapeutic applications including immunostimulatory, molluscicidal, hypocholesterolemic, antimicrobial, and antioxidant activities [32]. Cardiac glycoside is a cardiotonic steroid, that generally contains a steroid-like structure, and induces a cardiotonic effect via selective inhibition of  $\text{Na}^+/\text{K}^+$ -ATPase [33]. Tannins are moieties linked to a carbohydrate core (hydrolyzable tannins), and have wide therapeutic properties including antiproliferative and antimicrobial activities [34],

However, apart from the secondary metabolite which is responsible for the metabolic activity of the essential oil of *Aframomum melegueta* stem extract, the Gas chromatographic chemical constituent also reveals the presence of arrays of chemical architecture ranging from alkyl group to the alkanooates. The prominent ones are from Std+ fraction are Cycloheptasiloxane, tetradeca methyl( $\text{C}_{14}\text{H}_{42}\text{O}_7\text{Si}_7$ ), l-(+)-Ascorbic acid 2,6-dihexadecanoate( $\text{C}_{38}\text{H}_{68}\text{O}_8$ ) and Dodecanoic acid ( $\text{C}_{12}\text{H}_{36}\text{O}_6\text{Si}_6$ ) [34].

The first highest peak from (16) sixteen components in GCMS analysis of essential oil fraction (SD+) of *Aframomum melegueta* stem extract was Cycloheptasiloxane, tetradecamethyl. Cycloheptasiloxane, tetradecamethyl-(( $\text{C}_{14}\text{H}_{42}\text{O}_7\text{Si}_7$ ) compound which belongs to the class of organic compounds known as organo-heterosilanes. These are organosilicon compounds where the tetravalent silicon atom is linked to one or more heteroatoms. Cyclotrisiloxane (CAS No. 541-05-9) is the cyclic dimethyl polysiloxane that conforms to the generic structure of cyclic dimethyl polysiloxane compounds, where  $n = 3$ , and the other components of Cyclomethicone

(where  $n = 4, 5, 6,$  or  $7$ ) are present at the levels of less than 1%. Other names for cyclo-trisiloxane include cyclo-trisiloxane, hexamethyl- and hexamethylcyclotrisiloxane [35].

The presence of Cyclohexasiloxane, dodecamethyl in *Aframomum melegueta* stem extract is used in personal care products, antiperspirants and antifungals alluded to the report of (36) that the plant extract of *Aframomum melegueta* is an effective antifungal agent in the treatment of fungi infection and a promising alternative/adjunct/ supplement to the azole and allylamine group. Cyclohexasiloxane, dodecamethyl in *Aframomum melegueta* stem extract may be used as personal care compositions and cosmetic compositions can be in the form of a solution, emulsion, foam, mousse, cream, gel, lotion, ointment, solid, powder, paste, semi-solid, stick, spray or a combination thereof. Exemplary personal care compositions or cosmetic compositions include deodorants, antiperspirants, insect repellants, anesthetics, skin conditioners, skin lotions, skin moisturizers, skin toners, skin sanitizers, skin cleansing compositions, skin soothing and lubricating compositions, sunscreen, anti-aging products, concealer products, soaps, foaming bath products, shower gels, cleansing products, shampoos, hair conditioners, hair styling gels, hair anti-dandruff compositions, hair growth promoter compositions, hair colorant compositions, hair bleaching agent compositions, hair anti-frizzing agent compositions, hair shining compositions, hair relaxer compositions, mousses, styling gels, hair sprays, hair dyes, hair waving products, hair straightening products, shaving product compositions, personal lubricant compositions, spermicidal gel compositions, manicure products, nail polish, nail polish remover, nail creams and lotions, cuticle softeners, color cosmetics, lipsticks, lip balms, foundations, face powders, eye liners, eye shadows, blushes, makeup, mascaras and color cosmetic removers [37], this shows the various ways in which Cyclohexasiloxane, dodecamethyl in *Aframomum melegueta* stem extract can be put to biological use.

The second highest peak from (16) sixteen components in GC/MS analysis of essential oil fraction (SD+) of *Aframomum melegueta* stem extract was l-(+)-Ascorbic acid 2,6-dihexadecanoate ( $C_{38}H_{68}O_8$ ) It is a vitamin C compound. Ascorbic acid is often used for preventing and treating the common cold, gum disease, acne, and other skin infections, bronchitis, stomach ulcers, tuberculosis, dysentery, boils, and wounds [38], “Ascorbic acid is also used to prevent glaucoma, cataracts, gallbladder disease, dental cavities, constipation, hay fever, asthma,

arthritis, back pain, diabetes, chronic fatigue syndrome, osteoporosis and boosting the immune system” (38), “Ascorbic acid acts as an antioxidant in the skin by scavenging and quenching free radicals generated by ultraviolet radiation. The use of *Aframomum melegueta* stem in the treatment of diarrhoea, dysentery, stomach problems, ulcers, wound, fever and other form of health challenges in herbal medicine could be as a result of the presence of ascorbic acid 2,6-dihexadecanoate in the stem of the plant” [39,40]

“The combination of Cyclohexasiloxane and Dodecanoic acid also known as lauric acid has been a driving factor in the antimicrobial properties of *Aframomum melegueta* stem extract”[41]. “Tetradecanoic acid known as myristic acid is commonly added co-translationally to the penultimate, nitrogen-terminus, glycine in receptor-associated kinases to confer the membrane localization of the enzyme”[42].

“The first and second highest peak from (48) sixteen components in GCMS analysis of essential oil fraction (SD++) of *Afr. melegueta* stem extract were Phenol, 2,4-bis(1,1-dimethyl ethyl ( $C_{14}H_{22}O$ ) and Cyclooctasiloxane, hexadecamethyl-( $C_{16}H_{48}O_8Si_8$ ), Cyclooctasiloxane, hexadecamethyl-( $C_{16}H_{48}O_8Si_8$ ) has been discussed previously. Moreover, 2,4-Di-tert-butyl-phenol or 2,4-bis(1,1-dimethylethyl)-phenol (2,4-DTBP) is a common natural product that exhibits potent toxicity against almost all testing organisms, including the producing species. 2,4-DTBP can modulate the secreted EPS of *Serratia marcescens*, which in turn could facilitate the disruption of biofilms, as well as favor the diffusion of antimicrobials into the cell aggregates, resulting in the eradication of persistent and biofilms”[44].

“This compound can be used to enhance the efficacy of conventional antibiotics. Intercellular communication in bacteria (quorum sensing (QS)) is an important phenomenon in disease dissemination and pathogenesis that controls biofilm formation. 2,4-DTBP controls QS-mediated biofilm formation and simultaneously increases the hydration of the cell wall, which results in reduced biofilm formation” (43) and “3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium (MTT) and plaque reduction assays showed that 2,4-DTBP exhibited significant anti-coxsackievirus B-3 (CVB-3) and anti-herpes virus type 2 (HSV-2) activities”[45].

“The compound was found to be effective against an agriculturally important root-rot fungus *Fusarium oxysporum* by inhibiting spore germination and hyphal growth” (46,47,48). During the fungal spore germination, 2,4-DTBP completely inhibited the germination by

preventing the emergence of a normal germ tube and led to the abnormal branching and swelling of hyphae.

## **Conclusion :**

The study shows the richness of naturally occurring essential oil of *Afr. melegueta* which is deposited abundantly in the fresh stem extract and suggests its antimicrobial capability and potentials as a natural antimicrobial agent against selected multiple resistant microorganisms. *Afr. melegueta* essential oil has limitless potentials and its various uses, example of such uses was elucidated during the course of this research work. Our observation is mind blowing. Fresh stem extract of *Afra.melegueta* also contain many useful naturally occurring chemical compounds which are useful in nature. *Aframomum melegueta* extracts is indeed a very rich medicinal plant

## **FUNDING**

This work was supported by TETFUND Institution Based Research program, Nigeria.

## **ACKNOWLEDGEMENTS**

The authors wish to express their appreciation to TETFUND Institution Based Research program, Nigeria for sponsoring this research work and all the technical staff of the laboratory unit of the Department of Microbiology, Faculty of Science, Adekunle Ajasin University, Akungba-Akoko, Ondo State, Nigeria, Obafemi Awolowo University, Ile Ife, Osun State, Nigeria, for their technical support.

## **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

Corresponding author; Oludare temitope Osuntokun ; **ORCID-<https://orcid.org/0000-0002-39546778>, Web of Science Researcher ID; L-4314-2016**

[linkedin.com/in/dr-oludare-temitope-osuntokun-87b56477](https://www.linkedin.com/in/dr-oludare-temitope-osuntokun-87b56477)

## References

1. Omukoro S.; Ashorobi R. B(2007). Further studies on the antinociceptive action of aqueous seed extract of *Aframomum melegueta*. J. Ethnopharmacol. 2007, 109, 501–504

2. Osuntokun, O.T., Balogun, J.O., Akele, E.O., Adedayo, S.A., & Bello, O. A. (2021). *Aframomum melegueta* [Roscoe] K. Schum) Ethanolic Seed Extract against Asymptomatic Bacteriuria Isolates from Pregnant Women Attending Antenatal Clinic in Ondo State, Nigeria. *International Journal of Pathogen Research*, 8(2), 12-43.
3. Galal A. M(1996). Anti-microbial activity of 6-paradol and related compounds. *Int. J. Pharmacog.* 1996, 31, 64–69.
4. Kamtchouing P.; Mbongue G. Y. F.; Dimo T.; Watcho P.; Jatsa H. B.; Sokeng S. D. ( 2002) Effects of *Aframomum melegueta* and *Piper guineense* on sexual behaviour of male rats. *Behav. Pharmacol.* 2002, 13, 243–247.
5. Alo M. N.; Anyim C.; Igwe J. C.; Elom M.; Uchenna D. S.( 2012) Antibacterial activity of water, ethanol and methanol extracts of *Ocimum gratissimum*, *Vernonia amygdalina* and *Aframomum melegueta*. *Adv. Appl. Sci. Res.* 2012, 3, 844–848.
6. Chiejina N. V.; Ukeh J. A(2012). Antimicrobial properties and phytochemical analysis of methanolic extracts of *Aframomum melegueta* and *Zingiber officinale* on fungal diseases of tomato fruit. *J. Nat. Sci. Res.* 2012, 2, 10–15.
7. Mbongue G. Y. F.; Kamtchouing P.; Dimo T(2012,). Effects of the aqueous extract of dry seeds of *Aframomum melegueta* on some parameters of the reproductive function of mature male rats. *Andrologia* 2012, 44, 53–58
8. Umukoro S.; Ashorobi R. B(2001). Effect of *Aframomum melegueta* seed extract on thermal pain and on carrageenin-induced edema. *Nig. Q. J. Hosp. Med.* 2001, 11, 220–225.
9. Umukoro S.; Ashorobi R. B(2008). Further pharmacological studies on aqueous seed extract of *Aframomum melegueta* in rats. *J. Ethnopharmacol.* 2008, 115, 489–493
10. Lawal B. A. S.; Aderibigbe A. O.; Essiet G. A.; Essien A. D(2007). Hypotensive and antihypertensive effects of *Aframomum melegueta* seeds in humans. *Int. J. Pharmacol.* 2007, 3, 311–318.
11. Sugita J.; Yoneshiro T.; Hatano T.; Sayuri A.; Takeshi I.; Hideyo U.; Toshihiko I.; Toshimitsu K.; Yuko K.; Masayuki S( 2013). Grains of paradise (*Aframomum melegueta*) extract activates brown adipose tissue and increases whole-body energy expenditure in men. *Br. J. Nutr.* 2013, 110, 733–738.
12. Sugita J.; Yoneshiro T.; Sugishima Y.; Ikemoto T.; Uchiwa H.; Suzuki I.; Masayuki S. Daily ingestion of grains of paradise (*Aframomum melegueta*) extract increases whole-body

energy expenditure and decreases visceral fat in humans. *J. Nutr. Sci. Vitaminol.* 2014, 60, 22–27.

13. Ajaiyeoba EO, Ekundayo O(1999). Essential oil constituents of *Aframomum melegueta* (Roscoe) K. Schum. seeds (alligator pepper) from Nigeria. *Flavour and Fragrance Journal.* 1999; 14(2):109-111.

14. Owokotomo IA, Ekundayo O, Oguntuase BJ(2014). Chemical constituents of the leaf, stem, root and seed essential oils of *Aframomum melegueta* (K. Schum) from South West Nigeria. *International Research Journal of Pure and Applied Chemistry.* 2014; 4(4):395-401

15. Ntonifor NN, Mueller-Harvey I, van Emden HF, Brown RH(2006). Antifeedant activities of crude seed extracts of tropical African species against *Spodoptera littoralis* (Lepidoptera: Noctuidae). *International Journal of Tropical Insect Science* 2006; 26(2):78-85

16. Menut C, Lamaty G, Amvam Zollo PH, Atogho BM, Abondo R, Bessièrè JM(1991). Aromatic plants of tropical central Africa. V. volatile components of three Zingiberaceae from Cameroon: *Aframomum melegueta* (roscoe) K. Schum., *A. daniellii* (hook. f.) K. Schum. and *A. sulcatum* (oliv. and hanb.) K. Schum. *Flavour and Fragrance Journal* 1991; 6(3):183-186

17. Lamaty G, Menut C, Koudou J, Régnier P, Bessièrè JM(1993). Aromatic plants of tropical central Africa. XI. Essential oils of leaf and seed of *Aframomum melegueta* (Roscoe) K. Schum. from Central African Republic, *Journal of Essential Oil Research.* 1993; 5(1):81-83

18. Adefegha SA, Olasehinde TA, Oboh G(2017). Essential oil composition, antioxidant, antidiabetic and antihypertensive properties of two *Aframomum* species. *Journal of Oleo Science.* 2017; 66(1):51-63

19. Ukeh DA(2008). Bioactivities of essential oils of *Aframomum melegueta* and *Zingiber officinale* both (Zingiberaceae) against *Rhyzopertha dominica* (Fabricius). *Journal of Entomology.* 2008; 5:193-199

20. Uzeh RE, Oguntosin DO(2013),. Efficacy of essential oils from some African spices against two strains of *Bacillus cereus* isolated from vegetable salad. *Journal of Food Research.* 2013; 2(6):48-54.

21. Héritier VNV, Paulin MK, Nadège NK, Michel MP, Jude-Thaddée MN(2017). *In vitro* antifungal activity of essential oils extracted from some plants of *Tangawisi*® products on *Candida albicans*. *Journal of Pharmacognosy and Phytochemistry.* 2017; 6(3):01-05

22. Owokotomo IA, Ekundayo O, Abayomi TG, Chukwuka AV(2015). *In-vitro* anti-cholinesterase activity of essential oil from four tropical medicinal plants. *Toxicology Reports*. 2015; 2:850-857
23. Oludare Temitope Osuntokun,V. O. Olumekun, A. O. Ajayi, I. O. Omotuyi and A. Olonisakin (2020a)Assessment of in-vitro Antioxidant/Enzymes Inhibitory Potentials of *Aframomum melegueta* [Roscoe] K.Schum(Grains of Paradise)Leaf, Stem Bark, Seed Bark and Seed Extracts, *Archives of Current Research International* 20(2): 40-57
24. Olaposi Omotuyi,Oyekanmi Nash, Basiru O. Ajiboye,Victor O. Olumekun, Babatunji E., Oyinloye, Oludare Temitope. Osuntokun,Adebisi Olonisakin,A. Olajide Ajayi, Olasehinde Olusanya,Funmilola S. Akomolafe,Niyi Adelokun (2020) *Aframomum melegueta* secondary metabolites exhibit polypharmacology against SARS-CoV-2 drug targets: in vitro validation of furin inhibition, *Phytotherapy Research*. 2020;1–12. Wiley onlinelibrary .com /journal /ptr, DOI: 10.100 2/ptr.6843,Pp,1-12,
25. Osuntokun, O. T., Bakare, O. S., Bankole, O. M., & Ayodele, O. A. (2020b). Comparative Study of *Ocimum gratissimum* Mediated Nanoparticles and Conventional Antibiotics against Endophytic Leguminous Bacteria and Selected Clinical Isolates. *Asian Journal of Advanced Research and Reports*, 13(1), 42-59.
26. Njimoh DL, Assob JCN, Mokake SE, Nyhalah DJ, Yinda CK, Sandjon B. Antimicrobial activities of a plethora of medicinal plant extracts and hydrolates against human pathogens and their potential to reverse antibiotic resistance. *International Journal of Microbiology*, Volume 2015 Article ID 547156, 15 pages. 2015;
27. Martins AP, Salgueiro L, Gonçalves MJ, da Cunha AP, Vila R, Cañigüeral S *et al.* Essential oil composition and antimicrobial activity of three Zingiberaceae from S.Tomé e Príncipe. *Planta Medica*. 2001; 67(6):580-584
28. Uzeh Lawal OA, Kasali AA, Opoku RA, Ojekale AB, Oladimeji OS Bakare S(2015). Chemical composition and antibacterial activity of essential oil from the leaves of *Aframomum melegueta* (Roscoe) K. Schum from Nigeria. *Journal of Essential Oil-Bearing Plants*. 2015; 18(1):222-229
29. Goc A, Niedzwiecki A, Rath M(2015). In vitro evaluation of antibacterial activity of phytochemicals and micronutrients against *Borrelia burgdorferi* and *Borrelia garinii*. *J Appl Microbiol*. 2015;119(6):1561–72.

30. Semwal DK, Semwal RB, Combrinck S, Viljoen A. Myricetin(2016): A dietary molecule with diverse biological activities. *Nutrients* [Internet]. 2016 Feb 16 [cited 2017 Sep 13];8(2). Available:<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4772053>
31. Amoa Onguéné P, Ntie-Kang F, Lifongo LL, Ndom JC, Sippl W, Mbaze LM(2013). The potential of anti-malarial compounds derived from African medicinal plants, part I: a pharmacological evaluation of alkaloids and terpenoids. *Malar J.* 2013;12:449.
32. Francis G, Kerem Z, Makkar HPS, Becker K.( 2002) The biological action of saponins in animal systems: a review. *Br J Nutr.*;88(6):587–605.
33. Moses T, Papadopoulou KK, Osbourn A(2014);.Metabolic and functional diversity of saponins, biosynthetic intermediates and semi-synthetic derivatives. *Crit Rev Biochem Mol Biol.* ;49(6):439–62.
34. Yusuf-Babatunde A.M, Osuntokun, O.T.,Ige, O.O. Solaja O.(2019) Evaluation of Antimicrobial, Antioxidant and Gas Chromatography mass Spectrometry Profile of the Stem Bark Extract of *Bombax buonopozense* P. Beauv. (Bombacaceae) *FUOYE Journal of Pure and Applied Sciences, FJPAS Vol 4(1) ISSN: 2816-1419,*
35. Patel JR ; P Tripathi ; V Sharma ; NS Chauhan ; VK Dixit(2011). *Journal of Ethno pharmacology*, 138(2),286-313.
36. Kumar A, Kennedy-Boone D, Weisz HA, Capra BA, Uchida T, Jennings K(2015). Neuroprotective effects of *Aframomum melegueta* extract after experimental traumatic brain injury. *Natural Product Chemistry and Research.*; 3:167. doi: 10.4172/2329-6836.1000167
37. Thijssen MA; RP Mensink (2005). *Fatty Acids and Atherosclerotic Risk*. In Arnold Von Eckardstein (Ed).*Atherosclerosis :Diet and Drugs*. Springer, 171-172.
38. Olumekun, V. O., Osuntokun, O. T., Ajayi, A. O., Omotuyi, I. O., & Olonisakin, A. (2020). Estimation of Vitamins in *Aframomum melegueta* [Roscoe] K. Schum and Its Relevance as Natural Immune Booster. *Journal of Applied Life Sciences International*, 23(7), 25-39.
39. Osuntokun O. T., Peace O. Ukwuoma,. Yusuf-Babatunde A. M and Ige O. O(2017), Evaluation of Antimicrobial Activity, Phytochemical Screening and Gas Chromatography-Mass Spectrophotometric Profile of Essential Oil from *Persea Americana*&*Citrus sinensis*, *Asian Journal of Medicine and Health*, Volume 5(1): 2017, Pp 1-17.
40. Osuntokun O.T, Ayegbusi B.L,Yusuf-Babatunde A.M, Ige O.O.(2017)Assessment of Antimicrobial, Phytochemical Screening and Gas Chromatography-Mass Spectro photometric

Profile of Crude *Chrysophyllum albidum* Essential Oil, *Chemistry Research Journal*, 2017, Volume 2(5):68-85,

41. Ntonifor NN, Mueller-Harvey I, van Emden HF, Brown RH (2006). Antifeedant activities of crude seed extracts of tropical African species against *Spodoptera littoralis* (Lepidoptera: Noctuidae). *International Journal of Tropical Insect Science*; 26(2):78-85.

42. Osuntokun O.T, Idowu T.O, Jatto F.O. (2019) "In Vitro Analysis, Secondary Metabolite Screening and GC-MS Profile of *Ricinodendron Heudelotii* (Muh.Arg) Essential Oil Extracts Against Selected Multiple Drug Resistance Clinical Isolates" *International Journal of Research and Innovation in Applied Science -IJRIAS* 4(6) pp.07-25,

43. Padmavathi, A.R.; Periyasamy, M.; Pandian, S.K (2015). Assessment of 2,4-di-tert-butyl phenol induced modifications in extracellular polymeric substances of *Serratia marcescens*. *Bioresour. Technol.* 2015, 188, 185–189.

44. Padmavathi, A.R.; Abinaya, B.; Pandian, S.K (2014). Phenol, 2,4-bis(1,1-dimethylethyl) of marine bacterial origin inhibits quorum sensing mediated biofilm formation in the uropathogen *Serratia marcescens*. *Biofouling*, 30, 1111–1122

45. Leila, A.; Lamjed, B.; Roudaina, B.; Najla, T.; Taamalli, A.; Jellouli, S.; Mokhtar, Z (2019). Isolation of an antiviral compound from Tunisian olive twig cultivars. *Microb. Pathog.*, 128, 245–249

46. Dharni, S.; Maurya, A.; Samad, A.; Srivastava, S.K.; Sharma, A.; Patra, D.D (2014). Purification, characterization, and in vitro activity of 2,4-Di-tert-butylphenol from *Pseudomonas monteilii* PsF84: Conformational and molecular docking studies. *J. Agric. Food Chem.*, 62, 6138–6146.

47. Chung KT, Wong TY, Wei CI, Huang YW, Lin Y (1998). Tannins and human health: a review. *Crit Rev Food Sci Nutr*; 38(6):421–64.

48. Osuntokun, O. T., Bakare, O. S., Bankole, O. M., & Ayodele, O. A. (2020). Comparative Study of *Ocimum gratissimum* Mediated Nanoparticles and Conventional Antibiotics against Endophytic Leguminous Bacteria and Selected Clinical Isolates. *Asian Journal of Advanced Research and Reports*, 13(1), 42-59.