

1                   **SPATIAL VARIABILITY MAPPING OF SOIL CHEMICAL**  
2                   **PROPERTIES USING GIS & GPS**

3  
4   **ABSTRACT**

5 Excessive application of fertilizers can cause wastage of fertilizer which increases input cost  
6 and environmental pollution. Implementation of Precision Agriculture through site specific  
7 nutrient management is the best suitable solution to increase nutrient application efficiency  
8 and thereby increase crop productivity. The methodology follows the delineation of the study  
9 area, location of sampling points and soil samples were collected and analysed for the soil  
10 chemical properties such as pH, Electric Conductivity, Available Nitrogen, Available  
11 Phosphorous, Available Potassium, Boron and Sulphur by using standard methods. Spatial  
12 variability maps of soil chemical properties were prepared by using Inverse Distance  
13 Weighing method of interpolation in ArcGIS. From this study, it could be concluded that,  
14 GIS along with GPS could be used as an effective tool for preparation of spatial variability  
15 maps for determining the spatial distribution of chemical properties of soils and thereby helps  
16 to achieve site specific nutrient recommendation which improves crop productivity, quality  
17 and reduce environmental stress. It also involves lesser number of soil analysis and thus  
18 reduces the cost of operation compared to plot-to-plot analysis.

19 **Key words:** Geographic Information of system (GIS), Global Positioning System (GPS), Site  
20 specific nutrient management (SSNM), Inverse distant weighing method (IDW).


21 **Introduction:**

22 The population has been augmenting everywhere in the planet which leads to the need of  
23 accelerating food production. Use of various fertilizers has significant contribution in  
24 increasing food production in order to reduce the world food insecurity. Recent studies  
25 showed that nutrient inputs are responsible for 30–50% of the crop yield. However excessive  
26 application of fertilizers can cause wastage of fertilizer which increases input cost and  
27 environmental pollution. In order to solve all these problems, implementation of Precision  
28 Agriculture through site specific nutrient management is the best suitable solution to increase  
29 nutrient application efficiency and thereby increase crop productivity. Site Specific Nutrient  
30 Management (SSNM) is the real time feeding of crops with nutrients while recognizing the  
31 spatial variability within the fields.

32 Many technologies such as remote sensing, variable rate technologies, nano  
33 technology etc., are used for the implementation of site-specific nutrient management but GIS  
34 found to be the most promising tool due to its vast applications. GIS techniques in Precision  
35 Agriculture are used for variety of applications such as conservation of important plant  
36 species in land use planning, land use suitability evaluations, crop selections and rotations,  
37 irrigation and mechanisation planning. Spatial analysis is the most important component of  
38 SSNM which is determined through Geographical Information System (GIS). Agricultural  
39 management interacts with environmental parameters and natural resources that have a clear  
40 spatial character and hence, GIS play a critical role in agricultural productivity, notably in the  
41 application of fertilisers. In most research, GIS is used to process model inputs and to display  
42 outcomes, but they may also be used for other purposes. GIS plays a key role in unravelling  
43 more complex and specialised problems, such as fertiliser management difficulties.

#### 44 **Materials and Methods:**

45 Fertility status of the soil is an important factor for achieving sustainable agricultural  
46 production, which was declining day by day results in declining yields and environmental  
47 pollution. Fertilizer management varies with zones, and it has a significant impact on  
48 agricultural output and quality. (Jeya and Vasanthakumar, 2020). The spatial variability of  
49 soil properties is needed for agricultural productivity, food safety and environmental  
50 modelling (Bhunias,2018). Understanding of soil nutrient distributions and the factors  
51 affecting them are crucial for fertilizer management and environmental protection in  
52 vulnerable ecological regions (Gao,2019). Preparation of fertility maps with the help of both  
53 GPS and GIS by locating sampling points, thereby collecting and analysing of samples were  
54 done by using standard methods. Recommendations were given to farmers based on the  
55 spatial distribution maps. The area was delineated by using cadastral map of the study area  
56 and coordinates of the corner of the study area were found with the help of hand held GPS.  
57 Georeferencing of the map was done by using the georeferencing tool of Arc GIS 10.3. Shape  
58 file of the study area was prepared along with the features such as buildings, placemarks,  
59 road and river as shown in fig.1.



60

61

**Fig. 1**Shape file of the KCAET campus

62

**Soil sampling points by using gridding**

63

Gridding was done in order to locate the sampling points, by using gridding tool in Arc GIS.

64

A grid interval of 100 ×100 m was taken for the study. The grid map was then exported to

65

google earth which is easier for visual identification of sampling points. It can be achieved by

66

converting layer to kml file using conversion tool in Arc GIS tool box. The kml file is opened

67

in google earth and sampling points were identified. Sampling points shown in Fig.2 consists

68


of cultivable area and built-up area which was excluded while collecting samples. Soil

69

samples were collected from 40 sampling points in the study area which were numbered

70

sequentially from 1 to 40.



71


72

**Fig.2**View of grid map and sampling points in Google Earth

73

74 **Collection of soil samples**

75 The sampling points in the study area were identified with the help of hand-held Garmin  
76 Etrex 3X by using coordinates of that sampling points which obtained from GIS map (Fig. 3).  
77 Soil samples were collected from each sampling point as per the procedure and at each  
78 sampling point, four subsamples were collected at a depth of 15 cm. The surface trashes and  
79 litter were removed at sampling location and a 'V' shaped cut was made with the help of  
80 spade. Samples collected were mixed thoroughly and again checked for any small stones and  
81 other foreign materials. One kg of soil sample was obtained and taken as a representative  
82 sample by using four quartering rule. The samples were numbered and kept for air drying for  
83 two weeks for the analysis of soil nutrients.



84

85 **Fig. 3 Sampling points of KCAET campus**

86 **Analysis of soil samples**

87 The sieve analysis of air-dried soil samples was carried out for soil analysis with 2 mm sieve  
88 for pH and EC and 0.5 mm sieve for other soil nutrients. The soil analysis was carried out at  
89 soil testing laboratory, KVK, Malappuram. The soil samples were analysed for the soil  
90 chemical properties such as pH, Electric Conductivity, Available Nitrogen, Available  
91 Phosphorous, Available Potassium, Boron and Sulphur by using standard methods (Table 1).  
92 Based on the Table 2, soil chemical properties were classified as low medium and high in the  
93 study area.

94

95

96 **Table 1. Methods Used for soil analysis**

| Sl.No. | Soil Parameters               | Methodology   | Author                        |
|--------|-------------------------------|---|-------------------------------|
| 1)     | Soil pH                       | pH meter with glass Electrode (1:2 soil- water ratio) | Jackson (1973)                |
| 2)     | Electric conductivity (dS/m)  | EC meter  | Jackson (1973)                |
| 3)     | Organic Carbon                | Alkaline permanganate method                          | Walkley and Black (1934)      |
| 4)     | Available Nitrogen (kg/ha)    | Kjeldhal Method                                       | Subbiah and Asija (1956)      |
| 5)     | Available phosphorous (kg/ha) | Bray No1 extraction method                            | Bray and Kurtz. (1945)        |
| 6)     | Available potassium (kg/ha)   | Flame photometry                                      | Stanford and English (1949)   |
| 7)     | Boron (mg/kg)                 | Hot water extraction method                           | Gupta (1967)                  |
| 8)     | Sulphur (mg/kg)               | CaCl <sub>2</sub> Extraction method                   | Massoumi and Cornfield (1963) |

97 **Vasu et al. (2017)**

98 **Table2. Fertility Rating of soil chemical properties**

| Soil chemical property        | Nutrient status |           |       |
|-------------------------------|-----------------|-----------|-------|
|                               | Low             | Medium    | High  |
| pH                            | <6              | 6-7       | >7    |
| EC (dS/m)                     | <1              | 1 – 3     | >3    |
| Organic Carbon (%)            | <0.76           | 0.76 -1.5 | > 1.5 |
| Available Nitrogen (kg/ha)    | < 280           | 280– 450  | >450  |
| Available Potassium (kg/ha)   | < 115           | 115-275   | >275  |
| Available Phosphorous (kg/ha) | <10             | 10-24     | >24   |
| Boron (mg/kg)                 | < 0.5           | 0.5 -1    | >1    |
| Sulphur (mg/kg)               | < 10            | 10 – 15   | >15   |

99 **Kumar et al. (2018)**

100 **Preparation of fertility maps by using Arc GIS**

101 Fertility Maps were prepared by using Interpolation tool in Arc tool box  
 102 supported by Geoprocessing tool in Arc GIS.The Inverse Distance Weighting method (IDW)  
 103 in ArcGIS was used to interpolate the spatial distribution of soil pH, EC, N, P, K, B and S  
 104 from the soil samples collected from the study area. Inverse Distance Weighting method  
 105 (IDW) determines grid cell values by averaging of sample data points that are closer to the  
 106 cell. The closer point to the centre of the cell being estimated, the more influence or weight

107 has given in the averaging process(Anjana 2019).An Inverse Distance Weighted method of  
 108 interpolation created continuous maps for each soil parameter which helps to estimatethe soil  
 109 properties ofthe entire area.

## 110 **RESULTS AND DISCUSSION**

111 It is important to assess the fundamental nutrients of the soilto determine the available  
 112 nutritional status of the soil and to avoid the adverse effects of excess chemical fertilisers in  
 113 the soil, as well as to the environment.Table 3. showed the percentage of soil samples(40  
 114 samples) fall under low, medium and high range for different soil chemical properties. In the  
 115 study area,97.5% of the soil samples showed acidic in nature and all the samples showed low  
 116 electrical conductivity.In the case of organic carbon, 55 % of the samples showed the low  
 117 percentage of organic carbon, 20 % of the samples showed medium percentage and 25% of  
 118 the samples showed high percentage whereas in case of nitrogen about 75% of the soil  
 119 samples fall under low range and 25 % of the soil samples fall under medium range.About  
 120 32.5 % of samples fall under low and high range and 35 % of samples fall under the medium  
 121 level for phosphorous whereas for potassium, about 62.5% of samples fall under medium  
 122 range, 22.5% of samples under low range and 15% soil samples fall under high range of  
 123 potassium. In case of sulphur, about 85 % of the samples fall under low range and 15% of the  
 124 samples were under high level of sulphur whereas in case of boron,about 20% of the samples  
 125 falls under medium range and 80 % of the samples were under high range of boron.

126 **Table 3. Percentage of soil samples (40 samples) fall under low, medium and high range**  
 127 **for different soil chemical properties**

| Sl.No | Parameter               | Range  | Number of samples | Percentage of samples |
|-------|-------------------------|--------|-------------------|-----------------------|
| 1     | pH                      | Low    | 39                | 97.5%                 |
|       |                         | Medium | 1                 | 2.5%                  |
|       |                         | High   | -                 | -                     |
| 2     | Electrical Conductivity | Low    | 40                | 100%                  |
|       |                         | Medium | -                 | -                     |
|       |                         | High   | -                 | -                     |
| 3     | Organic Carbon          | Low    | 22                | 55%                   |
|       |                         | Medium | 8                 | 20%                   |
|       |                         | High   | 10                | 25%                   |

|   |             |        |    |       |
|---|-------------|--------|----|-------|
| 4 | Nitrogen    | Low    | 30 | 75%   |
|   |             | Medium | 10 | 25%   |
|   |             | High   | -  | -     |
| 5 | Phosphorous | Low    | 13 | 32.5% |
|   |             | Medium | 14 | 35%   |
|   |             | High   | 13 | 32.5% |
| 7 | Potassium   | Low    | 9  | 22.5% |
|   |             | Medium | 25 | 62.5% |
|   |             | High   | 6  | 15%   |
| 8 | Sulphur     | Low    | 34 | 85%   |
|   |             | Medium | -  | -     |
|   |             | High   | 6  | 15%   |
| 9 | Boron       | Low    | -  | -     |
|   |             | Medium | 8  | 32%   |
|   |             | High   | 32 | 80%   |


128

129 The spatial variability maps of soil fertility parameters helped to find the extent and  
130 magnitude of the nutrients. The spatial variability of soil nutrients such as N, P, K, B and S  
131 and pH and EC were plotted with the help of Arc GIS software. These maps showed the  
132 spatial distribution of soil nutrients in the study area. From the fig 4.a, it could be seen that  
133 the pH of the soils throughout the study area varying from strongly acidic to slightly acidic in  
134 nature. The pH of the major portion of the study area showed the strongly acidic in nature.  
135 The acidic nature of soil could be due to nature of parent material, micro topography, weather  
136 conditions and type of fertilizer used. Lime could be added in order to reclaim acid soils.  
137 Lime raises soil pH along with it provides calcium and magnesium to the soil. Electric  
138 conductivity was found to be low (<1 ds/m) in most parts of the study areas shown in Fig  
139 4.b which is mainly due to leaching of salts due to high rainfall.

140 Organic carbon was found to be low (< 0.76 %) in the most parts of the cultivated area of the  
141 study area as shown in fig 4.c. This may be due to erosion of top soils and decomposition of  
142 organic matter. Organic carbon was found to be medium and high in southern part of the  
143 study area. Available nitrogen varied from very low to medium in the study area. Available  
144 nitrogen was found to be low (<280 kg/ha) in cultivated parts of the study area (Fig 4.d). This  
145 is mainly due to low availability of Organic Carbon, increased rate of mineralisation and


146 removal of N by nutrient exhaustive crops. Nitrogen found to be medium in southern and  
 147 western parts of the study area. Results of the soil analysis showed that deficiency of nitrogen  
 148 in most parts of the study area. Therefore, proper soil management techniques should be  
 149 followed in order to improve the availability of nitrogen in the soil.

150 From the fig 4.e, it can be seen that, available phosphorus varied as low, medium and high  
 151 status in the study area and it was found to be high in cultivated parts of the study area. This  
 152 may be due to the application of phosphorous fertilizers or deposition from upland  
 153 areas. From the fig 4.f, it can be seen that available potassium varied from low to high range  
 154 in the study area. Potassium was found to be in medium range (115-275 kg/ha) in the study  
 155 area in all parts, except in some pockets, where it was found in the low range (<115k/ha).  
 156 Major part of the cultivated area showed the medium range of potassium. Low and medium  
 157 status of potassium may be due to significant loss of potassium due to excessive rainfall.  
 158 Potassium was found to be high (>275 kg/ha) in some pockets of southern part of the study  
 159 area. Sulphur varied from low to high status in the study area. In most parts of the study area  
 160 (Fig 4.g), sulphur was found to be in low range (<10 mg/kg). This is may be due to oxidation  
 161 of sulphur into sulphuric acid by soil micro-organisms which also resulted in low pH and also  
 162 leaching. In northern parts of the study area, sulphur was found to be high in some pockets.  
 163 From the fig.4.h, it could be seen that boron varied from medium to high range in the study  
 164 area. Boron was found to be high (>1 mg/ha) in most parts of the study area. This may be due  
 165 to irrigation of crops with well water and also application of fertilizers. Boron was found to  
 166 be satisfactory level in the study area as it was necessary for plant growth. From these maps, it  
 167 is evident that most of the soils were low in terms of Electrical Conductivity, Organic  
 168 Carbon, Nitrogen, and Sulphur. Potassium and Phosphorous were in medium range, whereas  
 169 boron was in the high range in the study area. Based on these maps nutrient recommendations  
 170 can be given to farmers to achieve site specific nutrient management.




177

178 a)




b)




186

c)




d)




187

188

e)



f)



197

198

g)

h)

199 **Fig.4 (a - h) shows the spatial distribution maps of soil chemical properties**

200 **Conclusions**

- 201 • In the study area, pH was ranging from 4.2 to 6.14 which indicated that soils were  
202 acidic and electric conductivity was ranging from 0.109 dS/m to 0.601 dS/m which  
203 was considered as low (< 1 dS/m) in the study area.
- 204 • Nitrogen (<280 kg/ha) and sulphur (<10 mg/kg) were in low range whereas boron  
205 (>1mg/kg) was in high range and the remaining chemical properties such as organic  
206 carbon (0.76-1.5 %), phosphorus (10-24 kg/ha) and potassium (115-275 kg/ha) were  
207 in medium range in the study area.
- 208 • GIS could be used as an effective tool for determining the spatial distribution of  
209 chemical properties of soils and thereby we can provide site specific nutrient  
210 recommendations.
- 211 • GIS based soil nutrient maps provide a better way to achieve right inputs in right  
212 quantity at right place at right time.
- 213 • GIS based soil nutrient maps, provide a way for achieving site specific nutrient  
214 recommendation as it involves lesser numbers of soil analysis and thus reduces the  
215 cost of operation compared to plot-to-plot analysis.

216 **Acknowledgement**

217 The authors are thankful to Kerala Agricultural University and Kelappaji College of  
218 Agricultural Engineering and Technology, Tavanur for sanctioning the research project and  
219 all financial assistance and also wish to express their gratitude to ICAR KVK, Malappuram  
220 and RARS, Pattambi for their assistance in soil analysis, providing information and helpful  
221 discussions.

222 **References**

- 223 Bhunia, G., Pravat Kumar Shit, Rabindranath Chattopadhyay.2018. Assessment of spatial  
224 variability of soil properties using geostatistical approach of lateritic soil (West Bengal,  
225 India). *An. of Agrarian Sci.* 16: 436–443.
- 226 Jeya, R., Vasanthakumar, R. and Dineshkumar, S., 2020. Knowledge level of sugarcane  
227 growers on precision farming practices. *Recent Trends in Agriculture towards Food*  
228 *Security and Rural Livelihood-Vol. 1*, p.36.

- 229 Kumar, D., Gopal, B. and Dhananjaya, B.C. 2018. GIS Based Site Specific Major Nutrient  
230 Maps and Recommendations for Coconut (*Cocos nucifera* L.) Gardens.Int.J.  
231 Curr.Microbiol. App.Sci (2018) 7(5): 3643-3654.
- 232 Gao Xue-song, XIAO Yi, DENG Liang-ji, LI Qi-quan, WANG Chang-quan, LI Bing, DENG  
233 Ou-ping, ZENG Min.2019. Spatial variability of soil total nitrogen, phosphorus and  
234 potassium in Renshou County of Sichuan Basin, China. J. of Integrative Agric.18(2):  
235 279–289.
- 236 Vasu, D., Singha, S.K., Nisha, S., Pramod, T. 2017. Assessment of spatial variability of soil  
237 properties using geospatial techniques for farm level nutrient management. *Soil and*  
238 *Tillage Research*. 169 (2017) 25–34.