

Original Research Article

Performance of different Common Hyacinth (*Hyacinthus orientalis*) to qualitative and quantitative traits of growth and propagation

ABSTRACT

Evaluation of Hyacinth cultivars including Jan Bose, Annabella, Aqua, Blue jacket, Purple sensation, Purple star, Yellow stone, Gipsy queen, Aladdin, Pink pearl genotypes was carried out during the year 2018 at Division Of Floriculture and Landscape Architecture SKUAST-K Shalimar ~~with an objective~~ to assess the performance of vegetative traits and propagation ratio. Data depicts the cultivar yellow stone with maximum plant height at maturity (17.97cm), Leaf length (12.98cm), Leaf width (3.64cm), Number of leaves per plant 24.58 and minimum values for Plant height at maturity (15.02cm), Leaf length (11.06cm), Leaf width (1.93cm), Number of leaves per plant 17.58 resulted with cultivar Pink pearl. Minimum to maximum values for bulb weight (28.87 to 34.53g), bulb Size (8.18 to 10.03cm), number of offsets per bulb (2.00 to 3.58), and weight of offsets (11.09 to 16.94g) were observed with cultivar pink pearl and yellow stone respectively. Most of the variations for growth and propagation were significant among the cultivars.

Keywords: Hyacinth cultivars, Evaluation, Vegetative traits, Propagation

1. Introduction

India imports around 20 crores of bulbs of Lilium, Tulip, Hyacinth, etc annually from Holland which costs huge foreign exchange. ~~Area-~~The area under bulbous crops globally is 50000 ha and India has only 3500 ha under this sector that too propagating Gladiolus in 1200 ha followed by tuberose at 800 ha. Ornamental temperate bulbous crops have an important place in the floriculture trade. Among these bulbous crops *Hyacinthus orientalis* one of the important bulbous crops which has been commercially used as it is having landscape use under temperate and subtropical climatic conditions. Hyacinth belongs to family Asparagaceae. These are commonly called as “Hyacinth”. *Hyacinthus* is native to the Eastern Mediterranean region including Turkey, Iraq, Turkmenistan, Iran, Lebanon, Syria, and the Palestine region. Mostly in European parts including the Netherlands, France, Italy, Croatia, Serbia, and Albania it is grown under naturalized conditions. Mass multiplication of bulbous plants is one of the important factors for the economic feasibility of commercial floriculture. To increase a particular variety of bulbs rapidly the primary concern is to see the feasibility of cultivars for propagation purposes. Most of the cultivated cultivars in India are imported ones and differ in performance as far as agro-agro-climatic conditions are concerned. It has been observed—a cultivars can perform ~~goodwell~~ from a flowering point of view and after the season is over the propagation ratio is almost a failure. This invites a thrust for regular import. To come over this issue screening and evaluation of cultivars is an important

Comment [11]: Insert values in abstract incorrect, remove, plz.

consideration and this can help in taking up a few cultivars which can be propagated for commercialization purposes. Keeping in view the huge cost of hyacinth bulbs ranging from Rs 40 to Rs 80 in the Indian market, this work has a significance for economic upliftment and minimizing the import.

Comment [12]: No references in introduction
?why?????

2. MATERIAL AND METHODS

2.1 Geographical features

Srinagar, the summer capital of Jammu and Kashmir, India is situated between 34°05' to 34°07' north latitude and 74°08' to 74°09' east longitude at an altitude of about 1587 m above mean sea level. It is flanked on the southeast and northeast by the lofty Himalayan ranges.

2.2 Morphological parameters

Studied parameters include plant height at maturity, leaf length, leaf width, number of leaves per plant, bulb weight, bulb size, number of offsets per bulb, the weight of offsets for Jan Bose, Annabella, Aqua, Blue jacket, Purple sensation, Purple star, Yellow stone, Gipsy queen, Aladdin, Pink pearl genotypes.

2.3 Statistical analysis

Statistical analysis of the data collected for different parameters during the present investigation was subjected to analysis of variance for a complete randomized block design with three replications (Gomez and Gomez, 1983).

3. Results and Discussion

Performance of Hyacinth genotypes (Table 1) showed significant variation for all the vegetative parameters. Maximum plant height at maturity (17.97cm), leaf length (12.98cm), leaf width (3.64cm), and number of leaves per plant 24.58 resulted with ~~in cultivar~~ Yellow stone ~~cultivar~~ where ~~as as~~ minimum values for plant height at maturity (15.02cm), leaf length (11.06cm), leaf width (1.93cm), number of leaves per plant 17.58 resulted with cultivar Pink pearl. Minimum to maximum values for bulb weight (28.87 to 34.53g), bulb size (8.18 to 10.03cm), number of offsets per bulb (2.00 to 3.58) and weight of offsets (11.09 to 16.94g) were observed with ~~cultivar~~ pink pearl and yellow stone ~~cultivars~~, respectively (Table 2). Most of the variations for growth and propagation traits were significant among the cultivars. Cultivar response to a-specific ~~agro-agro~~ climatic conditions is due to the suitability ~~in of a~~ particular ~~agro-agro~~ climatic condition. Vegetative growth depends upon the optimum and target climatic conditions received by particular species. Using metabolites is very important and varietal responses for the same differ which ultimately gives output for propagation ratio. Photoperiodic induction involves the production of a flowering stimulus in the leaves and its translocation to the stem apex under a certain day length. Endogenous circadian changes in activity, regulated by a physiological clock ~~is~~ ~~are~~ responsible for rhythmic quantitative and qualitative changes in ~~the~~ plants. This internal

clock allows the plants to measure time independently of the environmental periodicity and these factors contribute to its response ~~for to~~ vegetative growth and propagule ~~formation~~. ~~performance-Performance~~ of a cultivar differs in terms of its environmental suitability and genetic ~~potential~~. Masoodi *et al.* (2022) reported ~~co-co~~-related findings while working on altitude influence on different tulip cultivars. Phenotypic variation depends upon ~~the~~ genetic makeup of *Lilium* cultivars and the estimates of phenotypic and genotypic co-efficients of variance showed a low disparity for plant height and number of leaves indicating ~~the~~ effect of environment on different traits and phenotypic variability could be a reliable measure of genotypic variability (Grassotti *et al.* (1990), Balode (2010) and Masoodi *et al.* (2018)). Singh and Sen (2000) suggested that if the phenotypic coefficient of variation is greater than the genotypic co-efficient of variation, the apparent variation is not only due to genotypes, but also due to ~~the~~ influence of environment. Bhatia *et al.* (2013) reported variability in different cultivars of tulips for vegetative and flowering phases. ~~Co-Co~~-related findings were reported by Singh and Kumar (2008) in marigolds for plant height, Dhiman *et al.* (2015) in Asiatic hybrid lily, Masoodi *et al.* (2018) in Asiatic, Oriental, and La hybrids. Genotypic and phenotypic co-efficient of variation for ~~the~~ number of flowers per stem ~~were-was~~ reported in China Aster (Ravikumar and Patil 2003), Misra and Saini (1990) in gladiolus and ~~in~~ French marigold and ~~the~~ number of florets per spike in gladiolus. Singh and Sen (2000) suggested that if the phenotypic coefficient of variation is greater than the genotypic co-efficient of variation, the apparent variation is not only due to genotypes, but also due to ~~the~~ influence of ~~the~~ environment. The estimates of phenotypic and genotypic co-efficients of variance showed a low disparity for plant height, ~~the~~ number of leaves per plant, and inflorescence diameter indicating the least effect of environment on different traits and phenotypic variability could be a reliable measure of genotypic variability. Bhatia *et al.* (2013), and Panse and Sukhatme, (1985) reported low genotypic co-efficient of variation helps in the measurement of genetic diversity in the qualitative and quantitative characters. The estimation of heritability has a greater role to play in determining the effectiveness of selection provided it is considered in conjunction with the predicated genetic advance.

Comment [13]: Insert reference here, plz

Comment [14]: Insert reference here, plz.

Comment [15]: Rewriting plz.

Table : 1: Performance of Hyacinth cultivars to growth traits

Treatment Symbol	Plant height at maturity (cm)	Leaf length (cm)	Leaf width (cm)	Number of leaves per plant
Jan Bose	17.56	11.81	2.74	22.33
Annabella	17.05	11.10	2.03	21.58
Aqua	17.29	11.48	2.77	22.26
Blue jacket	16.73	11.18	2.31	19.33
Purple sensation	16.05	11.43	2.41	19.58
Purple star	17.29	11.86	2.51	20.16
Yellow stone	17.97	12.98	3.64	24.58
Gipsy queen	15.93	11.42	2.74	18.83
Aladdin	16.04	11.46	2.57	18.33
Pink pearl	15.02	11.06	1.93	17.58
C.D(p≤0.05)	0.07	0.07	0.20	0.19

Table-2: Response of Hyacinth cultivars to propagation ratio

reatment Symbol	Bulb weight (g)	Bulb Size(cm)	Number of offsets per bulb	Weight of offsets(g)
Jan Bose	30.25	9.59	3.22	16.79
Annabella	29.77	9.12	3.00	16.15
Aqua	29.40	9.34	3.16	16.39
Blue jacket	30.85	8.90	3.11	13.10
Purple sensation	31.44	9.30	3.11	14.49
Purple star	31.86	9.13	3.10	14.56
Yellow stone	34.53	10.03	3.58	16.94
Gipsy queen	29.61	8.30	2.66	11.17
Aladdin	30.63	9.63	2.91	11.05
Pink pearl	28.87	8.18	2.00	11.09
C.D($p \leq 0.05$)	0.76	0.05	0.18	0.49

Fig 1: Performance of Hyacinth cultivars to growth traits





Fig 2: Response of Hyacinth cultivars to propagation ratio



References:

- Balode, A. 2010. Phenotypic analysis of hybrids and their parents in liliium species. *Breeding, Annual 16th International Scientific Conference Proceedings. Research for Rural Development 1*: 65-70.
- Bhatia, R., Dhiman, M. R., Chander, P. and Dey, S. S. 2013. Genetic variability and character association in tulip (*Tulipa gesneriana*) for various quantitative traits. *Indian Journal of Agricultural Sciences* 83 (7): 773-780.
- Dhiman, M. R., Parkash, C., Kumar, R., Guleria, M. S. and Dhiman, M. 2015. Studies on genetic variability and heritability in Asiatic hybrid lily (*Lilium x elegans* L.). *Molecular Plant Breeding* 6: 1-8.
- Gómez, K. and Gómez, A. (1983) *Statistical Procedures for Agricultural Research*. 2nd Edition, John Wiley & Sons, Hoboken, 630 p.
- Grassotti A., Torrini F., Mercuri A., and Schiva T., 1990, Genetic improvement of liliium in Italy. *Acta Horticulture* 266: 339-347.
- Misra, R. L. and Saini, H. C. 1990. Correlation and path-coefficient studies in gladiolus. *Indian Journal of Horticulture* 47 (1): 127-132.
- Masoodi N.H. and Nayeem S.M. 2018. Evaluation Of Different Lilium Hybrids Under Climatic Conditions Of Kashmir Valley. *Agri Res & Tech :Open Access J.* 12 (8): 2581-8627
- Masoodi N.H. and Nayeem S.M. 2022. Morphological characterization of Tulip cultivars at 1587m above Mean sea level under Kashmir Himalayas. *International Journal of environment and climate chnage Tech :Open Access J.* 17 (1): 556008.
- Panse, V. G. and Sukhatme, P. V. 1985. Statistical methods for agricultural workers. *Indian Council of Agricultural Research, New Delhi* 152-156.
- Ravikumar, H. and Patil, V. S. 2003. Evaluation of China aster [*Calcestephus chinensis*(L.) Ness] genotypes under transitional zone of north Karnataka. *Indian Society of Ornamental Horticulture* 12-14.
- Singh, D., and Sen, N. L. 2000. Genetic variability, heritability and genetic advance in marigold. *Journal of Ornamental Horticulture* 3(2): 75-78.
- Singh, Deepthi. and Kumar, S. 2008. Studies on genetic variability, heritability, genetic advance and correlation in marigold. *Journal of Ornamental Horticulture* 11(1): 27-31.