

Original Research Article

EXTENT OF ADOPTION OF RECOMMENDED PRACTICES OF COTTON CULTIVATION BY THE COTTON FARMER

ABSTRACT

Aims: To know the adoption level of recommended cultivation practices of cotton crop.

Study Design: The research design used for this study was an Ex-post Facto.

Place and Duration of Study: The study was conducted in Khandwa district of Madhya Pradesh and duration 2020-2022

Methodology: The study was conducted in Khandwa district of Madhya Pradesh a total of 200 farmers selected by following the proportionate random sampling technique. An interview schedule was prepared for collection data relating to the adoption of recommended practices by the respondents.

Results: The study, ~~it could be concluded~~ that, with respect to the adoption, majority of the farmers (69.00%) had a medium level of adoption of recommended improved practices of cotton cultivation while, an equal number (15.50%) of the respondents had low and high level of adoption ~~of recommended cotton cultivation practices~~ the same practices. Among the socio-economic and personal, communication and psychological characteristics of the respondents, ~~namely~~ age, education, family size, family type, land holding, annual income, farm mechanization, extension participation, information seeking behaviour, cosmopolitaness, economic motivation, scientific orientation and risk orientation indicated significant association with the extent of adoption of the recommended practices of cotton.

Conclusion: The study concluded that majority of the cotton growers were ~~low adopted recommended cotton practices~~.

Comment [Ma1]: Differs from the final results.

Key words: Adoption level, Cotton, Growers, Production technology, Respondents

INTRODUCTION

Cotton (*Gossypium spp.*) is one of India's most significant cash crops and sources of fibre, and it is vital to the nation's industrial and agricultural economics (Jaglan et al. 2012). The total area planted with cotton in the world in 2020–2021 was 31.66 million hectares, and the crop produced 113.11 million bales, or 778 kg/hectare, on that area. In terms of both total area and production, India has overtaken other countries to become the world's top cotton producer. India ranked first among the top exporters of cotton in the world. India occupied with 5.5 million bales, Canada ranks third after the United States (16.25 million bales) and Brazil (10.70 million bales) (USDA, 2020-21). In Madhya

Pradesh, cotton occupies an area of about 0.6 million hectare with a production of 19 lakh bales and with a productivity of 527 kg/ha (2020-21) (Deshpande 2020). The productivity of cotton is very poor and this crop is prone to disease and pest infestation attacked by a number of insect, pests, diseases, nematodes and weeds. This marks the essentiality of Mmodern agricultural technologies are essential to improve yields and increase climatic as well as disease resistance to climatic risks and attack of insect, pests, diseases and nematodes. Present studySo, the aims of the study was to determine the adoption of improved cotton producing technologies among the level of cotton growers about cotton production technology and also finds association between selected socio-economic & personal, communication and psychological characteristics of cotton growers and their adoption of improved cotton production technology (Vishnuvardhan *et al.* 2022).

Comment [Ma2]: How can aim of a study be a reference of any other author? The aim has to be solely decided by the researcher.

MATERIALS AND METHODOLOGY

The research study on adoption level of cotton growers about cotton production practices was conducted during the year 2021-2022 in Pandhana and Khandwa block of Khandwa district in (Madhya Pradesh). In the present investigation, descriptive type of ex-post-facto research design was employed. This design was appropriate because the phenomenon had previously happened. Ex-post-facto research is the most logical empirical enquiry as in which the researcher does not have any control over independent variables; as their appearance has already occurred or as they are inherent and cannot be manipulated. Thus, inferences about relations among variables were made without direct intervention from concomitant variation of independent and dependent variables (Kerlinger, 1972). Khandwa district comprises of 7 blocks namely Khandwa, Punasa, Pandhana, Harsud, Chhaigaon, Baldi and Khalwa. Out of these two blocks namely Pandhana and Khandwa block were purposively selected, because of highest area under cotton crop. The Pandhana block comprise of 124 villages and Khandwa block comprise of 97 villages. Out of which ten villages were selected on the basis of larger area under cotton crop. From the ten selected villages, 200 cotton growers were selected from each selected village by using proportionate random sampling method to make the total sample size for the present study.

Comment [Ma3]: Mention the total sample size. If 200 growers are selected from each of ten villages. The sample size stands as 2000. Is it what the researcher mean?

Comment [Ma4]: Sampling of the respondents and total respondent number is not very clear. Please check it carefully.

The adoption level of cotton growers with respect to improved cultivation practices of cotton was studied by computing adoption score. Concerning adoption, the responses were rated on three-point continuum, namely complete adoption, partial adoption and no adoption. A numerical score of 2 was assigned for complete adoption, a score 1 was assigned for partial adoption and a zero was assigned for no adoption. The score of all identified practices was then summed up. This sum total indicates the adoption score of that particular individual respondent. The scores of all the practices earned by each individual were added together to work out the adoption score of an individual. The raw score then was converted into adoption index as mentioned below.

Comment [Ma5]: Indicates same meaning as the previous line.

Total adoption score obtained
 Adoption Index = ----- x 100
 Maximum obtainable adoption score

On the basis of adoption index, the respondents were categorized in to low, medium and high

groups based on mean \pm sd

The primary data were collected personally by the researcher by interviewing the selected respondents with the help of semi structured interview schedule.

RESULTS AND DISCUSSION

Adoption level of the cotton growers about recommended cotton production technology:

Adoption level of the cotton growers regarding different practices of cotton cultivation is presented in Table 1. The perusal of data clearly indicated that the majority of the cotton growers had were fully or partially adopted recommended land preparation practice (91.5%), method of sowing (74%), recommended dose of fertilizer (76.5%), application of FYM/compost (66.5%), use of improved varieties of cotton (63.5%), insect management and seed treatment (61.5%), whereas, nearly more than half of the respondents were fully or partially adopted recommended seed rate (56.50%), intercropping (55.5%) and time of irrigation (51.50%). Furthermore, the data indicated that recommended practices like majority of the respondents were not adopting recommended striking trap (74.50%), topping operation (62.50%), intercultural operations (61.50%), and disease management (54.50%) were not adopted by majority of the respondents. The findings are in line with the findings of Jat et al. (2022) and Bondarwad et al. (2010).

Table 1: Practice wise adoption level of the cotton growers about recommended cotton production technology

(n = 200)

S. No.	Practices	Adoption Level				
		Complete Adoption	Partial Adoption	No Adoption	Mean	Rank
1	Preparation of land	49 (24.50)	134 67.00	17 (8.50)	116	I
2	Use of Improved Varieties of cotton	38 (19.00)	89 (44.50)	73 (36.50)	82.50	V
3	Method of sowing	57 (28.50)	91 (45.50)	52 (26.00)	102.50	II
4	Seed rate	34 (17.00)	53 (26.50)	113 (56.50)	60.50	XI

Comment [Ma6]: Discussion part is missing. It is only the results that had been written without interpretations.

Comment [Ma7]: Give the data for full and partial adoption separately. Do it for all.

Comment [Ma8]: Does not match with the data as per table 1. PI Check.

Comment [Ma9]: Does not match as per data provided in Table 1.

Comment [Ma10]: Check the total number of respondent. As mentioned in the methodology, the number stands different. Cross check.

5	Seed treatment	28 (14.00)	93 (46.50)	79 (39.50)	74.50	VIII
6	Application of FYM/Compost	46 (23.00)	85 (42.50)	69 (34.50)	88.50	IV
7	Time of Irrigation	38 (19.00)	59 (29.50)	103 (51.50)	67.50	X
8	Recommended dose of fertilizer	36 (18.00)	117 (58.50)	47 (23.50)	94.50	III
9	Application of weedicide	42 (21.00)	59 (29.50)	89 (49.50)	71.50	IX
10	Intercultural operations	33 (16.50)	44 (22.00)	123 (61.50)	55.00	XIII
11	Insect management	40 (20.00)	81 (40.50)	79 (39.50)	80.50	VI
12	Disease management	22 (11.00)	69 (34.50)	109 (54.50)	56.50	XII
13	Intercropping	47 (23.50)	64 (32.00)	89 (44.50)	79.00	VII
14	Topping operation	24 (12.00)	51 (25.50)	125 (62.50)	49.50	XIV
15	Striking trap	24 (12.00)	27 (13.50)	149 (74.50)	37.50	XV

Figures in parenthesis indicate percentage

Overall adoption level of the cotton growers:

It is evident from the data reported in Table 2 revealed that, nearly two third of the respondents (69.00%) had a medium level of adoption of recommended improved practices of cotton cultivation while, an equal ~~number-percentage~~ (15.50%) of the respondents had low and high level of adoption of recommended cotton cultivation practices. Thus, it may be inferred from the data that the majority of cotton growers had medium level of adoption regarding recommended improved practices of cotton cultivation. The probable reason might be that the majority of the respondents had medium level of knowledge ~~would have that~~ motivated them to adopt ~~a number of the~~ technologies fully or partially. Besides ~~the respondents with~~ medium risk orientation, scientific orientation, extension participation and information seeking behaviour ~~of the respondents also influenced would have~~ adopted of the

Formatted Table

latest recommended technologies to some extent. The findings are in line with the findings of Rajput and Chinchmalatpure (2016), Kumar *et al.* (2017) and Mahendrakar *et al.* (2018).

Table 2: Distribution of cotton growers according to their adoption level

S. No.	Categories	Frequency	Percentage
1.	Low (Up to 39 Score)	31	15.50
2.	Medium (40 to 64 Score)	138	69.00
3.	High (Above 64 Score)	31	15.50
Total		200	100.00

(Mean = 51.31, S.D=12.20)

Association between Socio-economic personal, communication and psychological characteristics of cotton growers and their adoption level:

It is apparent from the Table 3 that out of fourteen independent variables seven variables namely age, education, farm mechanization, economic motivation, family size, scientific orientation and risk orientation had shown significant association at 0.01 level of probability with adoption level of the cotton growers, while other six variables namely type of family, annual income, land holding, extension participation, cosmopolitaness and information seeking behaviour had significant association at 0.05 level of probability with the dependent variable, adoption level of the cotton growers.

Therefore the null hypotheses were rejected and original propositions that there would be association between age, education, farm mechanization, economic motivation, scientific orientation, risk orientation, family size, type of family, annual income, land holding, extension participation, information seeking behaviour and cosmopolitaness of cotton growers and their adoption level were accepted. The findings are in line with the findings of Singh *et al.* (2019), and Bishnoi *et al.* (2016).

The remaining one variable viz; caste, had no significant association with adoption level of the cotton growers. Hence, the null hypothesis was accepted and original proposition that there would be association between caste, of cotton growers and their adoption level was rejected. The findings are in line with the findings of Verma *et al.* (2019).

Table 3: Association between socio-economic personal, communication and psychological characteristics of cotton growers and their adoption level

S. No.	Independent variables	Chi square value (χ^2)
1.	Age	35.44**
2.	Education	64.29**
3.	Caste	7.33 ^{ns}

4.	Family size	13.79**
5.	Type of family	6.99*
6.	Annual income	19.76*
7.	Land holding	24.61*
8.	Farm mechanization	40.98**
9.	Extension participation	10.19*
10.	Information seeking behaviour	10.97*
11.	Cosmopolitaness	17.13*
12.	Economic Motivation	44.39**
13.	Scientific Orientation	67.65**
14.	Risk orientation	39.93**

*Significant at 0.05 level of probability

** Significant at 0.01 level of probability

NS = Non-Significant

CONCLUSION

The study concluded that majority of the cotton growers ~~had were~~ adopted recommended ~~practices of methods for land~~ preparation of land, ~~recommended method of~~ sowing, ~~recommended dose/quantity of~~ fertilizer ~~dose~~, ~~recommended dose of~~ FYM/compost ~~application~~, use of improved varieties ~~of cotton~~ and ~~recommended~~ insect management, while, very few of them were adopted recommended seed treatment, ~~recommended application of~~ weedicide ~~application for weed control~~, ~~recommended timely of~~ irrigation, ~~recommended~~ seed rate, ~~recommended~~ disease management, ~~recommended~~ inter-cultural operations, ~~recommended~~ topping operation, ~~recommended~~ striking trap and use of light trap. The study also indicated that the majority of cotton growers had medium level of adoption ~~of the~~ ~~concerning~~ ~~practices~~ ~~cotton~~ ~~production~~ ~~technology~~. The study also concluded that adoption level of cotton growers was significantly associated with their age, education, farm mechanization, economic motivation, scientific orientation risk orientation, family size, family type, annual income, land holding, extension participation, cosmopolitaness and information seeking behaviour. Hence, to increase the adoption level of cotton growers, ~~the~~ various training programmes ~~and including~~ method ~~and~~ result demonstrations should be organised on farmers' fields, ~~the~~ Extension agencies may also ~~be~~ used information communication technologies for motivation of cotton growers towards cotton production technologies and required inputs should made available in time.

REFERENCES

Bishnoi M, Sisodiya SS, Kumar V. Adoption of farmers about bt. cotton production technology in Bhilwara district of Rajasthan. Int. J. Agric. Sci.2016;8(60):3353-3356.

Bondarwad SP, Wangikar SD, Deshmukh ND. Present status of adoption of bt. cotton production technology by farmers. Indian J. Ext. Educ.2010;5(3&4):322-324.

Deshpande T. State of agriculture in India. PRS Legislative Research 2020;53(8):6-7.

Jaglan RS, Singh N, KANU P. Impact of climate change on cotton and its production in India. Silver Jubilee International Symposium, 2012vol. 271, p. 271.

Jat M, Jaiswal DK, Saharawat YS. Extent of knowledge and adoption of recommended wheat production practices among wheat growers in malwa region (M.P.). Indian J. Ext. Educ. 2022;58(1):40-43.

Kerlinger FN. Draft report of the APA committee on ethical standards in psychological research: A critical reaction. Am. Psychol.1972;27(9):894-896.

Kumar P, Pavan R, Dhorey RK. Extent of adoption of farmers about bt. cotton practices in Warangal district of Telangana State, India. Int. J. Curr. Microbiol. Appl. Sci.2017;6(11):824-829.

Mahendrakar M, Syed J, Mazhar H. Knowledge and adoption level of respondents about bt. cotton production practices in Shahapur Taluk of Yadgir district Karnataka. Int. J. of Innov. Sci. Res. Technol.2018;3(6):9-12.

Rajput HD, Chinchmalatpure UR. Knowledge and adoption of bt. cotton cultivation practices. Indian J. Ext. Educ. 2016;52(1 & 2):121-123.

Singh S, Yadav D, Hariyale V. Study on adoption of integrated pest management practices by cotton growers in Khandwa district. Int. J. Curr. Microbiol. Appl. Sci. 2019;8(10):430-435.

Verma AK, Doharey RK, Dubey SK, Bishnoi SS, Prakash O, Prashad K 2019. Extent of knowledge and adoption of cotton growers in Bharuch district of Gujarat. Indian J. Ext. Educ. 2019;55(4):87-91.

Vishnuvardhan KM, Reddy BV, Kalyani DL, Krishna MS, Reddy YR. A high yielding cotton variety with high fibre quality and sucking pest tolerance suitable for both Central and South zones of India. Electronic Journal of Plant Breeding, 2022;13(2):739-744.

Comment [Ma11]: Use same format

Formatted: Normal, Indent: Left: 0", Hanging: 1", Space After: 6 pt, Line spacing: 1.5 lines, Pattern: Clear (White)