

## Case study

### MYOCARDIAL INFARCTION REVEALING NON-COMPACTION OF THE LEFT VENTRICLE: A RARE CASE-BASED DISCUSSION.

#### Abstract :

Leftventricular non compaction (LVNC)isacardiomyopathy due to disturbances in the normal development of the heart muscle **lead** to the persistence of hypertrabeculation. its incidence is more common in males.

Comment [MF1]: leading

The mostcommonclinicalpresentations at the time of **first diagnosis** are heartfailure, thromboembolicevents and variousrhythmdisorders.

Comment [MF2]: the

We report the case of a 52 yearold man, with a probable unexploredhistory of cerebrovascular accident, whoconsultedinthe emergency room for acute chest pain constrictive **restrosternale**.

Comment [MF3]: retrosternale

The electrocardiogramshowednegative T waves in the lateral and inferior septum. The echocardiographyshowed a trabeculated aspect of the apex of the leftventricle, intertrabecularrecesses takingcolor on color Doppler. Ultra-sensitive troponinswereelevated to 351 ng/l.

**A** coronaryangiographywasperformedreturning normal.

Comment [MF4]: delete

Myocardialischemiawithhealthycoronariessecondary to non-compaction of the LV remainspoorlyunderstood and maybesecondary to an anomaly of the microcirculation, coronaryembolization of a thrombus formed at the level of the ventriculartrabeculations.

The dreadful complications of **discovery** of LVNC must draw the cardiologist's attention to thispathology for bettersubsequent management.

Comment [MF5]: the

#### Keys words :

**Leftventricular non-compaction, myocardialinfarction, case report**

#### Introduction :

Comment [MF6]: Add references

Leftventricular non-compaction (LVNC) or previouslycalledembryonicspongyleftventricleis characterised by the presence of excessive trabeculations of the leftventricle, formingdeepcavitiesthatfillwithblood.

It is due to disturbances in the noncompactionprocess of multifactorialoriginleading to the persistence of hypertrabeculation and LVNC.

The most frequent clinical presentations at the time of first diagnosis are heart failure, thromboembolic events and various rhythm disorders.

Comment [MF7]: the

We report the case of a 52-year-old patient admitted with acute non-ST-segment elevation coronary syndrome in whom LVNC was discovered incidentally on echocardiography. There are few similar cases in the literature.

### Case presentation :

We report the case of a 52-year-old man, a 20-year-old smoker, with no personal or family history of cardiovascular pathology or sudden death. His symptoms date back to two months before his admission with the onset of constrictive retrosternal chest pain at rest, which gave way spontaneously; the patient consulted the emergency room. In addition, the patient reported a transient hemiparesis occurring 3 years before his admission, which was not the subject of a consultation or exploration.

On admission, the patient was conscious, hemodynamically and respiratory stable, eupneic, normocardial at 95 bpm and normotensive at 110/65 mmHg. The physical examination did not reveal any heart murmurs or signs of cardiac insufficiency with normal pleuropulmonary auscultation. The neurological examination did not reveal any motor or sensory deficit.

The electrocardiogram showed negative T waves in the lateral and inferior leads. The echocardiography showed a trabeculated aspect of the apex of the left ventricle, intertrabecular recesses taking colour on colour Doppler, an aspect of non-compacted tissue at the apical level with a ratio of compacted to non-compacted tissue at 3.14. (figure 1.2.3) The echocardiography did not reveal any segmental kinetic disorder, nor any individualizable thrombus, with a left ventricular ejection fraction (LVEF) preserved at 55%. The chest X-ray was without abnormality. The coronary angiography performed within 24 hours was normal with no atheromatous plaques or thrombotic lesions. A Holter ECG was performed and found no supraventricular rhythm disorder. Cardiac MRI was not performed because it was not available.

In view of the neurological symptoms described by the patient, a cerebral scanner was performed, showing ischemic lesions of a sequential nature.

On biological examination, the ultra-sensitive troponins were elevated to 351 ng/l, a microcytic hypochromic anaemia at 9.7 g/dL, and a correct renal function. A thrombophilia test did not reveal any abnormality.

Faced with a myocardial infarction with non-obstructive coronary arteries on a non-compaction of the left ventricle, the possible diagnoses were a myocardial infarction (MI) due to an attack on the microcirculation found in the non-compaction of the LV, or the embolisation of a thrombus formed at the apical level of the LV despite its non-visualisation on echocardiography and the absence of thrombus at the level of the coronaries on coronary angiography, or a thrombus linked to atrial fibrillation in the context of LVNC. Type 2 MI was also suspected in the presence of iron deficiency anaemia. The most likely mechanism

in the face of coexisting sequelae of ischemic brain lesions and embolisation from a thrombus in the trabeculations not visualised on echocardiography or paroxysmal atrial fibrillation (AF).

The patient was treated medically and put on anticoagulation with vitamin K antagonist (VKA).

Comment [MF8]: with vitamin

## DISCUSSION

Left ventricular non-compaction describes a phenotypic phenomenon characterised by the presence of excessive trabeculations of the left ventricle, forming deep cavities that fill with blood. The first autopsy report describing an "embryonic spongy pattern of the left ventricle" in newborns was published in 1926 [1]. The first case of an isolated sinusoidal anomaly of the left ventricle diagnosed by echocardiography in a 33-year-old woman was published in 1984 [2].

Its classification as a cardiomyopathy differs between the European Society of Cardiology and the American Heart Association. The former defines it as an "unclassified cardiomyopathy", and the latter as a "genetic cardiomyopathy" [3, 4].

The prevalence is 4.1-5%, and men are almost three times more affected [5].

Comment [MF9]: This paragraph moves to the introduction

Disturbances in the normal development of the heart muscle lead to the persistence of hypertrabeculation. Trabeculations appear at the end of the 4th week of gestation. The thin subepicardial part of the muscle forms the "compacted myocardium" while the trabecular part forms the "non-compacted" part. In the following weeks, the trabeculae thicken, increasing the volume of the compacted layer, while the intertrabecular recesses compress and form capillaries [6]. Disruptions in this process, for whatever reason, lead to the persistence of hypertrabeculation and LVNC. Normally, the process of compaction progresses from the epicardium to the endocardium, from the septum to the free wall and from the base to the apex in the LV. Moreover, it is more often observed in the LV than in the right ventricle. Therefore, apical non-compaction is the most common type.

Furthermore, the presence of many cases of familial LVNC argues for a genetic origin of the disease and about 66 genes have been considered to be associated with LVNC, 82% of which code for sarcomeric proteins [7]. In patients with LVNC, various concomitant congenital heart diseases and neuromuscular disorders have been reported.

Furthermore, hypertrabeculation has not always been pathological and may be reversible. It has been observed as an adaptive phenomenon to increased cardiac output in athletes, pregnant women and patients with haemoglobinopathy [8]. The trabeculated ventricle can function more efficiently by generating the same systolic ejection volume at lower strain and wall stress.

The most common clinical presentations at the time of first diagnosis are heart failure, thromboembolic events and various rhythm disorders.

Ventricular dysfunction may be systolic or diastolic. Although the mechanism of heart failure associated with LVNC is not clear, it has been suggested that myocardial ischaemia may play an important role. This is supported by

subendocardial perfusion deficits that have been observed on cardiac MRI and PET scans that will lead to subendocardial fibrosis and necrosis and these perfusion deficits unrelated to atherosclerosis are explained by an overly thick subendocardial uncompact layer, preventing the endothelial myocardium from contacting the epicardial coronary arterial system, thereby causing ischaemia.

Comment [MF10]: Add references

The incidence of atrial fibrillation ranges from 15 to 23% and may lead to systemic embolisation. It has been observed more frequently in patients with reduced left ventricular systolic function [9]. Ventricular tachyarrhythmias are more common and occur in 47% of cases [10]. Ventricular fibrillation is a common cause of sudden death in patients with LVNC. Thromboembolic complications can occur in up to 38% [11]. Excessive trabeculation and atrial fibrillation of the left ventricle have been shown to be a risk factor for thromboembolic event [12].

Echocardiography is the most widely used diagnostic tool. Many diagnostic criteria have been proposed and include the presence of at least three trabeculations, the presence of blood flow between the trabeculations. In addition, the ratio of compacted to non-compacted myocardium must be  $>2$  at end diastole, and without secondary causes. Finally, the absence of a "golden standard" makes diagnosis very difficult. Cardiac magnetic resonance has become the method of choice for diagnosing LVNC [13].

LVNC has no specific therapy to date. Treatment should be applied according to the clinical situation. In patients with decreased LVEF, treatment of heart failure should be applied according to guidelines [14].

In the prevention of embolic events, anti vitamin K and new oral anticoagulant (NOAC) can be considered. In patients with LVNC and AF or a history of embolic events, NOAC is the first choice. In patients with left ventricular thrombus, AVK are the drugs of choice.

There is no agreement on treating patients with left ventricular dysfunction without a history of atrial fibrillation with anticoagulants or aspirin. In patients with normal LVEF, the decision can be made taking into account the CHADS 2 score. Chimenti et al. suggested that patients with a CHADS 2 score  $>2$  should be started on NOACs [15].

Ischaemia in LVNC can be explained by several mechanisms. The association of LVNC with coronary artery disease is considered rare, and may be explained by a common genetic predisposition to LVNC and MI. There is evidence of an association between MI and the activity of cytochrome P450 2C9 and sarcomere-related proteins such as the calcium-binding protein S100A [17]. When LVNC is diagnosed and the genes for one or more of these proteins are mutated, an increased risk of MI should be suspected. A coronary angiography study in LVNC subjects showed that 29% of patients had significant coronary artery disease [18]. In addition, microcirculatory dysfunction has also been proposed as a mechanism of myocardial ischaemia in LVNC. This was supported by post-mortem analyses of LVNC hearts, which showed ischaemic subendocardial lesions. [19] A previous study concluded that coronary reserve decreases in LVNC patients in both compacted and non-compacted heart muscle, including the normal epicardial coronary system [18]. In

Comment [MF11]: Reference number 16 is not present, correct the sequence of references.

addition, LVNC may affect the progression of remodelling in patients with MI and may worsen their prognosis. Associated with these mechanisms is the embolic mechanism secondary to thrombus in the trabeculations of the LV or secondary to atrial fibrillation common in the LVNC.[20]

Although the relationship between LVNC and coronary heart disease is not well understood, a number of papers have reported an association between NCC and acute myocardial infarction.

A few cases have been reported including a 73-year-old patient with left ventricular non-compaction (LVNC) who was diagnosed with acute myocardial infarction (MI), with atheromatous involvement of all three coronary arteries and intraventricular thrombus [21].

Arun Gopi reported a 19-year-old man with signs of heart failure.[22] TTE revealed LVNC with severe LV dysfunction. He was put on medical treatment including oral anticoagulants. Despite this, 6 months later he presented with an anterior wall MI. The coronary angiography revealed a thrombotic .

Two other patients were reported, aged 20 years, admitted separately for ST-segment elevation ACS, in whom coronary angiography showed normal coronaries. And in whom the TTE found non-compaction of the LV.[23] These latter cases are similar to our patient who was admitted for non-ST-segment elevation ACS, with echocardiography showing LVNC without thrombus and coronary angiography showing normal arteries.

#### **Conclusion :**

LVNC is a rare condition but one that should be kept in mind by the cardiologist as it mimics several cardiac conditions.

LVNC is almost always discovered following complications such as heart failure, thromboembolic events or following a myocardial infarction as our case illustrates.

The limited but growing number of reported cases provides evidence of an association between NCVG and MI. And because of this, patients with NCVG are considered at high risk for MI. Further studies are needed for a better comprehension of the mechanisms incriminated and codification of the management of this pathology.

#### **REFERENCES**

- 1-Grant, R.T. An unusual anomaly of the coronary vessels in the malformed heart of a child. *Heart* **1926**, *1*, 273–283
- 2-Engberding, R.; Bender, F. Identification of a rare congenital anomaly of the myocardium by two-dimensional echocardiography: Persistence of isolated myocardial sinusoids. *Am. J. Cardiol.* **1984**, *53*, 1733–1734
- 3-Elliott, P.; Andersson, B.; Arbustini, E.; Bilinska, Z.; Cecchi, F.; Charron, P.; Dubourg, O.; Kühl, U.; Maisch, B.; McKenna, W.J.; et al. Classification of the cardiomyopathies: A position statement from

the European Society Of Cardiology Working Group on Myocardial and Pericardial Diseases. *Eur. Heart J.* **2008**, *29*, 270–276

4-Maron, B.J.; Towbin, J.A.; Thiene, G.; Antzelevitch, C.; Corrado, D.; Arnett, D.; Moss, A.J.; Seidman, C.E.; Young, J.B.; American, H.A.; et al. Contemporary definitions and classification of the cardiomyopathies: An American Heart Association Scientific Statement from the Council on Clinical Cardiology, Heart Failure and Transplantation Committee; Quality of Care and Outcomes Research and Functional Genomics and Translational Biology Interdisciplinary Working Groups; and Council on Epidemiology and Prevention. *Circulation* **2006**, *113*, 1807–1816

5-Gerecke, B.J.; Engberding, R. Noncompaction Cardiomyopathy-History and Current Knowledge for Clinical Practice. *J. Clin. Med.* **2021**, *10*, 2457.

6-Towbin, J.A.; Jefferies, J.L. Cardiomyopathies Due to Left Ventricular Noncompaction, Mitochondrial and Storage Diseases, and Inborn Errors of Metabolism. *Circ. Res.* **2017**, *121*, 838–854.

7-van Waning, J.I.; Caliskan, K.; Hoedemaekers, Y.M.; van Spaendonck-Zwarts, K.Y.; Baas, A.F.; Boekholdt, S.M.; van Melle, J.P.; Teske, A.J.; Asselbergs, F.W.; Backx, A.P.C.M.; et al. Genetics, Clinical Features, and Long-Term Outcome of Noncompaction Cardiomyopathy. *J. Am. Coll. Cardiol.* **2018**, *71*, 711–722.

8-D'Silva, A.; Jensen, B. Left ventricular non-compaction cardiomyopathy: How many needles in the haystack. *Heart* **2021**, *107*, 1344–1352.

9-Engberding, R.; Stöllberger, C.; Schneider, B.; Nothnagel, D.; Fehske, W.; Gerecke, B.J. Abstract 14769: Heart Failure in Noncompaction Cardiomyopathy - Data from the German Noncompaction Registry (ALKK). *Circulation* **2012**, *126*, A14769.

10-Towbin, J.A.; Lorts, A.; Jefferies, J.L. Left ventricular non-compaction cardiomyopathy. *Lancet* **2015**, *386*, 813–825.

11-Chimenti, C.; Lavallo, C.; Magnocavallo, M.; Alfarano, M.; Mariani, M.V.; Bernardini, F.; Della Rocca, D.G.; Galardo, G.; Severino, P.; Di Lullo, L.; et al. A proposed strategy for anticoagulation therapy in noncompaction cardiomyopathy. *ESC Heart Fail.* **2022**, *9*, 241–250.

12-Pöyhönen, P.; Kuusisto, J.; Järvinen, V.; Pirinen, J.; Rätty, H.; Lehmonen, L.; Paakkanen, R.; Martinez-Majander, N.; Putaala, J.; Sinisalo, J. Left ventricular non-compaction as a potential source for cryptogenic ischemic stroke in the young: A case-control study. *PLoS ONE* **2020**, *15*, e0237228.

13-Sparrow, P.; Merchant, N.; Provost, Y.; Doyle, D.; Nguyen, E.; Paul, N. Cardiac MRI and CT features of inheritable and congenital conditions associated with sudden cardiac death. *Eur. Radiol.* **2009**, *19*, 259–270.

14-McDonagh, T.A.; Metra, M.; Adamo, M.; Gardner, R.S.; Baumbach, A.; Böhm, M.; Burri, H.; Butler, J.; Čelutkienė, J.; Chioncel, O.; et al. ESC SDG. 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur. Heart J.* **2021**, *42*, 3599–3726

15-Chimenti, C.; Lavallo, C.; Magnocavallo, M.; Alfarano, M.; Mariani, M.V.; Bernardini, F.; Della Rocca, D.G.; Galardo, G.; Severino, P.; Di Lullo, L.; et al. A proposed strategy for anticoagulation therapy in noncompaction cardiomyopathy. *ESC Heart Fail.* **2022**, *9*, 241–250.

16-Engberding, R.; Stöllberger, C.; Schneider, B.; Nothnagel, D.; Fehske, W.; Gerecke, B.J. Abstract 14769: Heart Failure in Noncompaction Cardiomyopathy - Data from the German Noncompaction Registry (ALKK). *Circulation* **2012**, *126*, A14769.

17-L.E. Visser, R.H. van Schaik, A.H. Jan Danser, *et al.* **The risk of myocardial infarction in patients with reduced activity of cytochrome P450 2C9** *Pharmacogenetics Genom*, 17 (7) (2007), pp. 473-479

18-Aras D., Tufekcioglu O., Ergun K., Clinical features of isolated ventricular noncompaction in adults long-term clinical course, echocardiographic properties, and predictors of left ventricular failure. *J Card Fail*. 2006; 12 (9) : 726-733 .

19- Jenni, C.A. Wyss, E.N. Oechslin, *et al.* **Isolated ventricular noncompaction is associated with coronary microcirculatory dysfunction** *J Am Coll Cardiol*, 39 (2002), pp. 450-454

22- [Jerzy Paluszkiwicz](#) and al **Left Ventricular Non-Compaction Cardiomyopathy-Still More Questions than Answers.** *J. Clin. Med.* **2022**, 11(14), 4135

21-[Toufan](#),M [Shahvalizadeh](#). R ,[Khalili](#)M . Myocardial infarction in a patient with left ventricular noncompaction: a case report *Int J Gen Med*. 2012; 5: 661–665

22-A. Gopi, P. Sebastian, C.D. Ramakrishna **A rare malady with a rarer complication** *Indian Heart J*, 68 (Suppl 2) (2016), pp. S264-S266

23- Kaul A. Hussain .A\*Vinay K Sharma. Noncompaction of left ventricle presenting as acute ST-elevation myocardial infarction in a young male with normal coronaries—a case report *Journal of Indian College of Cardiology* Volume 1, Issue 2, September 2011, Pages 115-117.




Figure 1 : apical 4-chamber echocardiographic section showing a trabeculated aspect of the apex of the left ventricle




Figure 2 : apical 4-chamber echocardiographic section showing inter trabecular recesses taking colour on colour doppler



Figure 3 : Parasternal short-axis echocardiographic section showing an aspect of non-compacted tissue at the apical level with a ratio of compacted to non-compacted tissue of 3.14